

For all the questions

- $[k] := \{1, 2, \dots, k\}$  where  $k \in \mathbb{N}$ .
- $\mathcal{L}(\mathcal{H}) :=$  Linear operators on  $\mathcal{H}$
- $\mathcal{R}(\mathcal{H}) :=$  Self-adjoint or hermitian operators on  $\mathcal{H}$
- $\mathcal{P}(\mathcal{H}) :=$  Positive semi-definite operators on  $\mathcal{H}$
- $\mathcal{D}(\mathcal{H}) :=$  Density operators on  $\mathcal{H}$

### Problem 1

For  $T : \mathcal{H} \rightarrow \mathcal{H}$ , prove that

$$\sum_{i=1}^d \langle e_i | T e_i \rangle = \sum_{i=1}^d \langle f_i | T f_i \rangle$$

if  $\{|e_i\rangle \in \mathcal{H} \mid 1 \leq i \leq d\}$  and  $\{|f_i\rangle \in \mathcal{H} \mid 1 \leq i \leq d\}$  are ONB.

**Solution:** Let  $S : \mathcal{H} \rightarrow \mathcal{H}$  where it maps the basis vectors from  $|e_i\rangle \rightarrow |f_i\rangle$ . Then  $S|e_i\rangle = |f_i\rangle$ . Hence  $S$  is an orthonormal matrix since

$$\langle e_j | S^\dagger S | e_i \rangle = \langle f_j | f_i \rangle = \delta_{ji} \quad \text{and} \quad \langle f_j | S S^\dagger | f_i \rangle = \langle e_j | e_i \rangle = \delta_{ji}$$

Hence

$$\sum_{i=1}^d \langle f_i | T f_i \rangle = \sum_{i=1}^d \langle e_i | S^\dagger T S | e_i \rangle = \text{tr}(S^\dagger T S) = \text{tr}(S S^\dagger T) = \text{tr}(T) = \sum_{i=1}^d \langle e_i | T e_i \rangle$$

Therefore we have

$$\sum_{i=1}^d \langle e_i | T e_i \rangle = \sum_{i=1}^d \langle f_i | T f_i \rangle$$

□

### Problem 2

If  $\{|e_i\rangle \in \mathcal{H}_1 \mid 1 \leq i \leq d\}$  and  $\{|f_i\rangle \in \mathcal{H}_2 \mid 1 \leq i \leq d\}$  are ONB, then  $\{|e_i\rangle \otimes |f_j\rangle \mid 1 \leq i, j \leq d\} \subseteq \mathcal{H}_1 \otimes \mathcal{H}_2$  is ONB

**Solution:** Let  $|\psi\rangle \otimes |\phi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2$ . Then  $|\psi\rangle = \sum_{i=1}^d \alpha_i |e_i\rangle$  where  $\alpha_i \in \mathbb{C}$  for all  $i \in [d]$  since  $\{|e_i\rangle \in \mathcal{H}_1 \mid 1 \leq i \leq d\}$  is ONB for  $\mathcal{H}_1$ . Hence

$$|\psi\rangle \otimes |\phi\rangle = \sum_{i=1}^d \alpha_i |e_i\rangle \otimes |\phi\rangle$$

Now  $|\phi\rangle = \sum_{i=1}^d \beta_i |f_i\rangle$  where  $\beta_i \in \mathbb{C}$  for all  $i \in [d]$  since  $\{|f_i\rangle \in \mathcal{H}_2 \mid 1 \leq i \leq d\}$  is ONB for  $\mathcal{H}_2$ . Hence

$$\forall i \in [d] \quad |e_i\rangle \otimes |\phi\rangle = \sum_{j=1}^d \beta_j |e_i\rangle \otimes |f_j\rangle$$

Therefore we get

$$|\psi\rangle \otimes |\phi\rangle = \sum_{i=1}^d \alpha_i |e_i\rangle \otimes |\phi\rangle = \sum_{i=1}^d \alpha_i \sum_{j=1}^d \beta_j |e_i\rangle \otimes |f_j\rangle = \sum_{1 \leq i,j \leq d} \alpha_i \beta_j |e_i\rangle \otimes |f_j\rangle$$

Therefore  $\{|e_i\rangle \otimes |f_j\rangle \mid 1 \leq i, j \leq d\}$  is a basis of  $\mathcal{H}_1 \otimes \mathcal{H}_2$ .

Now for any  $i_1, i_2, j_1, j_2 \in [d]$

$$(\langle e_{i_1} | \otimes \langle f_{j_1} |)(|e_{i_2}\rangle \otimes |f_{j_2}\rangle) = \langle e_{i_1} | e_{i_2} \rangle \langle f_{j_1} | f_{j_2} \rangle = \delta_{i_1, i_2} \delta_{j_1, j_2}$$

Therefore  $\{|e_i\rangle \otimes |f_j\rangle \mid 1 \leq i, j \leq d\}$  is orthonormal. Therefore  $\{|e_i\rangle \otimes |f_j\rangle \mid 1 \leq i, j \leq d\}$  is a ONB for  $\mathcal{H}_1 \otimes \mathcal{H}_2$ . □

### Problem 3

Let  $\{|g_k\rangle \mid 1 \leq k \leq d_2\} \subseteq \mathcal{H}_2$  be ONB. For  $T \in \mathcal{L}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ , let  $\text{tr}_2(T) \in \mathcal{L}(\mathcal{H}_1)$  denote the operator satisfying

$$\langle u | \text{tr}_2(T) | v \rangle = \sum_k \langle u \otimes g_k | T | v \otimes g_k \rangle$$

for any choice  $|u\rangle, |v\rangle \in \mathcal{H}_1$ . Prove that  $\sum_k \langle u \otimes g_k | T | v \otimes g_k \rangle$  is invariant.

### Problem 4

Show that the Pauli matrices are all Hermitian, unitary, they square to the identity, and their eigenvalues are  $\pm 1$

### Problem 5 Mark Wilde: Exercise 3.3.3

For  $S, T \in \mathcal{L}(\mathcal{H})$ , show that

$$\text{tr}(T) = \text{tr}(T^+), \quad \text{tr}(ST) = \text{tr}(TS)$$

[Recall  $T^+$  denotes adjoint of  $T$ ]. For  $|x\rangle, |y\rangle \in \mathcal{H}$  show

$$\text{tr}(|x\rangle \langle y| T) = \text{tr}(T |x\rangle \langle y|) = \langle y | T x \rangle$$

### Problem 6

Suppose  $\mathcal{H}$  is finite dimensional complex inner product space with  $\dim(\mathcal{H}) = d$ . Show complex dimensionality of  $\mathcal{L}(\mathcal{H})$  is  $d^2$ , real dimensionality of  $\mathcal{R}(\mathcal{H})$  is  $d^2$ .

Suppose  $\mathcal{H}$  is a real inner product space of dim  $d$ , show  $\mathcal{L}(\mathcal{H})$  has dimension  $d$  and the space of all symmetric operators is a real vector space of dimension  $\frac{d(d+1)}{2}$ .

**Solution:** Suppose  $\{|e_i\rangle \in \mathcal{H} \mid 1 \leq i \leq d\}$  is an ONB of  $\mathcal{H}$ . Let  $T \in \mathcal{L}(\mathcal{H})$ . Then for all  $i \in [d]$

$$T |e_i\rangle = \sum_{j=1}^d \alpha_{i,j} |e_j\rangle$$

where  $\alpha_{i,j} \in \mathbb{C}$ . Hence the map  $T$  is uniquely decided by the numbers  $\alpha_{i,j} \in \mathbb{C}$  for all  $i, j \in [d]$ . Hence there are  $d^2$  many numbers which uniquely decides  $T$ . Therefore  $\dim(\mathcal{L}(\mathcal{H})) = d^2$ .

Now let  $T \in \mathcal{R}(\mathcal{H})$ . Then  $T^\dagger = T$ . Again suppose  $\{|e_i\rangle \in \mathcal{H} \mid 1 \leq i \leq d\}$  is an ONB of  $\mathcal{H}$ . Let  $(i, j)$ th element of  $T$  is denoted by  $t_{i,j}$ . Then for all  $i \in [d]$ ,  $T_{i,i} \in \mathbb{R}$  since  $T^\dagger = T$ . Now for all off diagonal entries  $t_{j,i} = t_{i,j}^*$ .

So there are  $\frac{n^2-n}{2}$  many complex numbers which decides  $T$  uniquely apart from the  $n$  real entries in the diagonal. Now for each  $i, j \in [d]$  let  $t_{i,j} = x_{i,j} + iy_{i,j}$  where  $x_{i,j}, y_{i,j} \in \mathbb{R}$ . Therefore

$$t_{j,i} = t_{i,j}^* = x_{i,j} - iy_{i,j}$$

So for each off-diagonal entries there are corresponding 2 real numbers. And there are total  $\frac{n^2-n}{2}$  many off-diagonal entries which participates in uniquely deciding  $T$ . Hence there are total  $2 \times \frac{n^2-n}{2} + n = n^2$  real numbers which uniquely decides  $T$ . Hence  $\dim(\mathcal{H}) = d^2$ .

□

### Problem 7

Show that  $\mathcal{D}(\mathcal{H})$  is a convex subset of the real vector space of all Hermitian operators on  $\mathcal{H}$ . Show that the extreme points of  $\mathcal{D}(\mathcal{H})$  are pure states, i.e. rank 1 projection operators.

### Problem 8

Show that if  $\dim(\mathcal{H}) = d$ , then  $\mathcal{D}(\mathcal{H})$  can be embedded into a real vector space of dimension  $n = d^2 - 1$

### Problem 9

Prove the Singular value decomposition theorem stated in class.

### Problem 10

Suppose  $|\psi\rangle_{AR_1} \in \mathcal{H}_A \otimes \mathcal{H}_{R_1}$ ,  $|\psi\rangle_{AR_2} \in \mathcal{H}_A \otimes \mathcal{H}_{R_2}$  are purifications of  $\rho_A \in \mathcal{D}(\mathcal{H}_A)$  and  $\dim(\mathcal{H}_{R_2}) \geq \dim(\mathcal{H}_{R_1})$ , then show that there exists an isometry  $V : \mathcal{H}_{R_1} \rightarrow \mathcal{H}_{R_2}$  such that

$$|\psi\rangle_{AR_2} = (V \otimes I) |\psi\rangle_{AR_1}$$

### Problem 11 Mark Wilde: Exercise 3.6.5

Show that the Bell states form an orthonormal basis:

$$\langle \Phi^{z_1 x_1} | \Phi^{z_2 x_2} \rangle = \delta_{z_1, z_2} \delta_{x_1, x_2}$$

### Problem 12 Mark Wilde: Exercise 3.7.11

Show that the set of states  $\{|\Phi^{x,z}\rangle_{AB}\}_{x,z=0}^{d-1}$  forms a complete, orthonormal basis:

$$\langle \Phi^{x_1, z_1} | \Phi^{x_2, z_2} \rangle = \delta_{x_1, x_2} \delta_{z_1, z_2} \quad \sum_{x,z=0}^d |\Phi^{x,z}\rangle \langle \Phi^{x,z}| = I_{AB}$$

### Problem 13 Mark Wilde: Exercise 4.1.5

Show that the following ensembles have the same density operator:  $\{\{\frac{1}{2}, |0\rangle\}, \{\frac{1}{2}, |1\rangle\}\}$  and  $\{\{\frac{1}{2}, |+\rangle\}, \{\frac{1}{2}, |-\rangle\}\}$

### Problem 14

Show that the set of states  $\{|\Phi^{x,z}\rangle_{AB}\}_{x,z=0}^{d-1}$  forms a complete, orthonormal basis:

$$\langle \Phi^{x_1, z_1} | \Phi^{x_2, z_2} \rangle = \delta_{x_1, x_2} \delta_{z_1, z_2} \quad \sum_{x,z=0}^d |\Phi^{x,z}\rangle \langle \Phi^{x,z}| = I_{AB}$$

**Problem 15** Mark Wilde: Exercise 4.1.3

Show that the following ensembles have the same density operator:  $\{\{\frac{1}{2}, |0\rangle\}, \{\frac{1}{2}, |1\rangle\}\}$  and  $\{\{\frac{1}{2}, |+\rangle\}, \{\frac{1}{2}, |-\rangle\}\}$

**Problem 16** Mark Wilde: Exercise 3.7.12

Show that the following ensembles have the same density operator:  $\{\{\frac{1}{2}, |0\rangle\}, \{\frac{1}{2}, |1\rangle\}\}$  and  $\{\{\frac{1}{2}, |+\rangle\}, \{\frac{1}{2}, |-\rangle\}\}$

**Problem 17**

Show that the following ensembles have the same density operator:  $\{\{\frac{1}{2}, |0\rangle\}, \{\frac{1}{2}, |1\rangle\}\}$  and  $\{\{\frac{1}{2}, |+\rangle\}, \{\frac{1}{2}, |-\rangle\}\}$

**Problem 18**

Show that the following ensembles have the same density operator:  $\{\{\frac{1}{2}, |0\rangle\}, \{\frac{1}{2}, |1\rangle\}\}$  and  $\{\{\frac{1}{2}, |+\rangle\}, \{\frac{1}{2}, |-\rangle\}\}$

**Problem 19**

Show that the following ensembles have the same density operator:  $\{\{\frac{1}{2}, |0\rangle\}, \{\frac{1}{2}, |1\rangle\}\}$  and  $\{\{\frac{1}{2}, |+\rangle\}, \{\frac{1}{2}, |-\rangle\}\}$