

# Super-Polynomial Lower Bound of TSP Extended Formula

Soham Chatterjee

May 2025

# Introduction

## Definition (Travelling Salesman)

Given a graph  $G = (V, E)$ ,  $S \subseteq V$  and weights  $w : E \rightarrow \mathbb{R}$  find minimum weight cycle which visits every vertex of  $S$  exactly once.

# Introduction

## Definition (Travelling Salesman)

Given a graph  $G = (V, E)$ ,  $S \subseteq V$  and weights  $w : E \rightarrow \mathbb{R}$  find minimum weight cycle which visits every vertex of  $S$  exactly once.

We will focus on  $S = V$ .

- We know Traveling Salesman Problem is NP-complete.
- In [Yannakakis, 1988, STOC] he proved every symmetric LP for the TSP has exponential size.
- Here we will show TSP admits no polynomial-size LP.
- This proof also shows unconditional super-polynomial lower bound on the number of inequalities.
- Therefore it is impossible to prove  $P = NP$  by means of a polynomial size LP.

## Definitions

Let  $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$  is a polytope with  $A \in \mathbb{R}^{m \times d}$ ,  $b \in \mathbb{R}^m$  and  $V \subseteq \mathbb{R}^d$ . We will consider  $V$  as the characteristic vector for all hamiltonian paths.

# Definitions

Let  $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$  is a polytope with  $A \in \mathbb{R}^{m \times d}$ ,  $b \in \mathbb{R}^m$  and  $V \subseteq \mathbb{R}^d$ . We will consider  $V$  as the characteristic vector for all hamiltonian paths.

## Definition (Extension Polytope)

An extension of  $P$  is a polytope  $Q \subseteq \mathbb{R}^{d+e}$  such that there is a linear map  $\pi : \mathbb{R}^{d+e} \rightarrow \mathbb{R}^d$  such that  $\pi(Q) = P$ .

# Definitions

Let  $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$  is a polytope with  $A \in \mathbb{R}^{m \times d}$ ,  $b \in \mathbb{R}^m$  and  $V \subseteq \mathbb{R}^d$ . We will consider  $V$  as the characteristic vector for all hamiltonian paths.

## Definition (Extension Polytope)

An extension of  $P$  is a polytope  $Q \subseteq \mathbb{R}^{d+e}$  such that there is a linear map  $\pi : \mathbb{R}^{d+e} \rightarrow \mathbb{R}^d$  such that  $\pi(Q) = P$ .

## Definition (Extended Formula)

An EF  $Q$  is an extension of  $P$  is a linear system in variable  $s(x,y)$  such that

$$x \in P \iff \exists y (x,y) \in Q$$

# Definitions

Let  $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$  is a polytope with  $A \in \mathbb{R}^{m \times d}$ ,  $b \in \mathbb{R}^m$  and  $V \subseteq \mathbb{R}^d$ . We will consider  $V$  as the characteristic vector for all hamiltonian paths.

## Definition (Extension Polytope)

An extension of  $P$  is a polytope  $Q \subseteq \mathbb{R}^{d+e}$  such that there is a linear map  $\pi : \mathbb{R}^{d+e} \rightarrow \mathbb{R}^d$  such that  $\pi(Q) = P$ .

## Definition (Extended Formula)

An EF  $Q$  is an extension of  $P$  is a linear system in variable  $s(x,y)$  such that

$$x \in P \iff \exists y (x,y) \in Q$$

Extension complexity of  $P$  is the minimum size EF of  $P$  where size of a polytope is the number inequalities. We denote by  $xc(P)$ .

## Some Polytopes

- $TSP(n)$  is the traveling salesman polytope for  $K_n = (V_n, E_n)$ . Let  $C \subseteq E_n$  denotes a tour of  $K_n$ . Then  $\chi^C$  denotes the characteristic vector of  $C$ . Then



## Some Polytopes

- $TSP(n)$  is the traveling salesman polytope for  $K_n = (V_n, E_n)$ . Let  $C \subseteq E_n$  denotes a tour of  $K_n$ . Then  $\chi^C$  denotes the characteristic vector of  $C$ . Then

$$TSP(n) := \text{conv}\{\chi^C \mid C \subseteq E_n \text{ is a tour of } K_n\}$$

## Some Polytopes

- $TSP(n)$  is the traveling salesman polytope for  $K_n = (V_n, E_n)$ . Let  $C \subseteq E_n$  denotes a tour of  $K_n$ . Then  $\chi^C$  denotes the characteristic vector of  $C$ . Then

$$TSP(n) := \text{conv}\{\chi^C \mid C \subseteq E_n \text{ is a tour of } K_n\}$$

- Given  $G = (V, E)$ , for any  $S \subseteq V$ ,  $\chi^S$  denote characteristic vector of  $S$ . Then

## Some Polytopes

- $TSP(n)$  is the traveling salesman polytope for  $K_n = (V_n, E_n)$ . Let  $C \subseteq E_n$  denotes a tour of  $K_n$ . Then  $\chi^C$  denotes the characteristic vector of  $C$ . Then

$$TSP(n) := \text{conv}\{\chi^C \mid C \subseteq E_n \text{ is a tour of } K_n\}$$

- Given  $G = (V, E)$ , for any  $S \subseteq V$ ,  $\chi^S$  denote characteristic vector of  $S$ . Then

$$IND(G) := \text{conv}\{\chi^S \mid S \text{ is independent set of } G\}$$

- The correlation polytope  $COR(n)$  is

$$COR(n) := \text{conv}\{bb^T \mid b \in \{0, 1\}^n\}$$