

Super-Polynomial Lower Bound of TSP Extended Formula

Soham Chatterjee

May 2025



Introduction

Introduction

Definition (Travelling Salesman)

Given a graph $G = (V, E)$, $S \subseteq V$ and weights $w : E \rightarrow \mathbb{R}$ find minimum weight cycle which visits every vertex of S exactly once.

Introduction

Definition (Travelling Salesman)

Given a graph $G = (V, E)$, $S \subseteq V$ and weights $w : E \rightarrow \mathbb{R}$ find minimum weight cycle which visits every vertex of S exactly once.

We will focus on $S = V$.

- We know Traveling Salesman Problem is NP-complete.
- In [Yannakakis, 1988, STOC] he proved every symmetric LP for the TSP has exponential size.
- Here we will show TSP admits no polynomial-size LP.
- This proof also shows unconditional super-polynomial lower bound on the number of inequalities.
- Therefore it is impossible to prove $P = NP$ by means of a polynomial size LP.

Definitions

Let $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$ is a polytope with $A \in \mathbb{R}^{m \times d}$, $b \in \mathbb{R}^m$ and $V \subseteq \mathbb{R}^d$. We will consider V as the characteristic vector for all hamiltonian paths.

Definitions

Let $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$ is a polytope with $A \in \mathbb{R}^{m \times d}$, $b \in \mathbb{R}^m$ and $V \subseteq \mathbb{R}^d$. We will consider V as the characteristic vector for all hamiltonian paths.

Definition (Extension Polytope)

An extension of P is a polytope $Q \subseteq \mathbb{R}^{d+e}$ such that there is a linear map $\pi : \mathbb{R}^{d+e} \rightarrow \mathbb{R}^d$ such that $\pi(Q) = P$.

Definitions

Let $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$ is a polytope with $A \in \mathbb{R}^{m \times d}$, $b \in \mathbb{R}^m$ and $V \subseteq \mathbb{R}^d$. We will consider V as the characteristic vector for all hamiltonian paths.

Definition (Extension Polytope)

An extension of P is a polytope $Q \subseteq \mathbb{R}^{d+e}$ such that there is a linear map $\pi : \mathbb{R}^{d+e} \rightarrow \mathbb{R}^d$ such that $\pi(Q) = P$.

Definition (Extended Formula)

An EF Q is an extension of P is a linear system in variable $s(x,y)$ such that

$$x \in P \iff \exists y (x,y) \in Q$$

Definitions

Let $P = \{x \in \mathbb{R}^n \mid Ax \leq b\} = \text{conv}(V)$ is a polytope with $A \in \mathbb{R}^{m \times d}$, $b \in \mathbb{R}^m$ and $V \subseteq \mathbb{R}^d$. We will consider V as the characteristic vector for all hamiltonian paths.

Definition (Extension Polytope)

An extension of P is a polytope $Q \subseteq \mathbb{R}^{d+e}$ such that there is a linear map $\pi : \mathbb{R}^{d+e} \rightarrow \mathbb{R}^d$ such that $\pi(Q) = P$.

Definition (Extended Formula)

An EF Q is an extension of P is a linear system in variable $s(x,y)$ such that

$$x \in P \iff \exists y (x,y) \in Q$$

Extension Complexity

Extension complexity of P is the minimum size EF of P where size of a polytope is the number inequalities. We denote by $xc(P)$.

Extension Complexity

Extension complexity of P is the minimum size EF of P where size of a polytope is the number inequalities. We denote by $xc(P)$.

Lemma

Let P, Q and F be polytopes. Then the following holds:

- (i) If F is an extension of P then $xc(F) \geq xc(P)$.*
- (ii) If F is a face of Q then $xc(Q) \geq xc(F)$.*

Some Polytopes

- $TSP(n)$ is the traveling salesman polytope for $K_n = (V_n, E_n)$. Let $C \subseteq E_n$ denotes a tour of K_n . Then χ^C denotes the characteristic vector of C . Then

Some Polytopes

- $TSP(n)$ is the traveling salesman polytope for $K_n = (V_n, E_n)$. Let $C \subseteq E_n$ denotes a tour of K_n . Then χ^C denotes the characteristic vector of C . Then

$$TSP(n) := \text{conv}\{\chi^C \mid C \subseteq E_n \text{ is a tour of } K_n\}$$

Some Polytopes

- $TSP(n)$ is the traveling salesman polytope for $K_n = (V_n, E_n)$. Let $C \subseteq E_n$ denotes a tour of K_n . Then χ^C denotes the characteristic vector of C . Then

$$TSP(n) := \text{conv}\{\chi^C \mid C \subseteq E_n \text{ is a tour of } K_n\}$$

- Given $G = (V, E)$, for any $S \subseteq V$, χ^S denote characteristic vector of S . Then

Some Polytopes

- $TSP(n)$ is the traveling salesman polytope for $K_n = (V_n, E_n)$. Let $C \subseteq E_n$ denotes a tour of K_n . Then χ^C denotes the characteristic vector of C . Then

$$TSP(n) := \text{conv}\{\chi^C \mid C \subseteq E_n \text{ is a tour of } K_n\}$$

- Given $G = (V, E)$, for any $S \subseteq V$, χ^S denote characteristic vector of S . Then

$$IND(G) := \text{conv}\{\chi^S \mid S \text{ is independent set of } G\}$$

- The correlation polytope $COR(n)$ is

$$COR(n) := \text{conv}\{bb^T \mid b \in \{0, 1\}^n\}$$

Proof Flow

Step 1: First we will prove $xc(COR(n)) = 2^{\Omega(n)}$

Proof Flow

Step 1: First we will prove $xc(COR(n)) = 2^{\Omega(n)}$

Step 2: For all n , \exists graph G_n with n vertices such that $xc(IND(G_n)) \geq xc(COR(n'))$
where $n' = n^{\frac{1}{d}}$ for some $d > 1$.

Proof Flow

Step 1: First we will prove $xc(COR(n)) = 2^{\Omega(n)}$

Step 2: For all n , \exists graph G_n with n vertices such that $xc(IND(G_n)) \geq xc(COR(n'))$
where $n' = n^{\frac{1}{d}}$ for some $d > 1$.

Step 3: For any n -vertex graph G , $IND(G)$ is linear projection of a face of $TSP(k)$
where $k = O(n^2)$.

Correlation Polytope Lower Bound