No.	Medicine Name	Dose
1	Amoxicillin	500 mg
2	Ampicillin	500 mg–2 g
3	Penicillin V	250–500 mg
4	Benzathine penicillin G	1.2–2.4 million units
5	Flucloxacillin	500 mg
6	Piperacillin + tazobactam	4.5 g
7	Cefazolin	1–2 g
8	Cefuroxime	250–500 mg
9	Cefixime	200–400 mg/day
10	Cefpodoxime	100–200 mg
11	Ceftriaxone	1–2 g
12	Cefotaxime	1–2 g
13	Ceftazidime	1–2 g
14	Cefepime	1–2 g
15	Cefoperazone + sulbactam	1.5–3 g
16	Meropenem	1 g
17	Imipenem + cilastatin	500 mg
18	Ertapenem	1 g
19	Aztreonam	1–2 g
20	Azithromycin	500 mg day 1 then 250 mg, or 500 mg daily
21	Clarithromycin	250–500 mg
22	Ciprofloxacin	500–750 mg
23	Levofloxacin	500–750 mg
24	Moxifloxacin	400 mg
25	Ofloxacin	200–400 mg
26	Gentamicin	5–7 mg/kg
27	Amikacin	15 mg/kg
28	Tobramycin	5–7 mg/kg
29	Streptomycin	15 mg/kg (max 1 g)
30	Vancomycin	15–20 mg/kg
31	Teicoplanin	6–12 mg/kg
32	Linezolid	600 mg
33	Daptomycin	4–6 mg/kg
34	Colistin	Weight-based loading and maintenance
35	Polymyxin B	1.5–2.5 mg/kg/day
36	Tigecycline	100 mg then 50 mg
37	Co-trimoxazole (trimethoprim + sulfamethox	160/800 mg
38	Nitrofurantoin	50–100 mg
39	Fosfomycin	3 g
40	Metronidazole oral	400–500 mg
41	Tinidazole	2 g single dose or 500 mg twice daily
42	Secnidazole	2 g
43	Nitazoxanide	500 mg
44	Chloramphenicol	50 mg/kg/day
45	Isoniazid	5 mg/kg (usual 300 mg)
46	Pyrazinamide	20–25 mg/kg
47	Rifapentine	600–1200 mg
48	Bedaquiline	400 mg daily for 2 weeks then 200 mg three t

49	Pretomanid	200 mg
50	Delamanid	100 mg
51	Ethionamide	15–20 mg/kg/day
52	Cycloserine	10–15 mg/kg/day
53	Voriconazole	6 mg/kg IV twice on day 1 then 4 mg/kg; oral 2
54	Posaconazole	300 mg twice day 1 then 300 mg
55	Isavuconazole	372 mg isavuconazonium sulfate
56	Caspofungin	70 mg then 50 mg
57	Micafungin	100–150 mg
58	Anidulafungin	200 mg then 100 mg
59	Flucytosine	100–150 mg/kg/day
60	Oseltamivir	75 mg
61	Baloxavir marboxil	40–80 mg (weight based)
62	Ganciclovir	5 mg/kg
63	Valganciclovir	900 mg
64	Remdesivir	200 mg then 100 mg
65	Nirmatrelvir + ritonavir	300 mg + 100 mg
66	Molnupiravir	800 mg
67	Sofosbuvir	400 mg
68	Sofosbuvir + velpatasvir	400 mg + 100 mg
69	Entecavir	0.5 mg
70	Tenofovir disoproxil fumarate	300 mg
71	Tenofovir alafenamide	25 mg
72	Lamivudine	150 mg twice daily or 300 mg
73	Zidovudine	300 mg
74	Abacavir	600 mg
75	Dolutegravir	50 mg
76	Efavirenz	600 mg
77	Lopinavir + ritonavir	400 mg + 100 mg
78	Atazanavir + ritonavir	300 mg + 100 mg
79	Darunavir + ritonavir	800 mg + 100 mg
80	Raltegravir	400 mg
81	Tenofovir + lamivudine + dolutegravir (TLD)	300 mg + 300 mg + 50 mg
82	Emtricitabine + tenofovir	200 mg + 300 mg
83	Ribavirin	Weight-based
84	Artemether + lumefantrine	20/120 mg tablets, weight-based (4 tablets fo
85	Artesunate	2.4 mg/kg
86	Chloroquine	600 mg base then 300 mg
87	Primaquine	0.25–0.5 mg/kg
88	Tafenoquine	300 mg
89	Atovaquone + proguanil	250 mg + 100 mg
90	Mefloquine	250 mg
91	Quinine	600 mg
92	Sulfadoxine + pyrimethamine	Weight-based
93	Mebendazole	100 mg twice daily for 3 days or 500 mg singl
94	Praziquantel	40–60 mg/kg
95	Pyrantel pamoate	11 mg/kg (max 1 g)
96	Paromomycin	25–35 mg/kg/day

Frequency
3 times daily
Every 6 hours
4 times daily
IM single dose or weekly per indicatic
4 times daily
IV every 6–8 hours
IV every 8 hours
Twice daily
Once/twice daily
Twice daily
IV/IM once daily
IV every 6–8 hours
IV every 8 hours
IV every 8–12 hours
IV every 12 hours
IV every 8 hours
IV every 6 hours
IV once daily
IV every 8 hours
Once daily
Twice daily
Twice daily
Once daily
Once daily
Twice daily
IV/IM once daily
IV/IM once daily
IV once daily
IM once daily
IV every 8–12 hours
IV/IM every 12 hours for loading, ther
Twice daily
IV once daily
IV in divided doses
IV in divided doses
IV every 12 hours
Twice daily
4 times daily or twice daily (MR)
Single dose
3 times daily
Single/twice daily
Single dose
Twice daily for 3 days
Divided every 6 hours
Once daily
Once daily
Weekly or daily per regimen
Per regimen

Once daily
Twice daily
Divided doses
Divided doses
Twice daily
Once daily
Every 8 hours for 6 doses, then once d
IV once daily
IV once daily
IV once daily
Divided every 6 hours
Twice daily for 5 days
Single dose
IV every 12 hours
Twice daily (induction) then once dai
IV once daily
Twice daily for 5 days
Twice daily for 5 days
Once daily
Once daily
Once daily
Once daily
Once daily
Once/twice daily
Twice daily
Once daily
Once daily
Once daily at bedtime
Twice daily
Once daily
Once daily with food
Twice daily
Once daily
Once daily
Divided twice daily
Twice daily for 3 days
IV at 0, 12, 24 hours then daily
Per regimen over 3 days
Once daily for 7–14 days
Single dose
Once daily
Weekly for prophylaxis
3 times daily
Single dose
Per indication
Single/divided doses
Single dose
Divided doses

Instructions
Complete prescribed course; check penicillin allergy
Oral or IV depending on indication; check penicillin allergy
Take on an empty stomach; complete course
Deep IM injection only; check allergy history
Take on an empty stomach; monitor liver function with prolonged use
Hospital use; adjust for renal function
Surgical prophylaxis and skin/soft-tissue infections
Take with food; complete course
Complete prescribed course
Take with food; complete course
Hospital/clinic use; do not mix with calcium-containing IV fluids in neonates
Hospital use; adjust for renal function
Hospital use; useful for Pseudomonas coverage
Hospital use; adjust for renal function; neurotoxicity risk in renal impairment
Hospital use; monitor for bleeding tendency
Hospital use; adjust for renal function; interacts with valproate
Hospital use; seizure risk in renal impairment
Hospital/outpatient infusion; adjust for renal function
Hospital use; option in severe beta-lactam allergy
Take as directed; check QT-prolonging interactions
Many drug interactions; check QT-prolonging drugs
Avoid dairy/antacids near dose; tendon and QT warnings
Tendon, QT and glucose warnings; hydrate well
QT prolongation warning; avoid with antacids
Tendon and CNS warnings; separate from antacids
Hospital use; monitor renal function and levels
Hospital use; monitor renal function and hearing
Hospital use; monitor renal function and levels
Specialist TB use; monitor hearing and renal function
Hospital use; monitor levels and renal function
Hospital use; monitor renal function
Monitor blood counts; avoid serotonergic drug combinations
Hospital use; monitor CK; not for pneumonia
Specialist hospital use; monitor renal function
Specialist hospital use; nephrotoxicity and neurotoxicity risk
Hospital use; nausea common; not first-line for bloodstream infection
Hydrate well; check sulfonamide allergy and potassium
Take with food; avoid if poor kidney function
Dissolve in water; uncomplicated UTI
Avoid alcohol during and 48 hours after
Take with food; avoid alcohol
Take with food; avoid alcohol
Take with food
Specialist use; monitor blood counts
Give with pyridoxine; monitor liver function
Monitor liver function and uric acid
Specialist TB use; many drug interactions
DR-TB specialist use; monitor QT

Specialist DR-TB regimen; take with food
Specialist DR-TB use; monitor QT
Take with food; monitor liver and thyroid function
Specialist DR-TB use; monitor mood/neurologic effects
Check interactions; monitor liver function and visual symptoms
Delayed-release tablet with food; check interactions
Specialist use; monitor liver function
Hospital use; adjust in liver impairment
Hospital use for candidemia
Hospital use for candidemia
Specialist use; monitor blood counts and renal function
Start within 48 hours; adjust in renal impairment
Influenza; avoid with polyvalent cation products
Hospital use; monitor blood counts and renal function
Take with food; monitor blood counts
Hospital use; monitor liver function
Many drug interactions; adjust in renal impairment
Avoid in pregnancy; start early
Hepatitis C; used within combination regimen
Hepatitis C; check hepatitis B status
Take on an empty stomach; monitor hepatitis B response
Monitor renal function and bone health
Hepatitis B; take with food
HIV/HBV use; adjust for renal function
Monitor blood counts; HIV regimen
HLA-B*5701 testing before use
Separate from polyvalent cations; HIV regimen
Take on empty stomach; CNS effects common
Many drug interactions; GI upset common
Take with food; check interactions
Check sulfonamide allergy and interactions
Separate from antacids containing aluminium/magnesium
First-line HIV regimen; adherence essential
HIV treatment/PrEP; monitor renal function
Teratogenic; monitor hemoglobin
Take with fatty food; treatment of uncomplicated falciparum malaria
Severe malaria; hospital use
Vivax malaria in sensitive regions; take with food
Test G6PD first; take with food
Test G6PD first; avoid in pregnancy
Take with fatty food; malaria prophylaxis/treatment
Avoid in psychiatric/seizure history
Cinchonism common; monitor glucose and QT
Used in combination regimens; check sulfonamide allergy
Chew or swallow with food
Take with food; indication-specific regimen
Roundworm/pinworm; repeat in 2 weeks if needed
Intestinal amoebiasis/leishmaniasis per regimen