No.	Medicine Name	Dose
1	Metformin	500–2000 mg/day
2	Gliclazide	40–320 mg (MR 30–120 mg)
3	Glimepiride	1–6 mg
4	Glipizide	2.5–20 mg
5	Glibenclamide (glyburide)	2.5–15 mg
6	Pioglitazone	15–45 mg
7	Sitagliptin	100 mg
8	Vildagliptin	50 mg
9	Linagliptin	5 mg
10	Saxagliptin	2.5–5 mg
11	Teneligliptin	20–40 mg
12	Empagliflozin	10–25 mg
13	Dapagliflozin	10 mg
14	Canagliflozin	100–300 mg
15	Semaglutide injection	0.25 mg increasing to 1–2 mg
16	Semaglutide oral	3 mg then 7–14 mg
17	Dulaglutide	0.75–4.5 mg
18	Liraglutide	0.6–1.8 mg
19	Tirzepatide	2.5 mg increasing to 5–15 mg
20	Acarbose	25–100 mg
21	Voglibose	0.2–0.3 mg
22	Repaglinide	0.5–4 mg
23	Insulin regular (human)	Individualised
24	Insulin NPH (isophane)	Individualised
25	Insulin glargine	Individualised
26	Insulin detemir	Individualised
27	Insulin degludec	Individualised
28	Insulin aspart	Individualised
29	Insulin lispro	Individualised
30	Insulin glulisine	Individualised
31	Premixed insulin 70/30	Individualised
32	Levothyroxine	1.6 mcg/kg/day (adjust to TSH)
33	Liothyronine	5–25 mcg
34	Carbimazole	5–60 mg/day
35	Methimazole	5–30 mg
36	Propylthiouracil	100–300 mg/day
37	Hydrocortisone oral	15–25 mg/day
38	Prednisone	5–60 mg
39	Methylprednisolone	4–48 mg oral; 500–1000 mg IV pulse
40	Dexamethasone	0.5–10 mg
41	Fludrocortisone	0.05–0.2 mg
42	Deflazacort	6–90 mg
43	Alendronate	70 mg
44	Risedronate	35 mg weekly or 5 mg daily
45	Ibandronate	150 mg
46	Zoledronic acid	5 mg
47	Denosumab	60 mg
48	Raloxifene	60 mg

49	Teriparatide	20 mcg
50	Calcitriol (oral)	0.25–0.5 mcg
51	Cinacalcet	30–180 mg
52	Cabergoline	0.25–1 mg
53	Bromocriptine	2.5–15 mg/day
54	Desmopressin	0.1–0.4 mg/day orally or 10–40 mcg intranas
55	Octreotide	100–500 mcg SC or 20–30 mg LAR
56	Testosterone (cypionate/enanthate)	50–100 mg
57	Orlistat	120 mg

Frequency
Once/divided with meals
Once/twice daily
Once daily with breakfast
Once/twice daily
Once/twice daily
Once daily
Once daily
Once/twice daily
Once daily
Once daily
Once daily
Once daily
Once daily
Once daily
Once weekly
Once daily
Once weekly
Once daily
Once weekly
3 times daily with meals
3 times daily before meals
Before meals up to 4 times daily
30 minutes before meals
Once/twice daily
Once daily
Once/twice daily
Once daily
With meals
With meals
With meals
Twice daily with meals
Once daily
Once–three times daily
Once/divided
Once daily
Divided 2–3 times daily
Divided 2–3 times daily
Once daily in the morning
Once daily or pulse
Once/divided
Once daily
Once daily
Once weekly
Weekly/daily
Once monthly
IV once yearly
Subcutaneous every 6 months
Once daily

Subcutaneous daily
Once/twice daily
Once daily with food
Once/twice weekly
Divided doses
Once/divided
Three times daily or every 4 weeks
Once weekly or every 2 weeks IM
3 times daily with meals

Instructions
Take with food; check renal function; hold around contrast procedures
Take with meals; hypoglycemia risk
Hypoglycemia risk; adjust in renal impairment
Take 30 minutes before meals; hypoglycemia risk
Higher hypoglycemia risk; avoid in older adults/renal impairment
Monitor for edema and heart failure; bone fracture risk
Adjust in renal impairment; report severe abdominal pain
Monitor liver enzymes
No renal adjustment needed; report severe abdominal pain
Adjust in renal impairment; monitor for heart failure
Monitor glucose; adjust per local guidance
Genital hygiene; hold during acute illness/dehydration; ketoacidosis risk
Also used in heart failure/CKD; hold during acute illness
Foot care advised; ketoacidosis and dehydration risk
Subcutaneous injection; titrate slowly; GI side effects common
Take on empty stomach with small amount of water; wait 30 minutes before eating
Subcutaneous injection; GI side effects common
Subcutaneous injection; titrate slowly
Subcutaneous injection; titrate every 4 weeks
Take with first bite of food; flatulence common
Take just before meals; GI effects possible
Skip dose if meal skipped; hypoglycemia risk
Monitor glucose; rotate injection sites
Monitor glucose; resuspend before use
Basal insulin; monitor glucose; do not mix
Basal insulin; monitor glucose
Basal insulin; flexible dosing time
Rapid-acting; inject at mealtime
Rapid-acting; inject at mealtime
Rapid-acting; inject at mealtime
Resuspend before use; consistent meal timing
Take on empty stomach 30–60 minutes before breakfast; separate from calcium/iron
Specialist use; monitor heart rate and TSH
Report sore throat/fever promptly; monitor counts and liver
Report sore throat/fever; monitor liver and counts
Reserved for specific situations; liver toxicity risk
Adrenal insufficiency replacement; sick-day rules
Take with food; taper if prolonged use
Take with food; monitor glucose and BP
Potent steroid; short-course use; monitor glucose
Monitor BP and potassium
Take with food; monitor for steroid adverse effects
Take upright with full glass of water; remain upright for 30 minutes
Take upright with water; remain upright
Take upright with water; remain upright for 1 hour
Hydrate; check calcium, vitamin D and renal function
Do not delay doses; ensure calcium and vitamin D
Venous thromboembolism warning

Osteoporosis specialist use; check calcium
Monitor calcium and phosphate
Monitor calcium; nausea common
Take with food; monitor for valvulopathy at high doses
Take with food; orthostatic hypotension possible
Monitor sodium; limit fluid intake
Specialist use; monitor glucose and gallbladder
Monitor hematocrit, PSA and lipids
Take with fat-containing meal; vitamin supplementation may be needed