

The diet, prostate inflammation, and the development of prostate cancer

William G. Nelson, Theodore L. DeWeese and Angelo M. DeMarzo
The Sidney Kimmel Cancer Center at Johns Hopkins, Baltimore, MD, USA

Key words: prostate cancer, glutathione *S*-transferases, proliferative inflammatory atrophy, oxidative stress

Summary

Evidence that somatic inactivation of *GSTP1*, encoding the human π -class glutathione *S*-transferase, may initiate prostatic carcinogenesis is reviewed along with epidemiological evidence implicating several environment and lifestyle factors, including the diet and sexually transmitted diseases, as prostate cancer risk factors. An integrated model is presented featuring *GSTP1* function as a ‘caretaker’ gene during the pathogenesis of prostate cancer, in which the early loss of GSTP1 activity renders prostate cells vulnerable to genome damage associated with chronic prostatic inflammation and repeated exposure to carcinogens. The model predicts that the critical prostate carcinogens will be those that are substrates for GSTP1 detoxification and are associated with high prostate cancer risk diet and lifestyle habits.

Both genetic and environmental factors likely contribute to the pathogenesis of human prostate cancer. In support of a role for inheritance in the development of prostate cancer, familial clusters of the disease have been reported, and segregation analyses have suggested that prostate cancer in some of these families is likely attributable to inheritance of prostate cancer susceptibility genes [1–3]. Over the past few years, a number of genetic loci have been identified that have been postulated to be responsible for inherited susceptibility to prostate cancer [4–17]. How do such genes lead to prostate cancer development? Until the suspected genes have been identified and characterized, the manner by which the genes increase prostate cancer risk will remain to be established. Nevertheless, in a recent study of cancer risks among 44,788 pairs of twins in Sweden, Denmark, and Finland [18], a statistically significant effect of genotype was observed only for some 42% of prostate cancer cases (with a 95% confidence interval of 29–50%), indicating that environment and lifestyle likely play a more dominant role than inheritance in the development of most prostate cancers.

A dominant role for environment and lifestyle in the development of life-threatening prostate cancer is further supported by ecological epidemiology data [19]. Prostate cancer incidence and mortality are well-known to vary greatly in different geographic regions

of the world, with low risks of prostate cancer mortality characteristic of Asia and high risks of prostate cancer mortality characteristic of the US and Western Europe [19,20]. In addition, Asian immigrants to the US tend to adopt higher prostate cancer risks, strong evidence that the environment and lifestyle may be the major cause of life-threatening prostate cancer in the US [21,22]. How do the environment and lifestyle promote prostatic carcinogenesis? Insights into the earliest steps in the molecular pathogenesis of human prostate cancer have begun to provide a clue: prostate cancer development appears to be initiated by somatic inactivation of *GSTP1*, encoding the human π -class glutathione *S*-transferase [23–26]. Cancer cell DNA typically contains many somatic alterations, including mutations, deletions, amplifications, translocations, and hypermethylated CpG islands, that affect the function of critical genes and contribute to the malignant phenotype [27,28]. Critical genes targeted include oncogenes, which promote transformation when activated, and tumor suppressor genes, which fail to prevent transformation when inactivated. During the development of prostate cancer, *GSTP1* does not appear to function either as an oncogene, or as a tumor suppressor gene [25]. Instead, *GSTP1* likely acts as a ‘caretaker’ gene in prostate cells, which when inactivated, fails to prevent further somatic genome alterations upon chronic exposure to genome-damaging stresses [25,29]. In




Figure 1. A new model for prostatic carcinogenesis. GSTP1 (gray) is constantly expressed in prostate basal epithelial cells, but can be induced in luminal epithelial cells upon exposure to reactive chemical species, including electrophiles and oxidants, that threaten genome damage. Loss of GSTP1 expression accompanying *GSTP1* CpG island hypermethylation increases vulnerability of luminal cells to neoplastic transformation and malignant progression.

this model of prostatic carcinogenesis, loss of *GSTP1* function increases the vulnerability of prostate cells to carcinogenic insults, contributed by environment and lifestyle, that promote prostate cancer development and progression (Figure 1).

Recognition that somatic inactivation of *GSTP1* may initiate prostatic carcinogenesis permits a reevaluation of the influence of environment and lifestyle on the development of life-threatening prostate cancer. Epidemiology studies have identified several environmental factors, including the diet, sexually transmitted diseases, and others, as candidate prostate cancer risk factors [30–32]. However, no specific prostate carcinogens have been identified. Glutathione *S*-transferases (GSTs), like GSTP1, are capable of detoxifying oxidant and electrophilic chemicals to prevent cell and genome damage [33]. If loss of *GSTP1* function plays a critical role in the pathogenesis of prostate cancer, we can expect that the relevant prostate carcinogens will be oxidants and electrophiles that (i) are substrates for GSTP1 detoxification, (ii) are associated with high prostate cancer risk diet and lifestyle habits, and (iii) are able to reach prostate cells with defective GSTP1 ‘caretaker’ activity to inflict cell and genome injury. In this review, we consider the influence of environment and lifestyle on the development of life-threatening prostate cancer, focusing on the acquired vulnerability of prostate cells to carcinogens and other genome-damaging stresses associated with loss of *GSTP1* function.

The molecular pathogenesis of prostate cancer: Somatic inactivation of *GSTP1*

The pathogenesis of prostate cancer likely proceeds over at least 30 or more years. Small prostate cancers have been found in >30% of men aged 30–40 years, while diagnoses of clinically significant prostate cancers are typically made in men aged 60–70 years [34]. Prostate cancer cells typically contain a myriad of somatic genome alterations, including gene mutations, gene amplifications, gene deletions, chromosomal rearrangements, and changes in DNA methylation (see Figure 2) [35]. In addition, cancer cells from different prostate cancer cases, from different prostate cancer lesions in individual cases, or from different areas within the same prostate cancer lesions, often display marked differences in somatic genome changes [35,36]. This heterogeneity of somatic genome defects in prostate cancer cells, accumulating over decades, suggests that prostate cancers may arise as a consequence of either chronic or prolonged exposure to genome-damaging stresses, defective maintenance of genome integrity, or a combination of both processes. Furthermore, ongoing genomic instability in prostate cancer cells may be what leads to metastasis, progression to androgen independence after attempts at hormonal therapy, and other malignant behaviors [37–42].

Over the past several years, evidence has accumulated in support of the concept that somatic inactivation of *GSTP1*, most commonly by CpG island hypermethylation, may be the initiating somatic genome lesion for prostatic carcinogenesis, increasing the rate at which additional somatic genome alterations appear in response to exposure to genome-damaging stresses [23–26]. GSTs stereotypically provide an inducible defense against macromolecular damage by reactive chemical species [33]. In the normal prostate, GSTP1 polypeptides are selectively present in basal epithelial cells, with little, if any, enzyme detectable in non-stressed columnar epithelial cells (Figure 1). Nonetheless, *GSTP1* expression appears strikingly induced in columnar epithelial cells subjected to genome damaging stresses, such as those accompanying inflammation, commonly present in the human prostate. Proliferative inflammatory atrophy (PIA) lesions have been proposed to arise as a consequence of inflammatory injury to the prostate epithelium followed by exuberant epithelial cell proliferation/regeneration, and to give rise directly to prostatic intraepithelial neoplasia (PIN) lesions, known




Figure 2. The molecular pathogenesis of prostate cancer. Somatic genome lesions accumulate during prostate cancer development. *GSTP1* CpG island hypermethylation appears to serve as the initiating somatic genome lesion.

precursors to prostate cancer [43,44]. In PIA lesions, atrophic luminal cells characteristically contain high levels of GSTP1 and other GSTs [43,45]. However, in PIN lesions, in contrast to PIA lesions, absent GSTP1 expression is a consistent finding [46]. Prostate cancer cells are also characteristically devoid of GSTP1. In almost all prostate cancer cases, the absence of GSTP1 expression can be attributed to somatic *GSTP1* CpG island hypermethylation, not present in DNA from normal tissues, but evident in DNA from some 70% of PIN lesions and >90% of prostate cancers [25,46]. As the most common somatic genome alteration yet reported for human prostate cancer, *GSTP1* CpG island hypermethylation, readily discriminated using various sensitive techniques, is an attractive candidate molecular biomarker for prostate cancer cells, under development for use in prostate cancer detection, diagnosis, and staging (Table 1) [23–25,46–54].

GSTP1 CpG island hypermethylation appears to prevent GSTP1 expression in prostate cancer cells via ‘silencing’ of *GSTP1* transcription. In support of this mechanism, LNCaP prostate cancer cells contain only hypermethylated *GSTP1* CpG island alleles, fail to express either *GSTP1* mRNA or GSTP1 polypeptides, and exhibit reduced *GSTP1* transcription, whereas treatment of LNCaP cells *in vitro* or *in vivo* with DNA methyltransferase inhibitors both reverses *GSTP1* CpG island hypermethylation and restores GSTP1

expression [24,25,55]. In addition, while unmethylated *GSTP1* transcriptional regulatory sequences readily promote abundant CAT reporter expression after transfection into LNCaP prostate cancer cells, *SssI* CpG-methylase treatment of the *GSTP1* promoter before transfection leads to a marked reduction in CAT reporter expression [25,56]. Finally, detailed molecular pathology analyses have revealed, for almost all prostate cancer cases, that each of the cancer cells contains only hypermethylated *GSTP1* CpG island alleles, and each fails to express GSTP1 [25]. In contrast, for the two prostate cancer cases thus far described in which prostate cancer cells have been found to contain unmethylated *GSTP1* CpG island alleles, each of the prostate cancer cells displayed abundant GSTP1 expression [25,54]. CpG island hypermethylation has been proposed to repress gene transcription directly, via interference with transcriptional trans-activator binding, or indirectly, via the actions of ^{5-m}C-binding proteins that affect chromatin structure, leading to a transcriptionally inactive chromatin conformation [57–63]. For the *GSTP1* CpG island, data collected thus far suggest that CpG island hypermethylation acts to repress *GSTP1* transcription indirectly [56]. Regardless of the mechanism by which CpG island hypermethylation prevents *GSTP1* transcription, because CpG methylation patterns are maintained through mitosis by the action of DNA methyltransferases,

Table 1. *GSTP1* CpG island hypermethylation as a molecular biomarker for prostate cancer

Reference	Assay technique	<i>GSTP1</i> CpG island hypermethylation*
Lee et al. (1994)	Southern blot	Tissue: 100% PCA; 0% BPH
Lee et al. (1997)	Restriction enzyme-PCR	Tissue: 91% PCA
Brooks et al. (1998)	Restriction enzyme-PCR	Tissue: 70% PIN
Santourlidis et al. (1999)	Restriction enzyme-PCR	Tissue: 75% PCA; 0% TCC
Millar et al. (1999)	Bisulfite genomic sequencing	Tissue: 83% PCA
Suh et al. (2000)	Restriction enzyme-PCR	Ejaculate: 44% PCA
Goessl et al. (2000)	Methylation-specific PCR	Tissue: 94% PCA; 0% BPH Urine: 36% PCA Ejaculate: 50% PCA Plasma: 50% PCA
Goessl et al. (2001)	Methylation-specific PCR	Urine: 73% PCA; 29% PIN; 2% BPH
Cairns et al. (2001)	Methylation-specific PCR	Tissue: 79% PCA Urine: 27% PCA
Lin et al. (2001)	Southern blot, restriction enzyme-PCR, bisulfite genomic sequencing	Tissue: 100% PCA
Goessl et al. (2001)	Methylation-specific PCR	Tissue: 90% PCA; 0% BPH Urine: 76% PCA Ejaculate: 50% PCA Plasma: 72% PCA
Jeronimo et al. (2001)	Quantitative methylation-specific PCR	Tissue: 91% PCA; 54% PIN; 29% BPH

*PCA-prostate cancer, BPH-benign prostatic hyperplasia, PIN-prostatic intraepithelial neoplasia, TCC-transitional cell carcinoma.

GSTP1 transcriptional 'silencing' associated with CpG island hypermethylation may be subject to selection during prostatic carcinogenesis. For this reason, the observation that most prostate cancers contain only prostate cancer cells with hypermethylated *GSTP1* CpG island alleles may be evidence that loss of *GSTP1* function likely provides some sort of selective growth or survival advantage at some point during the pathogenesis of prostate cancer.

Prostate cancer epidemiology: The diet, prostate inflammation, and prostate cancer risk

The etiology of prostate cancer has not been established. However, as mentioned previously, both prostate cancer incidence and mortality vary greatly in different geographic regions, with generally low risks of prostate cancer development characteristic of Asia, and generally high risks of prostate cancer development characteristic of the US and Western Europe [21,64–67]. Of note, prostate cancer risk among ethnic Asian immigrants to North America increases with duration of exposure to a Western lifestyle: Asian immigrants to North America have a higher risk of prostate cancer after living in North America for more than 25 years than after living in North America for less than 10 years [22]. Asian men born in the US have

a risk for life-threatening prostate cancer development similar to Caucasian men. Of course, these epidemiological observations underscore the critical role for environment and lifestyle in fostering the epidemic of prostate cancer afflicting men in the US. The major environment and lifestyle factor modulating prostate cancer risk appears to be the diet. Clearly, stereotypical Asian diets are quite different from stereotypical Western diets. Unfortunately, whether the stereotypical Western diet makes an error of *commission* (e.g. over-consumption of dietary components increasing prostate cancer risks), an error of *omission* (e.g. under-consumption of dietary components decreasing prostate cancer risks), or *both* has been difficult to establish. Nonetheless, if specific dietary components could be demonstrated to change prostate cancer risks, the component could be *avoided* if it promoted prostate cancer and *provided* if it protected against prostate cancer as a rational prostate cancer prevention strategy.

Ecological, case-control, and cohort epidemiology studies all have long implicated fats and meats as candidate risks factors for prostate cancer (for a review, see [30]). The challenge for epidemiologists has been to ascertain whether increased prostate cancer risks might be attributable to total fat intake, to increased energy intake associated with high-fat diets, to intake of specific fats, or to intake of fats from specific sources, such as red meats. For example, in the Health Professionals

Follow-up Study, a prospective cohort of 51,529 men, total fat intake, adjusted for energy intake, appeared to confer increased risks (a relative risk of 1.79 for high *versus* low quintile of intake, with 95% confidence interval of 1.04–3.07) of prostate cancer development [68]. However, saturated fat intake *per se* did not appear to be responsible for the increased prostate cancer risks. Rather, animal fat intake (a relative risk of 1.63 with a 95% confidence interval of 0.95–2.78), particularly red meat intake (a relative risk of 2.64 with a 95% confidence interval of 1.21–5.77), may have been responsible for the observed prostate cancer risks associated with the high total fat diets. A deleterious effect of red meat consumption on prostate cancer risk has also been seen in the Physicians Health Study [69], a prospective cohort of 14,916 men (a relative risk of 2.5 for red meat consumption 5 times/week *versus* less than once/week with a 95% confidence interval of 0.9–6.7), and in a large cohort study in Hawaii [70], involving 20,316 men of varying ethnicities (a relative risk of 1.6 for highest *versus* lowest tertile of beef consumption with a 95% confidence interval of 1.1–2.4). Intriguingly, cooking of meats at high temperatures or on charcoal grills, is known to lead to the formation of heterocyclic amine carcinogens and/or polycyclic aromatic hydrocarbon carcinogens [71,72]. However, the level of intake of these carcinogenic substances has been difficult to estimate for epidemiologic studies of prostate cancer risk.

While chronic consumption of animal fats and red meats may promote prostate cancer, intake of vegetables may protect against prostate cancer development (for a review, see [31]). Among potentially protective vegetables, attention has been most intensively focused on tomatoes [73–76], which contain an α -carotenoid anti-oxidant, lycopene, and on cruciferous vegetables [76–78], which contain an anti-carcinogenic isothiocyanate, sulforaphane [79]. Lycopene, which appears in prostate tissues after tomato ingestion, can scavenge oxidant species, including nitric oxide, to prevent oxidative cell and genome damage [80–83]. In a study of 578 prostate cancer cases and 1294 controls from the Physician's Health Study cohort, high plasma lycopene levels were associated with a decrease in the risk of aggressive prostate cancer (a relative risk of 0.56 for high *versus* low quintile of lycopene plasma level with a 95% confidence interval of 0.34–0.91). Of interest, other anti-oxidants, selenium and vitamin E, have been found to attenuate prostate cancer development in randomized clinical trials [84–86]. Sulforaphane has been proposed to protect against

cancer development by increasing the expression of carcinogen-detoxification enzymes, including GSTs and quinone oxidoreductases, that help prevent genome damage mediated by carcinogens [79,87–89]. In a study of 628 prostate cancer cases and 602 controls from King County, Washington, cruciferous vegetable consumption, adjusted for total vegetable intake, was associated with diminished prostate cancer risks (a relative risk of 0.59 for 3 or more servings/week *versus* less than one serving/week with a 95% confidence interval of 0.39–0.90) [77].

In addition to the diet, environment and lifestyle factors affecting prostate cancer risk include sexually transmitted diseases, risk factors that might be modified to prevent prostate cancer [32]. However, unlike cancer of the uterine cervix, nasopharyngeal carcinoma, and Kaposi's sarcoma, it has been difficult to identify a specific pathogen responsible for the direct transformation of prostatic cells. Instead, prostatic inflammation associated with sexually transmitted infections may play a more important role in prostatic carcinogenesis. In a recent population-based study involving 981 prostate cancer cases (479 black men, 502 white men) and 1315 controls (594 black men, 721 white men), prostate cancer risks were increased among men (i) who reported a history of gonorrhoea or syphilis (a relative risk of 1.6 with a 95% confidence interval of 1.2–2.1), (ii) who reported three or more episodes of gonorrhea (a relative risk of 3.3 with a 95% confidence interval of 1.4–7.8), and (iii) who displayed serological evidence of syphilis (a relative risk of 1.8 with a 95% confidence interval of 1.0–3.5) [90]. Inflammation, known to inflict oxidative damage, has been thought to contribute to the pathogenesis of many human cancers [91]. Prostatitis, whether the result of an identified infection or idiopathic, very commonly afflicts men as they grow older in the US [92]. Whether chronic or recurrent prostatic inflammation contributes to prostate cancer development, or might explain differences in prostate cancer risks between different geographic regions, has not been ascertained.

A 'caretaker' role for *GSTP1* during prostatic carcinogenesis: The integration of prostate cancer molecular biology and epidemiology into a unified model

Rather than function as a tumor suppressor during prostatic carcinogenesis, *GSTP1* appears most likely to act as a 'caretaker' [29], defending prostate cells




Figure 3. Loss of GSTP1 activity leads to a failure of cellular protection against oxidant and electrophilic carcinogens.

against genome damage mediated by carcinogens (Figure 3). GSTs have long been thought to protect against cancer development by catalyzing conjugation reactions between glutathione and a variety of reactive chemicals species, preventing carcinogen-induced cell and genome damage [33]. The enzymes are dimers composed of subunit polypeptides encoded by a complex collection of genes, organized into gene families α , μ , π , and θ . *GSTP1* encodes the single π -class GST subunit polypeptide; GSTP1-1 is a homodimer. The various different GSTs are normally expressed in different cells and tissues; however, GST activity can also be stereotypically induced in other cells and tissues in response to chemical stresses, via increases in *GST* subunit gene transcription [93,94]. The transcription factor Nrf2, binding to specific *cis*-regulatory sequences in *GST* subunit gene promoters, appears to play a critical role in activating *GST* subunit gene transcription in response to chemical stresses [95]. The inducible protection against reactive chemicals afforded by GSTs comprises a substantial barrier to chemical carcinogenesis. For example, mice carrying disrupted *Gstp1/2* genes, encoding the murine π -class GSTs, display increased skin tumorigenesis upon topical exposure to 7,12-dimethylbenz anthracene (DMBA) [96]. In addition, mice carrying disrupted *Nrf2* genes manifest increased gastric carcinogenesis upon exposure to benzo[a]pyrene [95]. By providing a barrier to cell and genome damage mediated by reactive chemicals, *GST* genes act as ‘caretaker’ genes for cancers arising as a consequence of carcinogen exposure.

As described above, prostate cancer epidemiology data have implicated heterocyclic amines (well-done meats), polycyclic aromatic hydrocarbons (‘char-broiled’ meats), and oxidants (inflammation) as candidate reactive chemicals that might threaten cell and

genome damage leading to prostate cancer development. If *GSTP1* ‘caretaker’ function provides a defense against such carcinogens, then the reactive chemical species ought to be substrates for GSTP1 detoxification. Recent data have suggested that heterocyclic amine carcinogens present in well-done meats may be detoxified by GSTP1 [97]. The heterocyclic aromatic amine carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) is known to cause mutations by adduction to DNA bases after metabolic activation by various cellular enzymes [71,98–101]. Rats fed PhIP have been reported to display mutations in prostate DNA and develop prostate cancer [102–104]. However, when LNCaP human prostate cancer cells devoid of GSTP1 were exposed to metabolically activated PhIP, high levels of PhIP-DNA adducts were detected, while LNCaP prostate cancer cells genetically modified to express GSTP1 exhibited substantial resistance to the formation of pro-mutagenic PhIP-DNA adducts [97]. Like the heterocyclic amine carcinogens, metabolically activated polycyclic aromatic hydrocarbon carcinogens are also GSTP1 substrates [105]. Of interest, polymorphic GSTP1 variants that are homodimers with subunit polypeptides containing isoleucine at amino acid 105, or containing valine at amino acid 105, appear to exhibit different catalytic properties when confronted with polycyclic aromatic hydrocarbons [106]. Furthermore, genetic epidemiology studies have suggested that homozygosity for *GSTP1*^{val-105} alleles may increase breast cancer risks (a relative risk of 1.97 for *GSTP1*^{val-105/val-105} with a 95% confidence interval of 0.77–5.02) [107]. The prostate cancer risks associated with homozygosity for *GSTP1*^{val-105} alleles have not been established. Finally, new preliminary data have suggested that oxidants may also be substrates for GSTP1: upon prolonged exposure to an oxidative stress, LNCaP prostate cancer cells genetically modified to express GSTP1 suffered less oxidative genome damage than LNCaP prostate cancer cells devoid of GSTP1 activity (DeWeese et al. unpublished data).

Remarkably, loss of *GSTP1* ‘caretaker’ function may provide a selective growth or survival advantage upon exposure to certain carcinogens. Usually, increased sensitivity to the genome damaging actions of chemical carcinogens, such as that seen upon metabolically activated PhIP treatment of LNCaP prostate cancer cells devoid of GSTP1 *versus* LNCaP cells expressing GSTP1, is accompanied by increased sensitivity to the cytotoxic effects of the same carcinogens [97]. However, new preliminary data have revealed that

when LNCaP prostate cancer cells, devoid of GSTP1 activity, are challenged by prolonged exposures to oxidant stresses, the cells appear to suffer much less cell death than LNCaP prostate cancer cells genetically modified to express GSTP1 (DeWeese et al. unpublished data). As a result, for LNCaP prostate cancer cells, loss of GSTP1 ‘caretaker’ activity appeared to result in increased genome damage and decreased cell death upon oxidant exposure. Perhaps, this phenotype of oxidation damage ‘tolerance’ may provide PIN cells or prostate cancer cells a selective growth or survival advantage in the face of an oxidative stress like chronic prostate inflammation [26,43]. A similar phenotype of cell and genome damage ‘tolerance’ associated with loss of π -class GST activity has been reported for studies of mice exposed to acetaminophen overdoses, in which *Gstp1/2^{+/+}* mice appeared to suffer markedly more hepatotoxicity than *Gstp1/2^{-/-}* mice after administration of high doses of acetaminophen [108]. The mechanism(s) by which π -class GST activity might be coupled to cell death upon exposure to oxidants or to acetaminophen could potentially include glutathione ‘bankruptcy’ associated with π -class GST metabolism of reactive chemical species [108], toxification of the chemical species by π -class GST conjugation of the chemical species with glutathione [109], and/or potential modulation of stress-associated signal transduction pathways by π -class GSTs [110–113].

Thus, with loss of *GSTP1* ‘caretaker’ function as the initiating somatic genome lesion in the pathogenesis of prostate cancer, several features of the molecular pathogenesis of prostate cancer and several features of prostate cancer epidemiology can be integrated into a coherent model (Figure 4) [26]. In this model, the earliest steps in prostatic carcinogenesis occur as a result of prostatic inflammation, associated with prostatic infections or with idiopathic prostatitis, that leads to prostate cell oxidant injury and regeneration, resulting in the appearance of PIA lesions. The PIA cells, expressing high levels of GSTP1 and other oxidant protection enzymes, give rise to PIN cells, devoid of GSTP1, and ultimately to prostate cancer cells. With loss of *GSTP1* ‘caretaker’ function in the PIN cells and prostate cancer cells, decades of chronic oxidant stress, and of exposure to dietary heterocyclic amine and polycyclic aromatic hydrocarbons, lead to an accumulation of somatic genome alterations that drive malignant progression. Of interest, prostatic carcinogenesis may proceed via a similar pathway in rats, where both prostatic inflammation and heterocyclic




Figure 4. An integrated model for prostate cancer development featuring *GSTP1* inactivation in prostate cells in the face of exposure to inflammatory oxidants and dietary oxidants and electrophiles.

aromatic amine exposures have been reported to lead to prostatic neoplasia [102,103,114,115].

Rational strategies for prostate cancer prevention

To interrupt prostate cancer pathogenesis and prevent life-threatening prostate cancer, several rational approaches might be considered, including: (i) *attenuation* or *abrogation* of genome damaging stresses via avoidance of exogenous carcinogens (such as heterocyclic amines and polycyclic aromatic hydrocarbons) and/or reduction of endogenous carcinogenic stresses (such as inflammatory oxidants), (ii) *restoration* of *GSTP1* expression via treatment with inhibitors of CpG methylation, and (iii) *compensation* for inadequate *GSTP1* activity via treatment with inducers of general GST activity. Some strategies for *attenuation* or *abrogation* of genome damaging stresses are already under evaluation for prostate cancer prevention. Antioxidant micronutrients, including vitamin E, selenium, and carotenoids, such as lycopene, might be expected to reduce oxidant damage to prostate cell DNA if used for prostate cancer prevention [73,74,116,117]. Each of these agents has reached human clinical studies. Selenium and vitamin E are to be tested in a large ($n = 32,400$) randomized trial (The Selenium and Vitamin E Chemoprevention Trial; SELECT) that is ongoing in the US to ascertain whether antioxidants selenium and vitamin E, alone or in combination, can reduce the incidence of prostate cancer in healthy men age 55 and older (age 50 and older for African-American men).

Early proof-of-principal clinical studies of lycopene, and/or tomato products, for prostate cancer have begun to be reported [118,119]. Anti-inflammatory agents might be expected to reduce inflammation-associated oxidant production in the prostate and reduce prostate cancer risks [120–122]. Several non-steroidal anti-inflammatory drugs are being examined as candidate drugs for prostate cancer prevention and/or treatment; ‘proof-of-principle’ clinical trials for prostate cancer are underway at Johns Hopkins featuring celecoxib (Celebrex®; Pharmacia) and sulindac.

Restoration of *GSTP1* function may be feasible as a prostate cancer prevention strategy (Figure 5). Silencing of *GSTP1* transcription in PIN and prostate cancer cells is likely maintained via the action of DNA methyltransferases, enzymes that could be targeted for therapeutic inhibition. We have collected data indicating that DNA methyltransferase inhibitors such as 5-aza-deoxycytidine and procainamide can restore *GSTP1* expression in LNCaP prostate cancer cells *in vivo* [55]. These agents may be considered candidate prostate cancer prevention drugs. In addition, data have been reported revealing interactions both between ^{5-m}C-binding proteins and histone deacetylases (HDACs), and between DNA methyltransferases and HDACs, in effecting transcriptional repression [60–62,123]. Perhaps, combinations of drugs active at inhibiting DNA methyltransferases and at inhibiting chromatin-remodeling enzymes might prove useful in restoring high level *GSTP1* expression in PIN or prostate cancer cells to slow life-threatening prostate cancer progression [61]. The key issue for development of this approach, especially if it is to be used for prostate cancer prevention as well as for prostate cancer treatment, will be whether restoration of *GSTP1*

expression (as well as the expression of other genes) can be accomplished with reasonable gene selectivity and with acceptable side effects. To this end, early phase I studies of combinations of the DNA methyltransferase inhibitor 5-aza-cytidine and the HDAC inhibitor phenylbutyrate have been initiated at Johns Hopkins.

Therapeutic compensation for inadequate ‘caretaker’ gene function may also hold promise for cancer prevention (Figure 6). Although *GSTP1* may be silenced early during the pathogenesis of prostate cancer, genes encoding other GST subunit polypeptides appear intact. As the genes encoding the other GST subunit polypeptides can be induced, via an Nrf2-dependent mechanism, this pathway can be exploited to better defend *GSTP1*-deficient prostate cells against injurious chemical stresses [93,94]. Augmentation of carcinogen-detoxification capacity, using a variety of such chemoprotective compounds, including isothiocyanates, 1,2-dithiole-3-thiones, terpenoids, etc., has been reported to prevent a variety of different cancers in different animal models by triggering the expression of carcinogen-detoxification enzymes [124]. Most or all of these compounds likely act to prevent cancer by activating carcinogen-detoxification enzyme gene expression via the Nrf2-dependent transcription induction pathway, as oltipraz, an anti-schistosomal 1,2-dithiole-3-thione compound known to protect against benzo[a]pyrene gastric carcinogenesis in murine models, had no effect on gastric tumor formation in mice carrying disrupted *Nrf2* alleles [95]. Induction of GST ‘caretaker’ activity in liver tissues, using oltipraz, a therapeutic inducer of GST activity, has been shown to reduce aflatoxin B₁




Figure 5. Therapeutic restoration of *GSTP1* function using DNA methyltransferase inhibitors.




Figure 6. Therapeutic compensation for lack of *GSTP1* activity using inducers of carcinogen detoxification enzymes, such as the isothiocyanate, sulforaphane, present in cruciferous vegetables.

damage when administered to a human clinical study cohort at high risk for aflatoxin exposure and liver cancer development in China [125–127]. Of greatest interest, many carcinogen-detoxification enzyme inducers have been detected in dietary components. For example, sulforaphane, an isothiocyanate that can trigger carcinogen-detoxification enzyme induction, is present in high amounts in cruciferous vegetables [79,87]. As described above, diets rich in carcinogen-inducers like sulforaphane have been associated with decreased cancer risks [77]. Early ‘proof-of-principle’ clinical trials of dietary carcinogen inducers as prevention for many different human cancers are currently underway [89,128].

Conclusions

Although both genes and the environment can contribute to the development of human prostate cancer, lifestyle factors, particularly the diet and possibly sexually transmitted diseases, likely play the dominant role in prostatic carcinogenesis. Loss of *GSTP1* ‘caretaker’ function, the earliest somatic genome alteration yet described for prostate cancer, appears to render prostate cells vulnerable to carcinogenic stresses. Heterocyclic amines, polycyclic aromatic hydrocarbons, and inflammatory oxidants, all carcinogens likely to be associated with high-risk prostate cancer diets and behaviors, are substrates for *GSTP1* detoxification. In the absence of *GSTP1* ‘caretaker’ activity, chronic exposure to such carcinogens promotes an accumulation of somatic genome abnormalities, targeting oncogenes and tumor suppressor genes, that leads to life-threatening prostate cancer progression. Therapeutic strategies to *abrogate* genome damaging stresses, to *restore GSTP1* ‘caretaker’ function, or to *compensate* for inadequate *GSTP1* ‘caretaker’ gene function via induction of carcinogen-detoxification enzymes, might be expected to attenuate prostatic carcinogenesis.

Acknowledgement

William G. Nelson has a patent (US Patent 5,552,277) entitled ‘Genetic Diagnosis of Prostate Cancer.’

References

1. Steinberg GD, Carter BS, Beaty TH, Childs B, Walsh PC: Family history and the risk of prostate cancer. *Prostate* 17: 337, 1990
2. Carter BS, Steinberg GD, Beaty TH, Childs B, Walsh PC: Familial risk factors for prostate cancer. *Cancer Surv* 11: 5, 1991
3. Carter BS, Beaty TH, Steinberg GD, Childs B, Walsh PC: Mendelian inheritance of familial prostate cancer. *Proc Natl Acad Sci USA* 89: 3367, 1992
4. Xu J, Zheng SL, Hawkins GA, Faith DA, Kelly B, Isaacs SD, Wiley KE, Chang B, Ewing CM, Bujnovszky P, Carpten JD, Bleecker ER, Walsh PC, Trent JM, Meyers DA, Isaacs WB: Linkage and association studies of prostate cancer susceptibility: evidence for linkage at 8p22-23. *Am J Hum Genet* 69: 341, 2001
5. Zheng SL, Xu J, Isaacs SD, Wiley K, Chang B, Bleecker ER, Walsh PC, Trent JM, Meyers DA, Isaacs WB: Evidence for a prostate cancer linkage to chromosome 20 in 159 hereditary prostate cancer families. *Hum Genet* 108: 430, 2001
6. Xu J, Zheng SL, Chang B, Smith JR, Carpten JD, Stine OC, Isaacs SD, Wiley KE, Henning L, Ewing C, Bujnovszky P, Bleecker ER, Walsh PC, Trent JM, Meyers DA, Isaacs WB: Linkage of prostate cancer susceptibility loci to chromosome 1. *Hum Genet* 108: 335, 2001
7. Xu J, Zheng SL, Carpten JD, Nupponen NN, Robbins CM, Mestre J, Moses TY, Faith DA, Kelly BD, Isaacs SD, Wiley KE, Ewing CM, Bujnovszky P, Chang B, Bailey-Wilson J, Bleecker ER, Walsh PC, Trent JM, Meyers DA, Isaacs WB: Evaluation of linkage and association of HPC2/ELAC2 in patients with familial or sporadic prostate cancer. *Am J Hum Genet* 68: 901, 2001
8. Gronberg H, Xu J, Smith JR, Carpten JD, Isaacs SD, Freije D, Bova GS, Danber JE, Bergh A, Walsh PC, Collins FS, Trent JM, Meyers DA, Isaacs WB: Early age at diagnosis in families providing evidence of linkage to the hereditary prostate cancer locus (HPC1) on chromosome 1. *Cancer Res* 57: 4707, 1997
9. Gronberg H, Isaacs SD, Smith JR, Carpten JD, Bova GS, Freije D, Xu J, Meyers DA, Collins FS, Trent JM, Walsh PC, Isaacs WB: Characteristics of prostate cancer in families potentially linked to the hereditary prostate cancer 1 (HPC1) locus. *JAMA* 278: 1251, 1997
10. Smith JR, Freije D, Carpten JD, Gronberg H, Xu J, Isaacs SD, Brownstein MJ, Bova GS, Guo H, Bujnovszky P, Nusskern DR, Damber JE, Bergh A, Emanuelsson M, Kallioniemi OP, Walker-Daniels J, Bailey-Wilson JE, Beaty TH, Meyers DA, Walsh PC, Collins FS, Trent JM, Isaacs WB: Major susceptibility locus for prostate cancer on chromosome 1 suggested by a genome-wide search. *Science* 274: 1371, 1996
11. Goode EL, Stanford JL, Peters MA, Janer M, Gibbs M, Kolb S, Badzioch MD, Hood L, Ostrander EA, Jarvik GP: Clinical characteristics of prostate cancer in an analysis of linkage to four putative susceptibility loci. *Clin Cancer Res* 7: 2739, 2001
12. Peters MA, Jarvik GP, Janer M, Chakrabarti L, Kolb S, Goode EL, Gibbs M, DuBois CC, Schuster EF, Hood L, Ostrander EA, Stanford JL: Genetic linkage analysis of prostate cancer families to Xq27-28. *Hum Hered* 51: 107, 2001

13. Gibbs M, Stanford JL, Jarvik GP, Janer M, Badzioch M, Peters MA, Goode EL, Kolb S, Chakrabarti L, Shook M, Basom R, Ostrander EA, Hood L: A genomic scan of families with prostate cancer identifies multiple regions of interest. *Am J Hum Genet* 67: 100, 2000
14. Gibbs M, Stanford JL, McIndoe RA, Jarvik GP, Kolb S, Goode EL, Chakrabarti L, Schuster EF, Buckley VA, Miller EL, Brandzel S, Li S, Hood L, Ostrander EA: Evidence for a rare prostate cancer-susceptibility locus at chromosome 1p36. *Am J Hum Genet* 64: 776, 1999
15. Xu J, Meyers D, Freije D, Isaacs S, Wiley K, Nusskern D, Ewing C, Wilkens E, Bujnovszky P, Bova GS, Walsh P, Isaacs W, Schleutker J, Matikainen M, Tammela T, Visakorpi T, Kallioniemi OP, Berry R, Schaid D, French A, McDonnell S, Schroeder J, Blute M, Thibodeau S, Trent J: Evidence for a prostate cancer susceptibility locus on the X chromosome. *Nat Genet* 20: 175, 1998
16. Berry R, Schroeder JJ, French AJ, McDonnell SK, Peterson BJ, Cunningham JM, Thibodeau SN, Schaid DJ: Evidence for a prostate cancer-susceptibility locus on chromosome 20. *Am J Hum Genet* 67: 82, 2000
17. Berthon P, Valeri A, Cohen-Akenine A, Drelon E, Paiss T, Wöhr G, Latil A, Millasseau P, Mellah I, Cohen N, Blanche H, Bellane-Chantelot C, Demenais F, Teillac P, Le Duc A, de Petriconi R, Hautmann R, Chumakov I, Bachner L, Maitland NJ, Lidereau R, Vogel W, Fournier G, Mangin P, Cussenot O: Predisposing gene for early-onset prostate cancer, localized on chromosome 1q42.2-43. *Am J Hum Genet* 62: 1416, 1998
18. Lichtenstein P, Holm NV, Verkasalo PK, Iliadou A, Kaprio J, Koskenvuo M, Pukkala E, Skytthe A, Hemminki K: Environmental and heritable factors in the causation of cancer – analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 343: 78, 2000
19. Hsing AW, Devesa SS: Trends and patterns of prostate cancer: What do they suggest? *Epidemiol Rev* 23: 3, 2001
20. Hsing AW, Tsao L, Devesa SS: International trends and patterns of prostate cancer incidence and mortality. *Int J Cancer* 85: 60, 2000
21. Shimizu H, Ross RK, Bernstein L, Yatani R, Henderson BE, Mack TM: Cancers of the prostate and breast among Japanese and white immigrants in Los Angeles County. *Br J Cancer* 63: 963, 1991
22. Whittemore AS, Kolonel LN, Wu AH, John EM, Gallagher RP, Howe GR, Burch JD, Hankin J, Dreon DM, West DW: Prostate cancer in relation to diet, physical activity, and body size in blacks, whites, and Asians in the United States and Canada. *J Natl Cancer Inst* 87: 652, 1995
23. Lee WH, Isaacs WB, Bova GS, Nelson WG: CG island methylation changes near the GSTP1 gene in prostatic carcinoma cells detected using the polymerase chain reaction: a new prostate cancer biomarker. *Cancer Epidemiol Biomarkers Prev* 6: 443, 1997
24. Lee WH, Morton RA, Epstein JI, Brooks JD, Campbell PA, Bova GS, Hsieh WS, Isaacs WB, Nelson WG: Cytidine methylation of regulatory sequences near the pi-class glutathione S-transferase gene accompanies human prostatic carcinogenesis. *Proc Natl Acad Sci USA* 91: 11733, 1994
25. Lin X, Tascilar M, Lee WH, Vles WJ, Lee BH, Veeraswamy R, Asgari K, Freije D, van Rees B, Gage WR, Bova GS, Isaacs WB, Brooks JD, DeWeese TL, De Marzo AM, Nelson WG: GSTP1 CpG island hypermethylation is responsible for the absence of GSTP1 expression in human prostate cancer cells. *Am J Pathol* 159: 1815, 2001
26. Nelson WG, De Marzo AM, DeWeese TL: The molecular pathogenesis of prostate cancer: Implications for prostate cancer prevention. *Urology* 57: 39, 2001
27. Fearon ER, Vogelstein B: A genetic model for colorectal tumorigenesis. *Cell* 61: 759, 1990
28. Baylin SB, Herman JG: DNA hypermethylation in tumorigenesis: Epigenetics joins genetics. *Trends Genet* 16: 168, 2000
29. Kinzler KW, Vogelstein B: Cancer-susceptibility genes. Gatekeepers and caretakers (news; comment) (see comments). *Nature* 386: 761, 1997
30. Kolonel LN: Fat, meat, and prostate cancer. *Epidemiol Rev* 23: 72, 2001
31. Chan JM, Giovannucci EL: Vegetables, fruits, associated micronutrients, and risk of prostate cancer. *Epidemiol Rev* 23: 82, 2001
32. Strickler HD, Goedert JJ: Sexual behavior and evidence for an infectious cause of prostate cancer. *Epidemiol Rev* 23: 144, 2001
33. Hayes JD, Pulford DJ: The glutathione S-transferase supergene family: Regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit Rev Biochem Mol Biol* 30: 445, 1995
34. Sakr WA, Grignon DJ, Crissman JD, Heilbrun LK, Cassin BJ, Pontes JJ, Haas GP: High grade prostatic intraepithelial neoplasia (HGPIN) and prostatic adenocarcinoma between the ages of 20–69: an autopsy study of 249 cases. *In vivo* 8: 439, 1994
35. Isaacs WB, Bova GS, Morton RA, Bussemakers MJ, Brooks JD, Ewing CM: Genetic alterations in prostate cancer. *Cold Spring Harb Symp Quant Biol* 59: 653, 1994
36. Ruijter ET, Miller GJ, van de Kaa CA, van Bokhoven A, Bussemakers MJ, Debruyne FM, Ruiter DJ, Schalken JA: Molecular analysis of multifocal prostate cancer lesions. *J Pathol* 188: 271, 1999
37. Suzuki H, Freije D, Nusskern DR, Okami K, Cairns P, Sidransky D, Isaacs WB, Bova GS: Interfocal heterogeneity of PTEN/MMAC1 gene alterations in multiple metastatic prostate cancer tissues. *Cancer Res* 58: 204, 1998
38. Cairns P, Okami K, Halachmi S, Halachmi N, Esteller M, Herman JG, Jen J, Isaacs WB, Bova GS, Sidransky D: Frequent inactivation of PTEN/MMAC1 in primary prostate cancer. *Cancer Res* 57: 4997, 1997
39. Jarrard DF, Bova GS, Ewing CM, Pin SS, Nguyen SH, Baylin SB, Cairns P, Sidransky D, Herman JG, Isaacs WB: Deletional, mutational, and methylation analyses of CDKN2 (p16/MTS1) in primary and metastatic prostate cancer. *Genes Chromosomes Cancer* 19: 90, 1997
40. Dong JT, Suzuki H, Pin SS, Bova GS, Schalken JA, Isaacs WB, Barrett JC, Isaacs JT: Down-regulation of the KAI1 metastasis suppressor gene during the progression of

- human prostatic cancer infrequently involves gene mutation or allelic loss. *Cancer Res* 56: 4387, 1996
41. Cher ML, Bova GS, Moore DH, Small EJ, Carroll PR, Pin SS, Epstein JI, Isaacs WB, Jensen RH: Genetic alterations in untreated metastases and androgen-independent prostate cancer detected by comparative genomic hybridization and allelotyping. *Cancer Res* 56: 3091, 1996
 42. Visakorpi T, Kallioniemi AH, Syvanen AC, Hyytinen ER, Karhu R, Tammela T, Isola JJ, Kallioniemi OP: Genetic changes in primary and recurrent prostate cancer by comparative genomic hybridization. *Cancer Res* 55: 342, 1995
 43. De Marzo AM, Marchi VL, Epstein JI, Nelson WG: Proliferative inflammatory atrophy of the prostate: implications for prostatic carcinogenesis. *Am J Pathol* 155: 1985, 1999
 44. Putzi MJ, De Marzo AM: Morphologic transitions between proliferative inflammatory atrophy and high-grade prostatic intraepithelial neoplasia (In process citation). *Urology* 56: 828, 2000
 45. Parsons JK, Nelson CP, Gage WR, Nelson WG, Kensler TW, De Marzo AM: GSTA1 expression in normal, preneoplastic, and neoplastic human prostate tissue. *Prostate* 49: 30, 2001
 46. Brooks JD, Weinstein M, Lin X, Sun Y, Pin SS, Bova GS, Epstein JI, Isaacs WB, Nelson WG: CG island methylation changes near the GSTP1 gene in prostatic intraepithelial neoplasia. *Cancer Epidemiol Biomarkers Prev* 7: 531, 1998
 47. Cairns P, Esteller M, Herman JG, Schoenberg M, Jeronimo C, Sanchez-Cespedes M, Chow NH, Grasso M, Wu L, Westra WB, Sidransky D: Molecular detection of prostate cancer in urine by GSTP1 hypermethylation. *Clin Cancer Res* 7: 2727, 2001
 48. Goessl C, Krause H, Muller M, Heicappell R, Schrader M, Sachsinger J, Miller K: Fluorescent methylation-specific polymerase chain reaction for DNA-based detection of prostate cancer in bodily fluids. *Cancer Res* 60: 5941, 2000
 49. Goessl C, Muller M, Heicappell R, Krause H, Straub B, Schrader M, Miller K: DNA-based detection of prostate cancer in urine after prostatic massage. *Urology* 58: 335, 2001
 50. Goessl C, Muller M, Heicappell R, Krause H, Miller K: DNA-based detection of prostate cancer in blood, urine, and ejaculates. *Ann N Y Acad Sci* 945: 51, 2001
 51. Jeronimo C, Usadel H, Henrique R, Oliveira J, Lopes C, Nelson WG, Sidransky D: Quantitation of GSTP1 methylation in non-neoplastic prostatic tissue and organ-confined prostate adenocarcinoma. *J Natl Cancer Inst* 93: 1747, 2001
 52. Suh CI, Shanafelt T, May DJ, Shroyer KR, Bobak JB, Crawford ED, Miller GJ, Markham N, Glode LM: Comparison of telomerase activity and GSTP1 promoter methylation in ejaculate as potential screening tests for prostate cancer. *Mol Cell Probes* 14: 211, 2000
 53. Santourlidis S, Florl A, Ackermann R, Wirtz HC, Schulz WA: High frequency of alterations in DNA methylation in adenocarcinoma of the prostate. *Prostate* 39: 166, 1999
 54. Millar DS, Ow KK, Paul CL, Russell PJ, Molloy PL, Clark SJ: Detailed methylation analysis of the glutathione S-transferase pi (GSTP1) gene in prostate cancer. *Oncogene* 18: 1313, 1999
 55. Lin X, Asgari K, Putzi MJ, Gage WR, Yu X, Cornblatt BS, Kumar A, Piantadosi S, DeWeese TL, De Marzo AM, Nelson WG: Reversal of GSTP1 CpG island hypermethylation and reactivation of pi-class glutathione S-transferase (GSTP1) expression in human prostate cancer cells by treatment with procainamide. *Cancer Res* 61: 8611, 2001
 56. Singal R, van Wert J, Bashambu M: Cytosine methylation represses glutathione S-transferase P1 (GSTP1) gene expression in human prostate cancer cells. *Cancer Res* 61: 4820, 2001
 57. Ng HH, Jeppesen P, Bird A: Active repression of methylated genes by the chromosomal protein MBD1. *Mol Cell Biol* 20: 1394, 2000
 58. Zhang Y, Ng HH, Erdjument-Bromage H, Tempst P, Bird A, Reinberg D: Analysis of the NuRD subunits reveals a histone deacetylase core complex and a connection with DNA methylation. *Genes Dev* 13: 1924, 1999
 59. Mielnicki LM, Ying AM, Head KL, Asch HL, Asch BB: Epigenetic regulation of gelsolin expression in human breast cancer cells. *Exp Cell Res* 249: 161, 1999
 60. Jones PL, Veenstra GJ, Wade PA, Vermaak D, Kass SU, Landsberger N, Strouboulis J, Wolffe AP: Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat Genet* 19: 187, 1998
 61. Cameron EE, Bachman KE, Myohanen S, Herman JG, Baylin SB: Synergy of demethylation and histone deacetylase inhibition in the re-expression of genes silenced in cancer. *Nat Genet* 21: 103, 1999
 62. Nan X, Ng HH, Johnson CA, Laherty CD, Turner BM, Eisenman RN, Bird A: Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393: 386, 1998
 63. Pikaart MJ, Recillas-Targa F, Felsenfeld G: Loss of transcriptional activity of a transgene is accompanied by DNA methylation and histone deacetylation and is prevented by insulators. *Genes Dev* 12: 2852, 1998
 64. Brawley OW, Knopf K, Thompson I: The epidemiology of prostate cancer part II: the risk factors. *Semin Urol Oncol* 16: 193, 1998
 65. Carter BS, Carter HB, Isaacs JT: Epidemiologic evidence regarding predisposing factors to prostate cancer. *Prostate* 16: 187, 1990
 66. Haenszel W, Kurihara M: Studies of Japanese migrants. I. Mortality from cancer and other diseases among Japanese in the United States. *J Natl Cancer Inst* 40: 43, 1968
 67. Danley KL, Richardson JL, Bernstein L, Langholz B, Ross RK: Prostate cancer: Trends in mortality and stage-specific incidence rates by racial/ethnic group in Los Angeles County, California (United States). *Cancer Causes Control* 6: 492, 1995
 68. Giovannucci E, Rimm EB, Colditz GA, Stampfer MJ, Ascherio A, Chute CC, Willett WC: A prospective study of dietary fat and risk of prostate cancer. *J Natl Cancer Inst* 85: 1571, 1993
 69. Gann PH, Hennekens CH, Sacks FM, Grodstein F, Giovannucci EL, Stampfer MJ: Prospective study of plasma fatty acids and risk of prostate cancer. *J Natl Cancer Inst* 86: 281, 1994

70. Le Marchand L, Kolonel LN, Wilkens LR, Myers BC, Hirohata T: Animal fat consumption and prostate cancer: A prospective study in Hawaii. *Epidemiology* 5: 276, 1994
71. Gross GA, Turesky RJ, Fay LB, Stillwell WG, Skipper PL, Tannenbaum SR: Heterocyclic aromatic amine formation in grilled bacon, beef and fish and in grill scrapings. *Carcinogenesis* 14: 2313, 1993
72. Lijinsky W, Shubik P: Benzo(a)pyrene and other polynuclear hydrocarbons in charcoal-broiled meat. *Science* 145: 53, 1964
73. Gann PH, Ma J, Giovannucci E, Willett W, Sacks FM, Hennekens CH, Stampfer MJ: Lower prostate cancer risk in men with elevated plasma lycopene levels: Results of a prospective analysis. *Cancer Res* 59: 1225, 1999
74. Giovannucci E, Ascherio A, Rimm EB, Stampfer MJ, Colditz GA, Willett WC: Intake of carotenoids and retinol in relation to risk of prostate cancer. *J Natl Cancer Inst* 87: 1767, 1995
75. Giovannucci E: Tomatoes, tomato-based products, lycopene, and cancer: Review of the epidemiologic literature. *J Natl Cancer Inst* 91: 317, 1999
76. Jain MG, Hislop GT, Howe GR, Ghadirian P: Plant foods, antioxidants, and prostate cancer risk: Findings from case-control studies in Canada. *Nutr Cancer* 34: 173, 1999
77. Cohen JH, Kristal AR, Stanford JL: Fruit and vegetable intakes and prostate cancer risk. *J Natl Cancer Inst* 92: 61, 2000
78. Kolonel LN, Hankin JH, Whittemore AS, Wu AH, Gallagher RP, Wilkens LR, John EM, Howe GR, Dreon DM, West DW, Paffenbarger RS Jr: Vegetables, fruits, legumes and prostate cancer: A multiethnic case-control study. *Cancer Epidemiol Biomarkers Prev* 9: 795, 2000
79. Zhang Y, Talalay P, Cho CG, Posner GH: A major inducer of anticarcinogenic protective enzymes from broccoli: Isolation and elucidation of structure. *Proc Natl Acad Sci USA* 89: 2399, 1992
80. Clinton SK: The dietary antioxidant network and prostate carcinoma. *Cancer* 86: 1629, 1999
81. Christen S, Woodall AA, Shigenaga MK, Southwell-Keely PT, Duncan MW, Ames BN: Gamma-tocopherol traps mutagenic electrophiles such as NO(X) and complements alpha-tocopherol: Physiological implications. *Proc Natl Acad Sci USA* 94: 3217, 1997
82. Freeman VL, Meydani M, Yong S, Pyle J, Wan Y, Arvizu-Durazo R, Liao Y: Prostatic levels of tocopherols, carotenoids, and retinol in relation to plasma levels and self-reported usual dietary intake. *Am J Epidemiol* 151: 109, 2000
83. Clinton SK, Emenhiser C, Schwartz SJ, Bostwick DG, Williams AW, Moore BJ, Erdman JW Jr: Cis-trans lycopene isomers, carotenoids, and retinol in the human prostate. *Cancer Epidemiol Biomarkers Prev* 5: 823, 1996
84. Clark LC, Combs GF Jr, Turnbull BW, Slate EH, Chalker DK, Chow J, Davis LS, Glover RA, Graham GF, Gross EG, Krongrad A, Lesher JL Jr, Park HK, Sanders BB Jr, Smith CL, Taylor JR: Effects of selenium supplementation for cancer prevention in patients with carcinoma of the skin. A randomized controlled trial. Nutritional Prevention of Cancer Study Group (see comments) (published erratum appears in *JAMA* 1997 May 21; 277(19): 1520). *JAMA* 276: 1957, 1996
85. Clark LC, Dalkin B, Krongrad A, Combs GF Jr, Turnbull BW, Slate EH, Witherington R, Herlong JH, Janosko E, Carpenter D, Borosso C, Falk S, Rounder J: Decreased incidence of prostate cancer with selenium supplementation: Results of a double-blind cancer prevention trial. *Br J Urol* 81: 730, 1998
86. Heinonen OP, Albanes D, Virtamo J, Taylor PR, Huttunen JK, Hartman AM, Haapakoski J, Malila N, Rautalahti M, Ripatti S, Maenpaa H, Teerenhovi L, Koss L, Virolainen M, Edwards BK: Prostate cancer and supplementation with alpha-tocopherol and beta-carotene: Incidence and mortality in a controlled trial (see comments). *J Natl Cancer Inst* 90: 440, 1998
87. Zhang Y, Kensler TW, Cho CG, Posner GH, Talalay P: Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proc Natl Acad Sci USA* 91: 3147, 1994
88. Fahey JW, Zhang Y, Talalay P: Broccoli sprouts: An exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci USA* 94: 10,367, 1997
89. Shapiro TA, Fahey JW, Wade KL, Stephenson KK, Talalay P: Chemoprotective glucosinolates and isothiocyanates of broccoli sprouts: Metabolism and excretion in humans. *Cancer Epidemiol Biomarkers Prev* 10: 501, 2001
90. Hayes RB, Pottern LM, Strickler H, Rabkin C, Pope V, Swanson GM, Greenberg RS, Schoenberg JB, Liff J, Schwartz AG, Hoover RN, Fraumeni JF Jr: Sexual behaviour, STDs and risks for prostate cancer. *Br J Cancer* 82: 718, 2000
91. Ames BN, Gold LS, Willett WC: The causes and prevention of cancer. *Proc Natl Acad Sci USA* 92: 5258, 1995
92. Roberts RO, Lieber MM, Rhodes T, Girman CJ, Bostwick DG, Jacobsen SJ: Prevalence of a physician-assigned diagnosis of prostatitis: The olmsted county study of urinary symptoms and health status among men. *Urology* 51: 578, 1998
93. Rushmore TH, King RG, Paulson KE, Pickett CB: Regulation of glutathione S-transferase Ya subunit gene expression: Identification of a unique xenobiotic-responsive element controlling inducible expression by planar aromatic compounds. *Proc Natl Acad Sci USA* 87: 3826, 1990
94. Rushmore TH, Pickett CB: Transcriptional regulation of the rat glutathione S-transferase Ya subunit gene. Characterization of a xenobiotic-responsive element controlling inducible expression by phenolic antioxidants. *J Biol Chem* 265: 14648, 1990
95. Ramos-Gomez M, Kwak MK, Dolan PM, Itoh K, Yamamoto M, Talalay P, Kensler TW: From the Cover: Sensitivity to carcinogenesis is increased and chemoprotective efficacy of enzyme inducers is lost in *nrf2* transcription factor-deficient mice. *Proc Natl Acad Sci USA* 98: 3410, 2001
96. Henderson CJ, Smith AG, Ure J, Brown K, Bacon EJ, Wolf CR: Increased skin tumorigenesis in mice lacking pi

- class glutathione S-transferases. *Proc Natl Acad Sci USA* 95: 5275, 1998
97. Nelson CP, Kidd LC, Sauvageot J, Isaacs WB, De Marzo AM, Groopman JD, Nelson WG, Kensler TW: Protection against 2-hydroxyamino-1-methyl-6-phenylimidazo[4,5-b]pyridine cytotoxicity and DNA adduct formation in human prostate by glutathione S-transferase P1. *Cancer Res* 61: 103, 2001
 98. Morgenthaler PM, Holzhauser D: Analysis of mutations induced by 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in human lymphoblastoid cells. *Carcinogenesis* 16: 713, 1995
 99. Nagaoka H, Wakabayashi K, Kim SB, Kim IS, Tanaka Y, Ochiai M, Tada A, Nukaya H, Sugimura T, Nagao M: Adduct formation at C-8 of guanine on in vitro reaction of the ultimate form of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine with 2'-deoxyguanosine and its phosphate esters. *Jpn J Cancer Res* 83: 1025, 1992
 100. Knize MG, Sinha R, Rothman N, Brown ED, Salmon CP, Levander OA, Cunningham PL, Felton JS: Heterocyclic amine content in fast-food meat products. *Food Chem Toxicol* 33: 545, 1995
 101. Davis CD, Schut HA, Snyderwine EG: Adduction of the heterocyclic amine food mutagens IQ and PhIP to mitochondrial and nuclear DNA in the liver of Fischer-344 rats. *Carcinogenesis* 15: 641, 1994
 102. Shirai T, Sano M, Tamano S, Takahashi S, Hirose M, Futakuchi M, Hasegawa R, Imaida K, Matsumoto K, Wakabayashi K, Sugimura T, Ito N: The prostate: A target for carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) derived from cooked foods. *Cancer Res* 57: 195, 1997
 103. Shirai T, Cui L, Takahashi S, Futakuchi M, Asamoto M, Kato K, Ito N: Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo [4,5-b]pyridine (PhIP) in the rat prostate and induction of invasive carcinomas by subsequent treatment with testosterone propionate. *Cancer Lett* 143: 217, 1999
 104. Stuart GR, Holcroft J, de Boer JG, Glickman BW: Prostate mutations in rats induced by the suspected human carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine. *Cancer Res* 60: 266, 2000
 105. Fields WR, Morrow CS, Doss AJ, Sundberg K, Jernstrom B, Townsend AJ: Overexpression of stably transfected human glutathione S-transferase P1-1 protects against DNA damage by benzo[a]pyrene diol-epoxide in human T47D cells. *Mol Pharmacol* 54: 298, 1998
 106. Sundberg K, Johansson AS, Stenberg G, Widersten M, Seidel A, Mannervik B, Jernstrom B: Differences in the catalytic efficiencies of allelic variants of glutathione transferase P1-1 towards carcinogenic diol epoxides of polycyclic aromatic hydrocarbons. *Carcinogenesis* 19: 433, 1998
 107. Helzlsouer KJ, Selmin O, Huang HY, Strickland PT, Hoffman S, Alberg AJ, Watson M, Comstock GW, Bell D: Association between glutathione S-transferase M1, P1, and T1 genetic polymorphisms and development of breast cancer. *J Natl Cancer Inst* 90: 512, 1998
 108. Henderson CJ, Wolf CR, Kitteringham N, Powell H, Otto D, Park BK: Increased resistance to acetaminophen hepatotoxicity in mice lacking glutathione S-transferase Pi. *Proc Natl Acad Sci USA* 97: 12,741, 2000
 109. Diah SK, Smitherman PK, Townsend AJ, Morrow CS: Detoxification of 1-chloro-2,4-dinitrobenzene in MCF7 breast cancer cells expressing glutathione S-transferase P1-1 and/or multidrug resistance protein 1. *Toxicol Appl Pharmacol* 157: 85, 1999
 110. Adler V, Yin Z, Fuchs SY, Benezra M, Rosario L, Tew KD, Pincus MR, Sardana M, Henderson CJ, Wolf CR, Davis RJ, Ronai Z: Regulation of JNK signaling by GSTp. *Embo J* 18: 1321, 1999
 111. Wang T, Arifoglu P, Ronai Z, Tew KD: Glutathione S-transferase P1-1 (GSTP1-1) Inhibits c-Jun N-terminal Kinase (JNK1) signaling through interaction with the C terminus. *J Biol Chem* 276: 20999, 2001
 112. Yin Z, Ivanov VN, Habelhah H, Tew K, Ronai Z: Glutathione S-transferase p elicits protection against H₂O₂-induced cell death via coordinated regulation of stress kinases. *Cancer Res* 60: 4053, 2000
 113. Ruscoe JE, Rosario LA, Wang T, Gate L, Arifoglu P, Wolf CR, Henderson CJ, Ronai Z, Tew KD: Pharmacologic or genetic manipulation of glutathione S-transferase P1-1 (GSTpi) influences cell proliferation pathways. *J Pharmacol Exp Ther* 298: 339, 2001
 114. Reznik G, Hamlin MH II, Ward JM, Stinson SF: Prostatic hyperplasia and neoplasia in aging F344 rats. *Prostate* 2: 261, 1981
 115. Gilardoni MB, Rabinovich GA, Oviedo M, Depiante-Depauli M: Prostate cancer induction in autoimmune rats and modulation of T cell apoptosis. *J Exp Clin Cancer Res* 18: 493, 1999
 116. Yoshizawa K, Willett WC, Morris SJ, Stampfer MJ, Spiegelman D, Rimm EB, Giovannucci E: Study of prediagnostic selenium level in toenails and the risk of advanced prostate cancer (see comments). *J Natl Cancer Inst* 90: 1219, 1998
 117. Helzlsouer KJ, Huang HY, Alberg AJ, Hoffman S, Burke A, Norkus EP, Morris JS, Comstock GW: Association between alpha-tocopherol, gamma-tocopherol, selenium, and subsequent prostate cancer. *J Natl Cancer Inst* 92: 2018, 2000
 118. Kucuk O, Sarkar FH, Sakr W, Djuric Z, Pollak MN, Khachik F, Li YW, Banerjee M, Grignon D, Bertram JS, Crissman JD, Pontes EJ, Wood DP Jr: Phase II randomized clinical trial of lycopene supplementation before radical prostatectomy. *Cancer Epidemiol Biomarkers Prev* 10: 861, 2001
 119. Chen L, Stacewicz-Sapuntzakis M, Duncan C, Sharifi R, Ghosh L, Breemen Rv R, Ashton D, Bowen PE: Oxidative DNA damage in prostate cancer patients consuming tomato sauce-based entrees as a whole-food intervention. *J Natl Cancer Inst* 93: 1872, 2001
 120. Wechter WJ, Leipold DD, Murray ED Jr, Quiggle D, McCracken JD, Barrios RS, Greenberg NM: E-7869 (R-flurbiprofen) inhibits progression of prostate cancer in the TRAMP mouse. *Cancer Res* 60: 2203, 2000

121. Nelson JE, Harris RE: Inverse association of prostate cancer and non-steroidal anti-inflammatory drugs (NSAIDs): Results of a case-control study. *Oncol Rep* 7: 169, 2000
122. Nelson WG, Wilding G: Prostate cancer prevention agent development: Criteria and pipeline for candidate chemoprevention agents. *Urology* 57: 56, 2001
123. Rountree MR, Bachman KE, Baylin SB: DNMT1 binds HDAC2 and a new co-repressor, DMAP1, to form a complex at replication foci. *Nat Genet* 25: 269, 2000
124. Kensler TW: Chemoprevention by inducers of carcinogen detoxication enzymes. *Environ Health Perspect* 105 Suppl(4): 965, 1997
125. Wang JS, Shen X, He X, Zhu YR, Zhang BC, Wang JB, Qian GS, Kuang SY, Zarba A, Egner PA, Jacobson LP, Munoz A, Helzlsouer KJ, Groopman JD, Kensler TW: Protective alterations in phase 1 and 2 metabolism of aflatoxin B1 by oltipraz in residents of Qidong, People's Republic of China. *J Natl Cancer Inst* 91: 347, 1999
126. Jacobson LP, Zhang BC, Zhu YR, Wang JB, Wu Y, Zhang QN, Yu LY, Qian GS, Kuang SY, Li YF, Fang X, Zarba A, Chen B, Enger C, Davidson NE, Gorman MB, Gordon GB, Prochaska HJ, Egner PA, Groopman JD, Munoz A, Helzlsouer KJ, Kensler TW: Oltipraz chemoprevention trial in Qidong, People's Republic of China: Study design and clinical outcomes. *Cancer Epidemiol Biomarkers Prev* 6: 257, 1997
127. Kensler TW, He X, Otieno M, Egner PA, Jacobson LP, Chen B, Wang JS, Zhu YR, Zhang BC, Wang JB, Wu Y, Zhang QN, Qian GS, Kuang SY, Fang X, Li YF, Yu LY, Prochaska HJ, Davidson NE, Gordon GB, Gorman MB, Zarba A, Enger C, Munoz A, Helzlsouer KJ: Oltipraz chemoprevention trial in Qidong, People's Republic of China: Modulation of serum aflatoxin albumin adduct biomarkers. *Cancer Epidemiol Biomarkers Prev* 7: 127, 1998
128. Shapiro TA, Fahey JW, Wade KL, Stephenson KK, Talalay P: Human metabolism and excretion of cancer chemoprotective glucosinolates and isothiocyanates of cruciferous vegetables. *Cancer Epidemiol Biomarkers Prev* 7: 1091, 1998