**Total Resources:** A = 10, B = 10, C = 10
**My ID:** 41220300573

The last two digits of my ID are 73. However, assigning "7 3" directly to the Allocation table would result in negative values in the Available resources, which is not valid in the Banker's Algorithm.
Therefore, I modified the allocation to 2 2 to ensure a valid and safe resource distribution.

| Process | Allocation | | | Max | | | Available | | | Remaining | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | A | B | C | A | B | C | A | B | C | A | B | C |
| P0 | 0 | 1 | 0 | 7 | 5 | 3 | 4 | 8 | 5 | 7 | 4 | 3 |
| P1 | 2 | 0 | 1 | 3 | 2 | 2 | 6 | 8 | 6 | 1 | 2 | 1 |
| P2 | 2 | 0 | 2 | 9 | 0 | 2 | 8 | 9 | 7 | 7 | 0 | 0 |
| P3 | 2 | 1 | 1 | 2 | 2 | 2 | 8 | 10 | 7 | 0 | 1 | 1 |
| P4 | 0 | 0 | 1 | 3 | 2 | 2 | 10 | 10 | 9 | 3 | 2 | 1 |
| | | | | | | | 10 | 10 | 10 | | | |

Safe Sequence:
P1->P3->P0->P2->P4

| Process | Allocation | | | Max | | | Available | | | Remaining | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | A | B | C | A | B | C | A | B | C | A | B | C |
| P0 | 0 | 1 | 0 | 7 | 5 | 3 | 4 | 8 | 5 | 7 | 4 | 3 |
| P1 | 2 | 0 | 1 | 3 | 2 | 2 | 6 | 8 | 6 | 1 | 2 | 1 |
| P2 | 2 | 0 | 2 | 9 | 0 | 2 | 8 | 9 | 7 | 7 | 0 | 0 |
| P3 | 2 | 1 | 1 | 2 | 2 | 2 | 8 | 9 | 8 | 0 | 1 | 1 |
| P4 | 0 | 0 | 1 | 3 | 2 | 2 | 8 | 10 | 8 | 3 | 2 | 1 |
| | | | | | | | 10 | 10 | 10 | | | |

Safe Sequence:
P1->P3->P4->P0->P2

**Total Resources:** A = 10, B = 10, C = 10
Find the value of x x so that it gives a safe sequence

| Process | Allocation | | | Max | | | Available | | | Remaining | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | A | B | C | A | B | C | A | B | C | A | B | C |
| P0 | 0 | 1 | 0 | 7 | 5 | 3 | | | | | | |
| P1 | 2 | 0 | 1 | 3 | 2 | 2 | | | | | | |
| P2 | x | 0 | x | 9 | 0 | 2 | | | | | | |
| P3 | 2 | 1 | 1 | 2 | 2 | 2 | | | | | | |
| P4 | 0 | 0 | 1 | 3 | 2 | 2 | | | | | | |
| | | | | | | | | | | | | |