

Assumption-Lean Post-Integrated Inference with Negative Control Outcomes

Jin-Hong Du^{†‡}

Kathryn Roeder^{†§}

Larry Wasserman^{†‡}

November 26, 2024

Abstract

Data integration methods aim to extract low-dimensional embeddings from high-dimensional outcomes to remove unwanted variations, such as batch effects and unmeasured covariates, across heterogeneous datasets. However, multiple hypothesis testing after integration can be biased due to data-dependent processes. We introduce a robust post-integrated inference (PII) method that adjusts for latent heterogeneity using negative control outcomes. Leveraging causal interpretations, we derive nonparametric identifiability of the direct effects, which motivates our semiparametric inference method. Our method extends to projected direct effect estimands, accounting for hidden mediators, confounders, and moderators. These estimands remain statistically meaningful under model misspecifications and with error-prone embeddings. We provide bias quantifications and finite-sample linear expansions with uniform concentration bounds. The proposed doubly robust estimators are consistent and efficient under minimal assumptions and potential misspecification, facilitating data-adaptive estimation with machine learning algorithms. Our proposal is evaluated with random forests through simulations and analysis of single-cell CRISPR perturbed datasets with potential unmeasured confounders.

Keywords: Batch correction; Confounder adjustment; Data integration; Hypothesis testing; Latent embedding; Model-free inference.

1 Introduction

In the big data era, integrating information from multiple heterogeneous sources has become increasingly crucial for achieving larger sample sizes and more diverse study populations. The applications of data integration are in a variety of fields, including but not limited to, causal inference on heterogeneous populations (Shi et al., 2023), survey sampling (Yang et al., 2020), health policy (Paddock et al., 2024), retrospective psychometrics (Howe and Brown, 2023), and multi-omics biological science (Du et al., 2022). Data integration methods have been proposed to mitigate the unwanted effects of heterogeneous datasets and unmeasured covariates, recovering the common variation across datasets. However, a critical and often overlooked question is whether reliable statistical inference can be made from integrated data. Directly performing statistical inference on integrated outcomes and covariates of interests fails to account for the complex correlation structures introduced by the data integration process, often leading to improper analyses that incorrectly assume the corrected data points are independent (Li et al., 2023).

While data integration is broadly utilized in various fields, our paper focuses on a challenging scenario with the presence of high-dimensional outcomes. Particularly in the context of genomics,

[†]Department of Statistics and Data Science, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

[‡]Machine Learning Department, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

[§]Computational Biology Department, Carnegie Mellon University, Pittsburgh, PA 15213, USA.




Figure 1: Overview of the post-integrated inference problem. **(a)** Data integration utilizes multiple outcomes $Y = (Y_1, \dots, Y_p)^\top$ and covariate X of interest to estimate the embeddings \hat{U} , and provides integrated outcomes $\tilde{Y} = (\tilde{Y}_1, \dots, \tilde{Y}_p)^\top$ for downstream analysis. **(b)** Inference on the direct associations between Y_j 's and X , and those between \tilde{Y}_j 's and X may be biased because of batch effects and observational dependency induced by data integration processes, respectively. **(c)** Post-integrated inference includes two strategies: the design-based approach that includes a batch indicator through a statistical model and the design-free approach that first estimates the latent embeddings and then treats them as extra covariates for downstream inference (the batch indicator can also be used as an observed confounder), where the latter is our focus.

experimental constraints often necessitate the collection of data in multiple batches (Luo and Wei, 2018; Luecken et al., 2022). Batch correction and data integration methods are commonly used in genomics to recover the *low-dimensional embeddings* or manifolds of each observation from the *high-dimensional outcomes*. The naive approach uses a batch indicator as a covariate in a regression model for inference, which may not be sufficient for adjusting for batch effects and unmeasured covariates (Li et al., 2023). Instead, two-step methods are commonly employed in practice as a separate data preprocessing step to produce integrated data, which can then be utilized for downstream inference. For instance, design-based methods, such as Combat (Johnson et al., 2007) and BUS (Luo and Wei, 2018), combine the batch or unknown subtype indicator into hierarchical Bayesian models and provide location and scale correction. Additionally, design-free methods, including RUV (Gagnon-Bartsch and Speed, 2012) and SVA (Leek et al., 2012) directly estimate the latent confounding factors, and users can use the estimated latent variables as extra covariates for the downstream inference. These methods apply to samples that share the same underlying biological variability, which is our focus in this paper; see Figure 1 for an illustration.

Despite different procedures and output formats, nearly all batch correction methods utilize information from multiple outcomes to estimate and align the underlying “embeddings” of observations. This approach is closely related to unmeasured confounder adjustment, particularly when each observation is viewed as a single dataset. Over the past decades, researchers have explored various methods to address unmeasured confounders in statistical analysis. In the presence of multiple outcomes, deconfounding techniques primarily employ two strategies: incorporating known




Figure 2: Histogram of t -statistics of $PTEN$ perturbation on 8320 cells and 4163 genes by four different confounder adjustment methods. The orange dashed curves represent the density of standard normal distribution. See Section 5 for more details about the methods and experiment setting.

negative control outcomes or leveraging sparsity assumptions (Wang et al., 2017; Zhou et al., 2024). Additionally, a line of research on proximal causal inference uses both negative control outcomes and/or exposures for deconfounding (Miao et al., 2018); see a review of related work in Appendix A. This paper focuses specifically on the negative control approach in the context of multiple outcomes, where the goal is to directly estimate and adjust for latent factors that may confound the treatment outcome relationships.

Mathematically, a high-dimensional outcome vector $Y \in \mathbb{R}^p$ is often related to a covariate vector $X \in \mathbb{R}^d$ and an unobserved low-dimensional latent vector $U \in \mathbb{R}^r$. Here, X includes variables such as disease status or treatment, and U , frequently referred to as the embedding vector, captures both the batch effects and the unmeasured covariates. Both of them serve as a compact representation of the outcome Y , with the dimensionality of the outcome space being significantly larger than that of the covariate and latent space, i.e., $p \gg d$ and $p \gg r$. Differences in how data are collected across datasets can result in shifts or distortions in the distribution of unobserved variable U , and can potentially affect the distribution of X as well. Our primary interest lies in the direct associations or causal relationships between the outcome Y_j and the covariate X for $j = 1, \dots, p$, after adjusting for the difference induced by unwanted variation U . When X and U are independent, the problem would be trivial because the direct effects can be estimated by regressing Y_j 's on X . However, when X and U are dependent, the direct regression approach targets the total effects and provides a biased estimate of the direct effects. Hence, proper data integration methods need to estimate U for integrating the outcomes from different sources and for multiple hypotheses testing.

Although two-step procedures are widely favored by practitioners, it is evident that the risk of making mistakes is propagated by the two steps. Specifically, the estimation of latent embeddings U and the subsequent statistical inference are both contingent on the assumptions made by their respective models. If either model is misspecified, the final inference results can be significantly biased. For instance, varying choices of hyperparameters, such as the latent dimension, can affect the accuracy of the first-stage estimation. It is, therefore, critical to understand whether such




Figure 3: Batch correction where the latent embedding U is (a) U is a mediator that contributes to the indirect effect from X to Y ; (b) a confounder that affects both X and Y ; and (c) a moderator that interacts with an independent variable X of interest to influence an outcome Y , but is not on the causal pathway.

approaches work in more general settings and how to remedy these existing post-integrated inference methods under possible misspecification.

In this paper, we rigorously investigate the validity of statistical inference on integrated data, focusing particularly on the use of negative control outcomes to ensure reliable inference. Our aim is to analyze the validity of two-step post-integrated inference under minimal assumptions about the data-generating processes. Further, we aim to provide a framework that not only ensures effective batch correction but also maintains the integrity and reliability of statistical inference, addressing two key challenges using flexible machine learning algorithms. This will allow researchers to retain the statistical power of their analyses while providing greater confidence in the validity of their inferences from integrated data.

Post-Integrated inference To demonstrate the challenges in post-integrated inference, we analyze high-throughput single-cell CRISPR data from a study on autism spectrum disorder-related gene perturbations and their effects on neuronal differentiation (Section 5). In this example, one cell can be viewed as a single dataset, where the heterogeneity among cells may not be fully explained by observed covariates. Our analysis focuses on testing nonlinear associations between 4163 genes and *PTEN* perturbation after accounting for covariates in neural development and unwanted variations from heterogeneous observations.

Figure 2 illustrates t-statistic distributions from four different methods. The unadjusted inference method yields overly conservative test statistic distributions compared to the expected $\mathcal{N}(0, 1)$ distribution. While batch correction and confounder adjustment methods produce distributions closer to the standard normal, some show anti-conservative tendencies. Importantly, only about half of the significant tests ($p\text{-value} < 0.05$) are consistent across the three confounder adjustment methods, raising concerns about their reliability. This inconsistency stems from varied model assumptions and algorithms tailored to specific data models, which may be misspecified for sparse single-cell data or due to inaccurate estimation of the number of latent factors.

The goal of this paper is to construct a robust statistical framework that uses embeddings from existing data integration methods to mitigate misspecification issues, ensuring valid statistical inference and enhancing current post-integrated inference methodologies.

Main contributions Our work makes several key contributions. First, in Section 2, we derive nonparametric identification conditions using negative control outcomes (Section 2.1), enhancing causal interpretations and forming the basis for our post-integrated inference (PII) method. In Section 2.2, introduce a robust and assumption-lean framework for post-integrated inference that effectively addresses hidden mediators, confounders, and moderators (Figure 3). This framework ensures reliable statistical inference despite possible confounding from batch effects and data heterogeneity. It eliminates confounding ambiguity (Remark 3), leverages negative control outcomes for accurate embedding estimation (Remark 4), and exhibits resilience to model misspecification,

supporting model-free inference (Remark 5).

Our second contribution in Section 3 analyzes the statistical error in target estimands using estimated embeddings. In Section 3.1, we use martingale interpretations to assess the bias caused by these embeddings. Under regularity conditions, we show in Theorem 2 that the bias of the projected target estimand with estimated embeddings is primarily determined by the L_2 -norm of the embedding estimation error, up to an invertible transformation. Furthermore, Lemma 3 shows that bias in linear models can be deterministically evaluated using the operator norm of projection matrices, regardless of latent dimensionality.

Our third main contribution, discussed in Section 3.2 and appendix E.3, involves developing efficient semiparametric inference methods for estimands with estimated covariates under both linear and nonlinear functions. These methods advance the assumption-lean approach of Vansteelandt and Dukes (2022) to accommodate multiple treatments and outcomes. Specifically, we derive finite-sample linear expansions for direct effect estimands (Theorem 4), provide a uniform concentration bound for residuals, and establish the asymptotic distribution for both linear and nonlinear effects under mild assumptions (Corollary 5, Theorem E.1) with triangular arrays. These results are essential for establishing guarantees related to multiple testing in high-dimensional outcomes (Proposition 6).

2 Post-Integrated inference

2.1 Nonparametric identification with negative control outcomes

In this section, we consider post-integrated inference with negative control outcomes. Similar to the causal inference analysis with observational data (Imbens and Rubin, 2015; Kennedy, 2022), we consider the case when the latent variable U is a confounder as in Figure 3(b). Let $\mathcal{X} \subseteq \mathbb{R}^d$ and $\mathcal{U} \subseteq \mathbb{R}^r$ be the support of X and U , respectively. We use f to denote a generic (conditional) probability density or mass function and require causal assumptions on the observational data (X, U, Y) and counterfactual outcome $Y(x)$ when X is interpreted as treatment.

Assumption 1. (i) Consistency: when $X = x$, $Y = Y(x)$.

(ii) Positivity: $f(x | u) > 0$ for all $u \in \mathcal{U}$.

(iii) Latent ignorability: $X \perp\!\!\!\perp Y(x) | U$ for all $x \in \mathcal{X}$.

Assumption 1i requires that no interference among the subjects, meaning that a subject's outcome is affected by its treatment but not by how others are treated. Assumption 1ii suggests that $X = x$ can be observed at any confounding levels of U with a positive probability. Assumption 1iii ensures that the treatment assignment is fully determined by the confounder U . These assumptions are required to estimate the counterfactual distribution of $Y(x)$ with observed variables (X, U, Y) by the g-formula $f_{Y(x)}(y) = \int f(y | u, x) f(u) du$. In our problem, because U is not observed, all information contained in the observed data is captured by $f(y, x)$, and one has to solve for $f(y, x, u)$ or equivalently $f(u | y, x)$ from the integral equation:

$$f(y, x) = \int f(y, x, u) du. \quad (1)$$

In general, the joint distribution $f(y, x, u)$ cannot be uniquely determined. With an auxiliary variable Z , the approach by Miao et al. (2023, Theorem 1) identifies the treatment effect from any admissible¹ distribution under exclusion restriction, equivalence, and completeness assumptions.

¹A joint distribution $\tilde{f}(y, x, u)$ is admissible if it conforms to the observed data distribution $f(y, x)$, that is, $f(y, x) = \int \tilde{f}(y, x, u) du$.

Because the negative control outcomes can also be viewed as a non-differentiable proxy of the confounder, their result also applies to our problem if taking $Z = Y_C$; however, when restricting to negative control outcomes, we can extend their approach on the identification of the counterfactual distributions with weaker assumptions. Now, we modify it as follows.

To present our first result on identification with negative control outcomes, we let $f(y, x, u; \alpha)$ denote a model for joint distribution indexed by a possibly infinite-dimensional parameter α , and conditional and marginal distributions are defined analogously. We require Assumption 2.

Assumption 2. The following hold for a set of control outcomes $\mathcal{C} \subset [p]$ and for any α :

- (i) (Negative control outcomes) $(Y_{\mathcal{C}^c}, X) \perp\!\!\!\perp Y_C | U$.
- (ii) (Equivalence) any $\tilde{f}(y_C, u)$ that solves $f(y_C; \alpha) = \int \tilde{f}(y_C, u; \alpha) du$ can be written as $\tilde{f}(y_C, u) = f(y_C, v^{-1}(u); \alpha)$ for some invertible but not necessarily known function v .
- (iii) (Completeness) for all $u \in \mathcal{U}$, $f(u) > 0$; for any square-integrable function g , $\mathbb{E}[g(U) | Y_C, X = x; \alpha] = 0$ almost surely if and only if $g(U) = 0$ almost surely.

The causal diagram under Assumption 2i is given by Figure 4. Assumption 2ii is a high-level assumption stating that at any level of covariates, the joint distribution of control outcomes and confounders lies in a class where each model is identified upon a one-to-one transformation of U . In contrast to Miao et al. (2023, Assumption 2 (ii)) that concern the joint distribution of (X, U, Y_C) , Assumption 2ii only requires equivalence on the joint distribution of (U, Y_C) ; though we also require an extra completeness assumption on U in Assumption 2iii for recovering an equivalent distribution of (X, U) . The completeness property plays a pivotal role in statistics (Lehmann and Scheffé, 2012). Intuitively, it precludes the degeneration of the (conditional) distributions on their supports, which guarantees the uniqueness of the solution to certain linear integral equations. At different levels of X , Assumption 2ii requires that any infinitesimal variability in U is accompanied by variability in Y_C , which implicitly requires the dimension of Y_C to be larger than the one of U . The completeness is viewed as a regularity condition, and more detailed discussions can be found in Miao et al. (2023, Appendix 2). Building upon the approach by Miao et al. (2023), we propose a modified identification approach.

Theorem 1 (Nonparametric identification). *Under Assumptions 1 and 2, for any admissible distribution $\tilde{f}(y_C, u)$ that solves $f(y_C) = \int \tilde{f}(y_C, u) du$ and let $\tilde{f}(u) := \int \tilde{f}(y_C, u) dy_C$, there exist a unique solution $\tilde{f}(x | u)$ to the equation*

$$f(x) = \int \tilde{f}(x | u) \tilde{f}(u) du. \quad (2)$$

Let $\tilde{f}(y_C, u | x) := \tilde{f}(y_C, u) \tilde{f}(x | u) / f(x)$, then there exists a unique solution $\tilde{f}(y_C^c | x, u)$ to the equation

$$f(y | x) = \int \tilde{f}(y_C^c | x, u) \tilde{f}(y_C, u | x) du, \quad (3)$$

Further, the potential outcome distribution is identified by

$$f_{Y(x)}(y) = \int \tilde{f}(y_C^c | u, x) \tilde{f}(y_C, u) du.$$




Figure 4: Causal diagram with negative control outcomes Y_C , from which an embedding function $Y_C \mapsto U$ can be estimated.

Theorem 1 suggests that if the joint distribution of (Y_C, U) can be estimated up to inverse transformation, then one can recover the joint distribution of potential outcome $Y(x)$. Based on Theorem 1, an operational strategy is given in two steps. The first step is to derive $\tilde{f}(y_C, u)$, which retrieves a proxy of U using the information from multiple control outcomes Y_C . Given $\tilde{f}(y_C, u)$, the conditional treatment distribution $\tilde{f}(x | u)$ and the condition outcome distribution can be obtained by solving integral equations (2) and (3). Even though $\tilde{f}(y_C, u)$ might not be unique, the estimated condition distributions $\tilde{f}(x | u)$ and $\tilde{f}(y_{C^c} | x, u)$ are guaranteed to be unique for any given $\tilde{f}(y_C, u)$. Motivated by the nonparametric identification condition presented in Theorem 1, we will provide a detailed description of the deconfounding strategy for recovering the true main effect under more relaxed assumptions in the next subsection.

Remark 1 (Deconfounding with negative control outcomes). The deconfounding strategy given in Theorem 1 is similar to previous negative control outcome approaches (Wang et al., 2017; Zhou et al., 2024) under parametric modeling assumptions but somewhat different from Miao et al. (2023, Theorem 1) under nonparametric modeling assumptions. More specifically, Theorem 1 of Miao et al. (2023) aims to recover the joint distribution of three variables (Z, X, U) , where Z is an auxiliary variable that satisfies exclusion restriction condition $Z \perp\!\!\!\perp Y_{C^c} | (X, U)$. When Z is negative control outcome Y_C , we are able to factorize the joint distribution into two conditional distributions of $X | U$ and $Y_C | U$. This property allows us to derive nonparametric identification with weaker assumptions in Theorem 1.

Another related approach is the proximal causal inference framework that uses both negative control outcomes and negative control exposures (Miao et al., 2024). The key to their method is a bridge function $b(Y_C, a)$ such that

$$p(Y_{C^c} | U, A = a) = \int b(y_C, a)p(y_C | U, A = a) dy_C = \int b(y_C, a)p(y_C | U) dy_C.$$

If the bridge function b is known, then the counterfactual distributions of $Y_{C^c}(a)$ can be recovered under classical causal assumptions. The proximal causal inference framework aims to bypass the estimation of the unmeasured confounders by estimating the bridge function using other extra information (e.g. negative control exposures), while our strategy relies on multiple control outcomes to estimate the distribution of confounders (up to invertible transformation) directly. With multiple negative control outcomes as in Figure 4, one can also split these outcomes into two nonoverlapping sets to serve the role of negative control outcomes and exposures in order to apply the proximal causal inference method; however, our approach avoids the splitting.

Remark 2 (Deconfounding with multiple treatments). When there is a single outcome, and the information of confounders solely comes from multiple treatments, we can marginalize the unknown conditional distribution $f(u | y, x)$ over the response y to obtain $f(u | x) = \int f(u | y, x)f(y | x) dy$. This suggests a two-stage procedure as in Miao et al. (2023), for successively identifying solutions $f(u, x)$ and $f(y | u, x)$ from two integral equations: $f(x) = \int f(u, x) du$ and $f(y | x) = \int f(y | u, x)f(u | x) du$. The information used to estimate the confounders in their setting is from multiple null treatments instead of multiple outcomes. For this reason, they require strong assumptions to distinguish the set of confounded treatments associated with confounders.

2.2 Assumption-Lean semiparametric inference

The nonparametric identification results aim to reveal the counterfactual distributions from confounded observational data, which is useful for designing general deconfounding strategies, yet remains impractical. When restricted to semiparametric models, however, one can design more efficient estimation and inferential procedures. A leading example of semiparametric regression models is the partially linear regression (Robinson, 1988; Härdle et al., 2000):

$$\mathbb{E}[Y | X, U] = \beta^\top X + h(U), \quad (4)$$

where Y is a high-dimensional vector of response, X is a low-dimensional vector of covariates (including the treatment of interest), $U \in \mathbb{R}^r$ is a low-dimensional latent vector, i.e., an unmeasured confounder, $\beta \in \mathbb{R}^{d \times p}$ is the coefficient to be estimated, and $h : \mathbb{R}^r \rightarrow \mathbb{R}^p$ is an unknown function. In the past decades, much attention has been on estimating and testing partially linear models.

When U is known, the coefficient β can be obtained with the double residual methodology (Robinson, 1988), by noting that

$$\mathbb{E}[Y | X, U] - \mathbb{E}[Y | U] = \beta^\top (X - \mathbb{E}[X | U]),$$

More specifically, the double residual methodology proceeds in two steps: (1) regressing Y on U to obtain the residual $Y - \widehat{\mathbb{E}}[Y | U]$, and regress X on U to obtain the residual $X - \widehat{\mathbb{E}}[X | U]$; and (2) regressing the residual $Y - \widehat{\mathbb{E}}[Y | U]$ on the residual $X - \widehat{\mathbb{E}}[X | U]$. Here, the notation $\widehat{\mathbb{E}}$ denotes the estimated regression function. The resulting regression coefficient is an estimator of β . Intuitively, this procedure removes the confounding effect of U by taking the residuals, so that the final regression only captures the relationship between X and Y conditional on U , which is β under the partial linear model assumption. In the special case with binary treatments, the resulting estimator is called E-estimator (Robins et al., 1992).

Even when the model (4) is misspecified, the estimator from the two-step procedure is directly informative about the conditional association between X and U . Under mild moment assumptions on the conditional covariance matrix of X given U , it returns a meaningful estimand

$$\begin{aligned} \beta &= \mathbb{E}[\text{Cov}(X | U)]^{-1} \mathbb{E}[\text{Cov}(X, \mathbb{E}[Y | X, U] | U)] \\ &= \mathbb{E}[\text{Cov}(X | U)]^{-1} \mathbb{E}[\text{Cov}(X, Y | U)], \end{aligned} \tag{5}$$

which itself does not crucially rely on the restrictions imposed by the outcome model (4).

Remark 3 (Relaxation of causal relationship). Under the causal setting in Section 2.1, when U is not a confounder but a moderator as in Figure 3, adjusting for U can also help to reduce the variance. If U is a confounder, it is necessary to adjust for U to have a proper interpretation of the main effect of X on Y . However, when U is missing, in general, we will not be certain whether U is a confounder or not. In particular, each entry of U can either be a confounder, a mediator, or a moderator (as in Figure 5). When targeting the estimand (5), we do not need to impose specific causal assumptions. In contrast, (5) allows us to relax the relationship between U and X , as long as the variability of X given U persists.

In summary, statistical inference targeting at projected direct effect (5) is model-free and assumption-lean. Because U is unmeasured, we rely on the strategy offered by Theorem 1 to estimate and perform inference with negative control outcomes. Our deconfounding procedure is summarized in Algorithm 1 for general link functions. Below, we describe the main steps of the procedure with an identity link as a special case.

(1) Reduction Suppose that $\mathcal{C} \subseteq [p]$ is the set of negative control outcomes such that $\beta_{\mathcal{C}} = 0$. In the first step, we aim to estimate U from the negative control outcomes $Y_{\mathcal{C}}$ independently of X . To distinguish from the previous causal setting, we call U as the embedding of $Y_{\mathcal{C}}$. This typically involves learning some (nonlinear) embedding map $f_e : \mathbb{R}^{|\mathcal{C}|} \rightarrow \mathbb{R}^r$ with $Y_{\mathcal{C}} \mapsto U$.

One can use the same set of data to learn the embedding function \widehat{f}_e and obtain the transformed embedding $\widehat{U} = \widehat{f}_e(Y_{\mathcal{C}})$. For example, perform the principle component analysis and use the first few principal components as the estimation embedding \widehat{U} . In a more general scenario, we can also borrow extra datasets to estimate the embedding function. For genomic studies, many single-cell atlas of healthy cells can be used to estimate it, which helps to improve the estimation of latent embedding and is commonly used in practice for transfer learning (Hao et al., 2021).




Figure 5: The causal relationship between X and U in Figure 4 can be further relaxed.

Algorithm 1 Post-Integrated inference (PII) with negative control outcomes

Input: A data set \mathcal{D} that contains N i.i.d. samples of $(X, Y) \in \mathbb{R}^d \times \mathbb{R}^p$, a set of control genes $\mathcal{C} \subset [p]$, and a user-specified link function g .

- 1: (Optional) Split sample $\mathcal{D} = \mathcal{D}_0 \cup \mathcal{D}_1$ with $|\mathcal{D}_0| = m$, $|\mathcal{D}_1| = n$ and $N = m+n$; otherwise set $\mathcal{D} = \mathcal{D}_0 = \mathcal{D}_1$ and $N = m = n$.
- 2: **Estimation of the embedding functional:** Based on samples in \mathcal{D}_0 , obtain an estimate $\hat{f}_e : \mathbb{R}^{|\mathcal{C}|} \rightarrow \mathbb{R}^r$ for the embedding map $f_e : Y_{\mathcal{C}} \mapsto U$.
- 3: **Extract estimated latent embeddings:** Compute $\hat{U} = \hat{f}_e(Y_{\mathcal{C}})$ on \mathcal{D}_1 .
- 4: **Semiparametric inference of the main effect estimand:** Use Algorithm E.3 to estimate

$$\tilde{\beta}_{\cdot j} = \mathbb{E}[\text{Cov}(X | \hat{U})]^{-1} \mathbb{E}[\text{Cov}(X, g(\mathbb{E}[Y_j | X, \hat{U}] | \hat{U}))], \quad j \in \mathcal{C}^c$$

and the empirical variance. Construct the confidence interval or compute p-values according to the asymptotic distribution of $\tilde{\beta}$.

Output: Return the confidence intervals or p-values.

Remark 4 (Negative control genes). For genomic studies, housekeeping genes can serve as negative control outcomes. Furthermore, even though most of the genes are measured, typically only the top thousands of highly variable genes are used for the subsequent differential expression testing. It is believed that the remaining genes with low expression behave similarly under different experimental conditions. As we demonstrate later in Section 5, we can ideally utilize these extra genes as pseudo-negative control outcomes to improve statistical inference. Of course, there are chances that some of the genes with low expression are indeed affected by the conditions; our framework would still provide reasonable interpretability as well as robustness against such misspecification of the negative controls.

(2) Estimation In the second stage, recall that our target estimand is β in (5). Because U is unobserved, the best we can do is to use \hat{U} as the estimated embedding and focus on the estimand:

$$\tilde{\beta}_{\cdot j} = \mathbb{E}[\text{Cov}(X | \hat{U})]^{-1} \mathbb{E}[\text{Cov}(X, Y_j | \hat{U})], \quad j \in \mathcal{C}^c. \quad (6)$$

This estimand quantifies the conditional associations of X and Y given \hat{U} . One would typically restrict the estimation of main effects to the complement set of control genes \mathcal{C}^c , while for notational simplicity, we simply set $\tilde{\beta}_{\cdot C} = 0_{d \times |\mathcal{C}|}$ and present the main effect matrix $\tilde{\beta} \in \mathbb{R}^{d \times p}$ in its whole. Note that for $j \in \mathcal{C}$, one always has $\tilde{\beta}_{\cdot j} = 0_d$, because $\mathbb{E}[Y_j | X, U] = \mathbb{E}[Y_j | U]$ does not depend on X and the conditional covariance between X and $\mathbb{E}[Y_j | X, U]$ is always zero.

(3) Inference In the last step, to provide uncertainty quantification, we rely on the efficient influence function for $\tilde{\beta}$, similar to E-estimator (Chernozhukov et al., 2018) and two-stage least squares estimators (Robins et al., 1992; Vansteelandt and Dukes, 2022). The details of semiparametric inference will be given later in Section 3.2 and Appendix E.3 for linear and nonlinear link functions, respectively.

Remark 5 (Assumption-lean and model-free inference). The above procedure is minimally dependent on assumptions regarding the data-generating process. It operates independently of any underlying data model, making it truly model-free. To compute an estimate of (6), arbitrary nonparametric methods can be employed to estimate the nuisance regression function. Inference can then be performed using the efficient influence function within the semiparametric framework (Vansteelandt and Dukes, 2022). As we will see in the next section, this approach only requires mild moment conditions on the true regression function and consistency assumptions on the nuisance function estimation.

The procedure is straightforward and easy to understand. However, caution is warranted for nuisance regression functions and variance estimation (Vansteelandt and Dukes, 2022). To understand the exact conditions under which this method is effective, a more sophisticated analysis is required to quantify the bias using estimated latent embeddings. Additionally, theoretical guarantees of valid inference need to take into account the presence of multivariate covariates and multiple outcomes. The next section serves these purposes.

3 Statistical properties with estimated embeddings

3.1 Bias of main effects

Before presenting our analysis of the estimation errors, we introduce several technical assumptions. To begin with, we consider a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and let \widehat{U}_m explicitly indicate the dependency of \widehat{U} on $m \in \mathbb{N}$, which is the sample size used to estimate the embedding functional \widehat{f}_e . In general, \widehat{U}_m can have different dimensions than U ; to ease our theoretical analysis, we will treat the latent dimension r as known so that $\widehat{U}_m \in \mathbb{R}^r$. As we will see later, such a requirement can be weakened under certain working models. Let $\{\mathcal{F}_m\}_{m \in \mathbb{N}}$ be a filtration generated by $\{\widehat{U}_m\}_{m \in \mathbb{N}}$ such that $\mathcal{F}_m = \sigma(\widehat{U}_m)$ and $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \dots$, and define the sub- σ -field $\mathcal{F}_\infty = \sigma(\cup_m \mathcal{F}_m) \subseteq \mathcal{F}$. We require the following assumption.

Assumption 3 (Latent embedding estimation). There exists a \mathcal{F}_∞ -measurable and invertible function v such that $\widehat{U}_m \xrightarrow{\text{a.s.}} v(U)$. Further, $\ell_m := \|\widehat{U}_m - v(U)\|_{L_2} < \infty$.

In many scenarios when we have prior information on the embedding function f_e , both the number of latent dimensions and the embedding can be consistently estimated. For example, consistent estimation of the number of latent variables has been well established under factor models (Bai and Ng, 2002) and under mixture models (Chen et al., 2012). Generally, a rate of $\ell_m = \mathcal{O}_{\mathbb{P}}(m^{-\frac{1}{2}})$ can be obtained for factor analysis when there are sufficient many negative control outcomes such that $|\mathcal{C}| > m$ (Bai and Li, 2012). For mixture models, this reduces estimating the cluster membership because one can treat the one hot vector of cluster memberships as the embedding and the cluster centers as the loading, akin to factor analysis. When f_e is estimated nonparametrically by \widehat{f}_e , the estimated embedding \widehat{U}_m can be viewed as nonparametrically generated covariates. In this context, Assumption 3 only requires the (conditional) L_2 -norm of the estimation error $\widehat{f}_e - f_e$ decays to zero in probability to ensure meaningful and accurate estimation of U , which is weaker than Assumption 2 of Mammen et al. (2012) that requires the (conditional) L_∞ -norm of $\widehat{f}_e - f_e$ is $o_{\mathbb{P}}(1)$. Finally, we also remark that one can use extra data sources to obtain a better estimate of \widehat{f}_e with a larger sample size m . In many applications, such as single-cell data analysis, the embedding function can be derived from previous alas studies so that m will be sufficiently large enough.

The following Assumption 4 imposes boundedness condition on the population quantities, and Assumption 5 imposes smoothness assumption on the regression function.

Assumption 4 (Regularity conditions). There exists constants $\bar{\sigma} \geq \sigma > 0$ and $M > 0$ such that $\sigma I_d \preceq \mathbb{E}[\text{Cov}(X | U)] \preceq \bar{\sigma} I_d$, $\sigma I_d \preceq \mathbb{E}[\text{Cov}(X | \widehat{U}_m)]$, $\|\beta\|_{2,\infty} \leq M$, $\|X\|_{L_2} \leq M$, $\max_{j \in \mathcal{C}^c} \|Y_j\|_{L_2} \leq M$.

Assumption 5 (Lipschitzness of regression functions). The regression functions satisfy Lipschitz conditions:

$$\begin{aligned} \|\mathbb{E}[X | U = u_1] - \mathbb{E}[X | U = u_2]\| &\leq L_X \|u_1 - u_2\| \\ \|\mathbb{E}[Y_j | X, U = u_1] - \mathbb{E}[Y_j | X, U = u_2]\| &\leq L_Y \|u_1 - u_2\|, \quad \forall j \in \mathcal{C}^c, \end{aligned}$$

almost surely for all $u_1, u_2 \in \mathcal{U}$ and some constants L_X and L_Y .

Assumption 5 imposes certain smoothness restrictions on the conditional expectation. In certain applications, the Lipschitz condition holds for many continuous multivariate distributions. For example, suppose W and V are jointly normally distributed with

$$\begin{pmatrix} W \\ V \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} \mu_W \\ \mu_V \end{pmatrix}, \begin{pmatrix} \Sigma_W & \Sigma_{WV} \\ \Sigma_{WV}^\top & \Sigma_V \end{pmatrix} \right).$$

Then $h(v) = \mathbb{E}[W \mid V = v] = \mu_W + \Sigma_{WV}\Sigma_V^{-1}(v - \mu_V)$ is L -Lipschitz in ℓ_2 -norm, with $L = \|\Sigma_{WV}\Sigma_V^{-1}\|$. Other examples of such a regression function include the posterior mean of the exponential and Poisson distributions under their conjugate prior, as in Bayesian inference. Similar conditions have been employed for nonparametric regression with generated covariates; see, for example, Assumption 4 in Mammen et al. (2012). In particular, Mammen et al. (2012) require differentiability and Lipschitz condition in ℓ_∞ of the condition expectation, which is much stronger than Assumption 5.

Consider two population coefficients β and $\tilde{\beta}$ as defined in (5) and (6), respectively. We next quantify the difference between the two in Theorem 2.

Theorem 2 (Bias for two-stage regression with estimated covaraites). *Under Assumptions 3–5, when $\|\mathbb{E}[X \mid \hat{U}] - \mathbb{E}[X \mid U]\|_{\ell_2} < \sigma/(2M)$, it holds that*

$$\max_{j \in \mathcal{C}^c} \|\tilde{\beta}_{\cdot j} - \beta_{\cdot j}\| \lesssim \left(\|X\|_{\ell_2} (L_X^{\frac{1}{2}} + L_Y^{\frac{1}{2}}) + \max_{j \in \mathcal{C}^c} \|Y_j\|_{\ell_2} L_Y^{\frac{1}{2}} \right) \ell_m.$$

Theorem 2 suggests that the upper bound of estimation error using estimated embeddings is related to the second moments of X and Y , as well as the accuracy of latent embedding estimation. This deterministic result only concerns the population quantities. Given i.i.d. samples of (X, U, Y) , the corresponding estimator of $\beta_{\cdot j}$ based on finite samples is given by

$$b_{\cdot j} = (\mathbb{P}_n\{(X - \hat{\mathbb{E}}[X \mid U])^{\otimes 2}\})^{-1} \mathbb{P}_n\{(X - \hat{\mathbb{E}}[X \mid U])(Y_j - \hat{\mathbb{E}}[Y_j \mid U])\}, \quad (7)$$

where $A^{\otimes 2} := AA^\top$ denotes Gram matrix of A^\top , and $\hat{\mathbb{E}}[X \mid U]$ and $\hat{\mathbb{E}}[Y \mid U]$ are the estimated nuisance functions. Because U is unobserved, we treat \hat{U} as the truth and estimate $\tilde{\beta}_{\cdot j}$ with:

$$\tilde{b}_{\cdot j} = (\mathbb{P}_n\{(X - \hat{\mathbb{E}}[X \mid \hat{U}])^{\otimes 2}\})^{-1} \mathbb{P}_n\{(X - \hat{\mathbb{E}}[X \mid \hat{U}])(Y_j - \hat{\mathbb{E}}[Y_j \mid \hat{U}])\}. \quad (8)$$

As an example, we consider a special case when the regression functions are linear models. To distinguish from previous notations, we use bold font to indicate the latent embedding matrix $\mathbf{U} \in \mathbb{R}^{n \times r}$ and its estimate $\hat{\mathbf{U}} \in \mathbb{R}^{n \times \hat{r}}$, where the latter may have a different dimension \hat{r} than the truth r . Lemma 3 below shows that we are still able to quantify the empirical estimation error of the main effects in terms of the estimation error of linear projection matrices in finite samples.

Lemma 3 (Empirical bias with estimated embeddings under linear models). *Define $S = \mathbb{P}_n\{(X - \mathbb{E}[X \mid U])^{\otimes 2}\}$, $\tilde{S} = \mathbb{P}_n\{(X - \mathbb{E}[X \mid \hat{U}])^{\otimes 2}\}$, and $\Gamma = \text{diag}(\mathbb{P}_n\{YY^\top\})$. Assume S and \tilde{S} have full rank, and $\kappa(S)\|P_{\hat{\mathbf{U}}}^\perp - P_{\mathbf{U}}^\perp\| < 1$, where for any matrix $A \in \mathbb{R}^{n \times p}$, $P_A = A(A^\top A)^{-1}A^\top$ denotes the projection matrix and $\kappa(A) = \|A\|\|A^{-1}\|$ denotes the condition number of matrix A . When $\hat{\mathbb{E}}[X \mid U]$, $\hat{\mathbb{E}}[X \mid \hat{U}]$, and $\hat{\mathbb{E}}[Y \mid U]$ are linear functions, it holds that*

$$\max_{j \in \mathcal{C}^c} \|\tilde{b}_{\cdot j} - b_{\cdot j}\| \leq \left(\|b\|_{2,\infty} + \|S\|_{\text{op}}^{-\frac{1}{2}} \|\Gamma\|_\infty \right) \frac{\kappa(S)\|P_{\hat{\mathbf{U}}}^\perp - P_{\mathbf{U}}^\perp\|}{1 - \kappa(S)\|P_{\hat{\mathbf{U}}}^\perp - P_{\mathbf{U}}^\perp\|},$$

where $\|A\|_{2,\infty} = \max_{j \in [p]} \|A_{\cdot j}\|$ is the maximum column euclidean norm for matrix $A \in \mathbb{R}^{n \times p}$.

Compared to Theorem 2, Lemma 3 suggests that the rate condition of \widehat{U} can be weakened to the rate condition of the linear projection $P_{\widehat{U}}^\perp$. The conclusion of Lemma 3 is fully deterministic and its proof relies on the backward error analysis in numerical linear algebra (Trefethen and Bau, 2022). The dimension of the estimated embedding is allowed to differ from the truth, as long as the column space of \widehat{U} captures essential information of the column space of U . Analogously, it is possible to relax Assumption 3 to varying latent dimension settings for Theorem 2 under general data models. In this regard, one can consider a decomposition of $\lim_m \widehat{U}_m = T + A$, where T and A are a sufficient statistic and an ancillary statistic, respectively, as when U is viewed as a parameter. We leave such an extension as future work.

3.2 Doubly robust semiparametric inference

In the previous section, we showed that the target estimands $\tilde{\beta}$ and β are similar whenever \widehat{U} is consistent to U up to any invertible transformation. Based on the estimated embedding \widehat{U} , our target of estimation and inference becomes $\tilde{\beta}$ as defined in (6). To consider potential nonparametric models for the nuisance functions, in what follows, we require the estimated nuisance functions $\widehat{\mathbb{E}}[X | U]$ and $\widehat{\mathbb{E}}[Y | U]$ to be computed from independent samples of \mathbb{P}_n . The required independence is very standard in recent developments of double machine learning and causal inference (Vansteelandt and Dukes, 2022; Kennedy, 2022), because sample splitting and cross-fitting can be used to fulfill this requirement, though one can also restrict to Donsker classes to avoid sample splitting (Kennedy, 2022).

Before we inspect the estimation error of \tilde{b} to the target estimand $\tilde{\beta}$, we introduce one extra assumption on the moments and consistency of nuisance estimation.

Assumption 6 (Bounded moments and consistency). There exists $\delta \in (0, 1]$, $M > 0$, such that

$$\begin{aligned} \|X - \mathbb{E}[X | \widehat{U}]\|_{L_{2(1+\delta^{-1})}} &\vee \|X - \widehat{\mathbb{E}}[X | \widehat{U}]\|_{L_{2(1+\delta^{-1})}} \vee \|Y - \mathbb{E}[Y | \widehat{U}]\|_{L_{2(1+\delta^{-1})}} < M, \\ \|\mathbb{E}[X | \widehat{U}] - \widehat{\mathbb{E}}[X | \widehat{U}]\|_{L_{2(1+\delta)}} &, \|\|\mathbb{E}[Y | \widehat{U}] - \widehat{\mathbb{E}}[Y | \widehat{U}]\|_\infty\|_{L_{2(1+\delta)}} = o_{\mathbb{P}}(1). \end{aligned}$$

Let $O = (X, \widehat{U}, Y) \in \mathbb{R}^d \times \mathbb{R}^r \times \mathbb{R}^p$ denote the observation when the estimated embedding function \widehat{f}_e is treated as fixed. The following theorem shows the linear expansion of the estimator \tilde{b} and gives the error bound of the residual term with high probability.

Theorem 4 (Linear expansion). *Consider the above inferential procedure, suppose Assumptions 4 and 6 hold and two nuisance functions $\widehat{\mathbb{E}}[X | \widehat{U}]$ and $\widehat{\mathbb{E}}[Y | \widehat{U}]$ are estimated from independent samples of \mathbb{P}_n . Then, the estimator \tilde{b} admits a linear expansion:*

$$\sqrt{n}(\tilde{b} - \tilde{\beta}) = \sqrt{n}\tilde{\Sigma}^{-1}(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}(O; \mathbb{P})\} + \xi,$$

where $\tilde{\Sigma} := \mathbb{E}[\text{Cov}(X | \widehat{U})]$ and $\tilde{\varphi}$ is the influence function of $\tilde{\Sigma}\tilde{\beta}$ defined as

$$\tilde{\varphi}(O; \mathbb{P}) := (X - \mathbb{E}[X | \widehat{U}])(((Y - \mathbb{E}[Y | X]) - \tilde{\beta}^\top(X - \mathbb{E}[X | \widehat{U}]))^\top. \quad (9)$$

For any $\epsilon > 0$, there exists a constant $C = C(\epsilon, \sigma, M, L)$, such that with probability at least $1 - \epsilon$, the remainder term ξ satisfies that

$$\begin{aligned} \|\xi\|_{2,\infty} &\leq C\{\|(\mathbb{P}_n - \mathbb{P})\{(X - \mathbb{E}[X | \widehat{U}])^{\otimes 2}\}\|_{\text{op}} \\ &\quad + \|\mathbb{E}[X | \widehat{U}] - \widehat{\mathbb{E}}[X | \widehat{U}]\|_{L_{2(1+\delta)}} + \|\|\mathbb{E}[Y | \widehat{U}] - \widehat{\mathbb{E}}[Y | \widehat{U}]\|_\infty\|_{L_{2(1+\delta)}}\} \\ &\quad + C\sqrt{n}\{\|\mathbb{E}[X | \widehat{U}] - \widehat{\mathbb{E}}[X | \widehat{U}]\|_{L_2}^2 \\ &\quad + ML\|\mathbb{E}[Y | \widehat{U}] - \widehat{\mathbb{E}}[Y | \widehat{U}]\|_{L_2, \infty}^2 \\ &\quad + \|\mathbb{E}[Y | \widehat{U}] - \widehat{\mathbb{E}}[Y | \widehat{U}]\|_{L_2, \infty}\|\mathbb{E}[X | \widehat{U}] - \widehat{\mathbb{E}}[X | \widehat{U}]\|_{L_2}\}. \end{aligned}$$

Theorem 4 provide a non-asymptotic uniform error bound for the residual terms over multiple outcomes. With the law of large numbers and the consistency in Assumption 6, we know that the first term of the upper bound is $o_{\mathbb{P}}(1)$. On the other hand, the secondary term is also negligible under specific rate conditions on the estimation errors of nuisances. Considering an asymptotic regime when viewing m and p as sequences indexed by n and $n, m, p \rightarrow \infty$, the above result suggests the asymptotic normality, as presented in the following corollary.

Corollary 5 (Doubly robust inference with estimated emebeddings). *Under conditions in Theorem 4, if further, the estimation error rates of nuisance functions satisfy that $\|\mathbb{E}[X | \hat{U}] - \hat{\mathbb{E}}[X | \hat{U}]\|_{L_2}^2 = o_{\mathbb{P}}(n^{-\frac{1}{2}})$, $\|\mathbb{E}[Y | \hat{U}] - \hat{\mathbb{E}}[Y | \hat{U}]\|_{L_2, \infty}^2 = o_{\mathbb{P}}(n^{-\frac{1}{2}})$, $\|\mathbb{E}[Y | \hat{U}] - \hat{\mathbb{E}}[Y | \hat{U}]\|_{L_2, \infty} \|\mathbb{E}[X | \hat{U}] - \hat{\mathbb{E}}[X | \hat{U}]\|_{L_2} = o_{\mathbb{P}}(n^{-\frac{1}{2}})$, then the estimator \tilde{b} is asymptotically normal:*

$$\sqrt{n}(\tilde{b}_{\cdot j} - \tilde{\beta}_{\cdot j}) \xrightarrow{d} \mathcal{N}_d(0, \tilde{\Sigma}^{-1} \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} \tilde{\Sigma}^{-1}), \quad j = 1, \dots, p.$$

Furthermore, if the conditions of Theorem 2 hold with $\ell_m = o(n^{-\frac{1}{2}})$, then we have

$$\sqrt{n}(\tilde{b}_{\cdot j} - \beta_{\cdot j}) \xrightarrow{d} \mathcal{N}_d(0, \tilde{\Sigma}^{-1} \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} \tilde{\Sigma}^{-1}), \quad j = 1, \dots, p.$$

In the presence of estimated embedding \hat{U}_m , the influence function $\tilde{\varphi}$ implicitly depends on the sample size m . Therefore, establishing the asymptotic normality requires verification of the Lindeberg condition for triangular array of random variables. In Corollary 5, the rate of estimation for the two nuisance functions may be slower than the parametric rate $n^{-\frac{1}{2}}$, as long as each individual estimation rate is faster than $n^{-\frac{1}{4}}$. This flexibility enables us to employ more versatile machine learning algorithms for nuisance function estimation while maintaining the validity of our inference. Furthermore, Corollary 5 suggests that efficient influence regarding the true main effect β is possible when the rate of consistently estimating the embedding is $\ell_m = o_{\mathbb{P}}(n^{-\frac{1}{2}})$. As discussed above, under factor models, one has $\ell_m = \mathcal{O}_{\mathbb{P}}(m^{-\frac{1}{2}})$, this requires $n = o(m)$, i.e., the factor loadings need to be estimated from more observations than those used for the estimation and inference of b .

Based on Corollary 5, the data-adaptive procedure to obtain the confidence intervals and p-values is given in Algorithm 2. To fulfill the independence assumptions, one can use cross-fitting to ensure that different samples are used for step 1 and step 2. When this holds, the following proposition shows that overall Type-I error control can be controlled at the desired level. In Proposition 6, when the unit vector v is chosen to be the basis vector, it reduces to testing whether a specific covariate has zero association with individual outcomes.

Proposition 6 (Multiple linear hypothesis testing). *Let $t_j = \sqrt{n} \mathbb{V}_n \{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\}^{\frac{1}{2}} \tilde{\Sigma}^{-1} (\tilde{b}_{\cdot j} - \tilde{\beta}_{\cdot j})$ be the standardized vector. For any unit vector $v \in \mathbb{R}^d$, consider the hypothesis $\mathcal{H}_{0j} : v^\top \beta_{\cdot j} = 0$. Let $\mathcal{N}_p = \{j \mid v^\top \beta_{\cdot j} = 0, j = 1, \dots, p\}$ be the true null hypotheses. Under the assumptions of Corollary 5, as $m, n, p, |\mathcal{N}_p| \rightarrow \infty$ such that $\ell_m = o(n^{-1/2})$, it holds that $|\mathcal{N}_p|^{-1} \sum_{j \in \mathcal{N}_p} \mathbf{1}\{|v^\top t_j| > z_{\frac{\alpha}{2}}\} \xrightarrow{p} \alpha$.*

Remark 6 (Multiple testing). The condition $\|\|\mathbb{E}[Y | \hat{U}] - \hat{\mathbb{E}}[Y | \hat{U}]\|_\infty\|_{L_2(1+\delta)} = o_{\mathbb{P}}(1)$ in Assumption 6 controls the envelope of the regression function estimation errors. This is useful when the number of outcomes p grows in the number of sample size n , when multiple testing procedures based on multiplier bootstrap can be applied to control both the family-wise error rate and the false discovery rate (Du et al., 2024b). Alternatively, one can simply apply the Benjamini–Hochberg procedure for multiple testing corrections.

Algorithm 2 Semiparametric inference for main effects

Input: Responses Y , covariate X , and estimated latent embedding \hat{U} .

- 1: Use machine learning methods to obtain nuisance estimates $\hat{\mathbb{E}}[Y | \hat{U}]$ and $\hat{\mathbb{E}}[X | \hat{U}]$.
- 2: Fit a linear regression of $Y - \hat{\mathbb{E}}[Y | \hat{U}] \sim X - \hat{\mathbb{E}}[X | \hat{U}]$ without an intercept to obtain an estimate \tilde{b} as defined in (8) of $\tilde{\beta}$ as defined in (8).
- 3: Estimate the variance of $\tilde{b}_{j,i}$ by \hat{S}_j/n based on Theorem 4, where $\hat{S}_j = \hat{\Sigma}^{-1}\mathbb{V}_n\{\tilde{\varphi}_j(O; \hat{\mathbb{P}})\}\hat{\Sigma}^{-1}$ and $\hat{\Sigma} = \mathbb{P}_n\{(X - \hat{\mathbb{E}}[X | \hat{U}])^{\otimes 2}\}$.

Output: Confidence intervals and p-values based on asymptotic null distribution $\tilde{b}_{j,i} \sim \mathcal{N}_d(\tilde{\beta}_{j,i}, \frac{\hat{S}_j}{n})$.




Figure 6: Simulation results with 500 negative control outcomes out of a total of 1000 outcomes. For PII, the nuisance functions are estimated using random forests. The data model is the Logistic regression model. The first and second rows have noise levels $\sigma_\epsilon = 0.8$ and $\sigma_\epsilon = 1$, respectively, for the latent variables.

4 Simulation

We generate the data from generalized partial linear models. The covariate $X \in \mathbb{R}$ is sampled from $\mathcal{N}(0, 1)$; the latent variable $U = X\alpha + \epsilon \in \mathbb{R}^r$ is a linear function of X , where $r = 10$, $\alpha_{1j} \sim \text{Unif}(-1, 1)$ and $\epsilon_j \sim \mathcal{N}(0, \sigma_\epsilon^2)$ independently for $j \in [r]$; and the response is generated from generalized linear models with a Logistic link $\text{logit}(\mathbb{E}[Y | X, U]) = X\beta + U\eta$, where $\beta_{1j} \sim 2 \times \text{Bernoulli}(0.2)$ and $\sqrt{r} \cdot \eta_{ij} \sim \text{Unif}(-1, 1)$ independently for $i \in [r]$ and $j \in [p]$. We set the total number of outcomes to be $p = 1000$, and use 500 null outcomes as the negative outcomes.

We evaluate four methods: (1) GLM (X): naive generalized linear models that use Logistic regression that only uses observed covariate X to predict Y ; (2) GLM (X, U): oracle Logistic regression that uses both observed covariate X and latent variable U to predict Y ; (3) PII (X, U): the proposed post-integrated inference method that uses observed covariate X and latent embedding U to predict Y ; and (4) PII (X, \hat{U}): the proposal method that uses the first r PCs of the outcome matrix are selected as \hat{U} .

For PII, we use the random forest to estimate the nuisance functions $\mathbb{E}[X | U]$, $\mathbb{E}[Y | X, U]$, and $\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U]$ and apply extrapolated cross-validation (ECV) (Du et al., 2024a) to select the hyperparameter that minimizes the estimated mean squared error. ECV allows us to use a smaller number of trees for estimating the out-of-sample prediction errors based on out-of-bag observations and extrapolate the risk estimation up to a larger number of trees consistently without sample splitting. In our experiment, we use 25 trees to perform ECV and the hyperparameters we consider include: ‘max_depth’ in {1, 3, 5} for the depth of each tree, ‘max_samples’ in {0.25, 0.5, 0.75, 1} for bootstrap samples and the number of trees in {1, ..., 50}.

To compare the performance of different methods, the power, type-I error, and false discovery

proportion (FDP) for hypothesis testing are analyzed. For both the type-I error and power, we set the significance level to be 0.05. For FDP, we use the Benjamini-Hochberg procedure with FDR controlled at 0.05. As shown in the first two columns of Figure 6, the GLM-NAIVE regression method fails to control the inflated type-I error, resulting in numerous false positives. Furthermore, as the sample size increases, this method becomes even more anti-conservative. Conversely, the GLM-ORACLE regression method exhibits tight control over type-I error, as expected. When the latent embedding U is known, we observe that PII also effectively controls type-I error. Additionally, under certain conditions, PII provides greater power than the GLM-ORACLE. This may be attributed to PII’s ability to address the effect of collinearity between X and U on the nonlinear outcome models through a two-step procedure, whereas GLM-ORACLE does not, leading to conservative results.

When the latent embedding U is unknown, we evaluate the performance of the estimated \hat{U} . As shown in the third panel of Figure 6, the error of embedding projection matrix $\|P_{\hat{U}} - P_U\|_{\text{op}}$ decreases rapidly as the sample size n increases. When U can be well approximated, PII experiences a slightly inflated type-I error because it targets the modified main effect $\tilde{\beta}$ instead of the true effect β . However, the statistical error remains reasonable, the FDP is controlled at the desired level, and PII achieves greater power compared to the oracle GLM in many cases. Lastly, PII exhibits greater power when the conditional variation of X given U is large (i.e., $\mathbb{V}(\epsilon)$ is relatively larger than the linear projected signal strength $\|\gamma\|$). One could potentially use the ratio of these two quantities as a metric to quantify the level of confounding.

5 Application on single-cell CRISPR data analysis

Background In a recent study by Lalli et al. (2020), the molecular mechanisms of genes associated with neurodevelopmental disorders, particularly Autism Spectrum Disorder (ASD), were investigated using a modified CRISPR-Cas9 system. Experiments focused on 13 ASD-linked gene knockdowns in Lund Human Mesencephalic neural progenitor cells, with gene expression changes assessed through single-cell RNA sequencing. The progression of neuronal differentiation was estimated via a pseudotime trajectory (Figure F2), revealing that some genetic perturbations impact this progression.

Confounders significantly affect the interpretation of single-cell CRISPR perturbation results, as these experiments often resemble observational studies. Variables like cell size, cycle stage, and microenvironment heterogeneity can alter gene expression patterns, obscuring true genetic effects. To address these challenges, we use 4000 lowly variable genes as negative control outcomes for adjustment, focusing on 4163 highly variable genes for differential expression analysis on 8320 cells. The data preprocessing procedure is detailed in Appendix F.2.

Compared methods and embedding estimation. We compare the proposed method with four methods for hypothesis testing: (1) GLM: Score tests based on generalized linear models with Negative Binomial likelihood and log link function. The covariance matrix is estimated using the HC3-type robust estimator. This method does not adjust for potential confounding effects. (2) RUV4: A heuristic method proposed by Gagnon-Bartsch and Speed (2012) that uses principle components on the residual matrix of regressing the negative control outcomes on the covariate of interest to estimate the latent embeddings. Based on heuristic calculations, the authors claim that the RUV-4 estimator has approximately the oracle variance. (3) CATE-NC: The deconfounding method CATE proposed by Wang et al. (2017) with negative controls, which uses maximum likelihood estimation to estimate the latent embedding. Under simplified Gaussian linear models, they show that their estimator has asymptotical type I error control when the number of negative controls is large. (4) CATE-RR: A variant of CATE method (Wang et al., 2017) with robust regression, which is similar to the heuristic algorithm LEAPP (Sun et al., 2012) and utilizes the sparsity of effects to estimate the latent embeddings.




Figure 7: Histogram of t -statistics of *PTEN* perturbation by different methods. PCA with 50 components, RUV4, CATE-NC, and CATE-RR.

For PII, we use four methods to estimate the cell embeddings, including PCA, RUV4, CATE-NC, and CATE-RR. The first three methods use negative control to estimate the embedding, while the last is only valid under the sparsity assumption on the effects. Before running PCA, we follow the preprocessing procedure in single-cell data analysis to adjust the library size of each cell to be 10^4 , add one pseudo count, and take the logarithm. We then select the top 50 principal components as the estimated embeddings. For the last three embedding estimation methods, we supply all 13086 genes as input, specify the set of pseudo-negative control genes when applicable, and set the number of factors to 10, a value commonly used by researchers based on empirical evidence. Though not presented in the paper, we observed similar results with higher numbers of factors.

Results. The study by Lalli et al. (2020) indicates that some perturbations affect changes in gene expressions along pseudotime, potentially altering development speed. Using pseudotime as a covariate allows us to examine if perturbations explain effects beyond developmental changes; see Appendix F.2 for the extended background of the data. Biologically, we expect more signals on pseudotime states (Figure F3) than on perturbation conditions.

We focus on the target gene *PTEN*, which is crucial in neural development and differentiation and influences other genes in a cascading manner when it is knockdown (Lalli et al., 2020). Examining the empirical distribution of test statistics for perturbation conditions using GLM reveals conservative results for genes like *CTNND2*, *MECP2*, and *MYT1L* (Figure F4). This suggests that GLM without adjusting for hidden confounders leads to biased hypothesis testing. PII corrects these biases (Figure 2, Figure 7), and even PCA-based simple embedding estimation effectively calibrates test distributions. Comparing methods RUV4, CATE-NC, CATE-RR with PII, we see PII reduces distinct discoveries from 45.6% to 38.8% (Figure 8(a)), indicating more coherent outcomes with PII.

To further assess the biological significance of the discoveries, we examine 276 and 203 common discoveries (Figure 8(a) and Figure 8(b)). Significant discrepancies are noted (Figure 8(c)). The associated gene ontology terms on biological processes using `clusterProfiler` package with default




Figure 8: Summary of significant genes ($p\text{-values} < 0.05$) associated with *PTEN* perturbation by different confounder adjustment methods. (a) Upset plot of discoveries by three methods: RUV4, CATE-NC, and CATE-RR, as in Figure F4. (b) Upset plot of discoveries by PII with embedding estimated by three methods: RUV4, CATE-NC, and CATE-RR, as in Figure 7. (c) The Venn plot of two sets of discoveries. One set includes 276 common discoveries by RUV4, CATE-NC, and CATE-RR, while the other includes 203 common discoveries by PII with the same estimated embeddings given by the three methods. (d) Gene ontology analysis of 137 distinct discoveries by PII.

false discovery control threshold (Yu et al., 2012) reveal that unlike the 210 genes from RUV4, CATE-NC, and CATE-RR, which had no associated GO terms, the 137 genes unique to PII align with ribosome-related processes (Figure 8(d)), supporting studies on *PTEN*'s impact on these processes (Liang et al., 2017; Cheung et al., 2023).

6 Discussion

A potential concern of the proposed method is whether the estimated embeddings might act as colliders, especially if \widehat{U} is influenced by both X and Y_{C^c} . However, our fundamental assumption is that Y_C is driven by a low-dimensional embedding U but not the covariate X , which inherently mitigates the risk of \widehat{U} becoming a collider. If this foundational assumption does not hold, the direct effect estimand (5) might not align with researchers' interests, necessitating the use of domain knowledge to identify and investigate alternative target estimands.

Our paper focuses on design-free deconfounding with negative control outcomes, though other strategies exist (detailed comparison in Appendix B). While our framework allows for flexible machine learning algorithms, it introduces computational complexity, especially with increasing outcomes and hyperparameter tuning. For practical applications, specialized models like variational autoencoders for joint outcome function fitting (Du et al., 2022; Moon et al., 2024) and efficient cross-validation methods can be beneficial.

Further extensions involve incorporating interaction effects (Vansteelandt and Dukes, 2022), developing tests for nonparametric confounding (Miao et al., 2018). Explorations into settings with

high-dimensional latent embeddings and covariates (Miao et al., 2023; Zeng et al., 2024) could also be of interest. In real data analysis, we use pseudo-negative control outcomes, which can be viewed as one variant of the synthetic control approaches (Abadie et al., 2010). Providing theoretical guarantees for the valid construction of negative control outcomes from data remains an area of practical interest.

Supplement

The appendix includes related work, comparisons with related deconfounding strategies, proof of all the theoretical results, and extra experiment results. The outline of the appendix and the summary of the notation are given below.

References

- Abadie, A., Diamond, A., and Hainmueller, J. (2010). Synthetic control methods for comparative case studies: Estimating the effect of California's tobacco control program. *Journal of the American Statistical Association*, 105(490):493–505.
- Adams, J. and Hansen, N. R. (2024). Substitute adjustment via recovery of latent variables. *arXiv preprint arXiv:2403.00202*.
- Andrews, I., Stock, J. H., and Sun, L. (2019). Weak instruments in instrumental variables regression: Theory and practice. *Annual Review of Economics*, 11(1):727–753.
- Antonsson, S. E. and Melsted, P. (2024). Batch correction methods used in single cell RNA-seq analyses are often poorly calibrated. *bioRxiv*, pages 2024–03.
- Bai, J. and Li, K. (2012). Statistical analysis of factor models of high dimension. *The Annals of Statistics*, pages 436–465.
- Bai, J. and Ng, S. (2002). Determining the number of factors in approximate factor models. *Econometrica*, 70(1):191–221.
- Berk, R., Buja, A., Brown, L., George, E., Kuchibhotla, A. K., Su, W., and Zhao, L. (2021). Assumption lean regression. *The American Statistician*.
- Biau, G. (2012). Analysis of a random forests model. *The Journal of Machine Learning Research*, 13(1):1063–1095.
- Billingsley, P. (1995). *Probability and measure*. Wiley Series in Probability and Statistics. Wiley.
- Bing, X., Cheng, W., Feng, H., and Ning, Y. (2023). Inference in high-dimensional multivariate response regression with hidden variables. *Journal of the American Statistical Association*, pages 1–12.
- Bing, X., Ning, Y., and Xu, Y. (2022). Adaptive estimation in multivariate response regression with hidden variables. *The Annals of Statistics*, 50(2):640–672.
- Chen, J., Li, P., and Fu, Y. (2012). Inference on the order of a normal mixture. *Journal of the American Statistical Association*, 107(499):1096–1105.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters.
- Cheung, S. K., Kwok, J., Or, P. M., Wong, C. W., Feng, B., Choy, K. W., Chang, R. C., Burbach, J. P. H., Cheng, A. S., and Chan, A. M. (2023). Neuropathological signatures revealed by transcriptomic and proteomic analysis in pten-deficient mouse models. *Scientific Reports*, 13(1):6763.
- Du, J.-H., Cai, Z., and Roeder, K. (2022). Robust probabilistic modeling for single-cell multimodal mosaic integration and imputation via scvaeit. *Proceedings of the National Academy of Sciences*, 119(49):e2214414119.
- Du, J.-H., Patil, P., Roeder, K., and Kuchibhotla, A. K. (2024a). Extrapolated cross-validation for randomized ensembles. *Journal of Computational and Graphical Statistics*, pages 1–12.
- Du, J.-H., Wasserman, L., and Roeder, K. (2023). Simultaneous inference for generalized linear models with unmeasured confounders. *arXiv preprint arXiv:2309.07261*.

- Du, J.-H., Zeng, Z., Kennedy, E. H., Wasserman, L., and Roeder, K. (2024b). Causal inference for genomic data with multiple heterogeneous outcomes. *arXiv preprint arXiv:2404.09119*.
- Gagnon-Bartsch, J. A. and Speed, T. P. (2012). Using control genes to correct for unwanted variation in microarray data. *Biostatistics*, 13(3):539–552.
- Galbraith, J. W. and Zinde-Walsh, V. (2020). Simple and reliable estimators of coefficients of interest in a model with high-dimensional confounding effects. *Journal of econometrics*, 218(2):609–632.
- Hao, Y., Hao, S., Andersen-Nissen, E., III, W. M. M., Zheng, S., Butler, A., Lee, M. J., Wilk, A. J., Darby, C., Zagar, M., Hoffman, P., Stoeckius, M., Papalexis, E., Mimitou, E. P., Jain, J., Srivastava, A., Stuart, T., Fleming, L. B., Yeung, B., Rogers, A. J., McElrath, J. M., Blish, C. A., Gottardo, R., Smibert, P., and Satija, R. (2021). Integrated analysis of multimodal single-cell data. *Cell*.
- Härdle, W., Liang, H., and Gao, J. (2000). *Partially linear models*. Springer Science & Business Media.
- Härdle, W., Mammen, E., and Müller, M. (1998). Testing parametric versus semiparametric modeling in generalized linear models. *Journal of the American Statistical Association*, 93(444):1461–1474.
- Howe, G. W. and Brown, C. H. (2023). Retrospective psychometrics and effect heterogeneity in integrated data analysis: Commentary on the special issue. *Prevention Science*, 24(8):1672–1681.
- Imbens, G. W. and Rubin, D. B. (2015). *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press.
- Jaglin, X. H., Poirier, K., Saillour, Y., Buhler, E., Tian, G., Bahi-Buisson, N., Fallet-Bianco, C., Phan-Dinh-Tuy, F., Kong, X. P., Bomont, P., et al. (2009). Mutations in the β -tubulin gene *tubb2b* result in asymmetrical polymicrogyria. *Nature genetics*, 41(6):746–752.
- Jia, Q., Nie, H., Yu, P., Xie, B., Wang, C., Yang, F., Wei, G., and Ni, T. (2019). Hnrnpal-mediated 3' utr length changes of hn1 contributes to cancer-and senescence-associated phenotypes. *Aging (Albany NY)*, 11(13):4407.
- Johnson, W. E., Li, C., and Rabinovic, A. (2007). Adjusting batch effects in microarray expression data using empirical bayes methods. *Biostatistics*, 8(1):118–127.
- Kennedy, E. H. (2022). Semiparametric doubly robust targeted double machine learning: a review. *arXiv preprint arXiv:2203.06469*.
- Kim, K., Li, B., Yu, Z., and Li, L. (2020). On post dimension reduction statistical inference. *Annals of Statistics*, 48(3):1567–1592.
- Lalli, M. A., Avey, D., Dougherty, J. D., Milbrandt, J., and Mitra, R. D. (2020). High-throughput single-cell functional elucidation of neurodevelopmental disease-associated genes reveals convergent mechanisms altering neuronal differentiation. *Genome research*, 30(9):1317–1331.
- Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E., and Storey, J. D. (2012). The sva package for removing batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics*, 28(6):882–883.
- Lehmann, E. L. and Scheffé, H. (2012). *Completeness, similar regions, and unbiased estimation—part II*. Springer.

- Li, T., Zhang, Y., Patil, P., and Johnson, W. E. (2023). Overcoming the impacts of two-step batch effect correction on gene expression estimation and inference. *Biostatistics*, 24(3):635–652.
- Liang, H., Chen, X., Yin, Q., Ruan, D., Zhao, X., Zhang, C., McNutt, M. A., and Yin, Y. (2017). $\text{Pten}\beta$ is an alternatively translated isoform of pten that regulates rdna transcription. *Nature communications*, 8(1):1–14.
- Luecken, M. D., Büttner, M., Chaichoompu, K., Danese, A., Interlandi, M., Müller, M. F., Strobl, D. C., Zappia, L., Dugas, M., Colomé-Tatché, M., et al. (2022). Benchmarking atlas-level data integration in single-cell genomics. *Nature methods*, 19(1):41–50.
- Luo, X. and Wei, Y. (2018). Batch effects correction with unknown subtypes. *Journal of the American Statistical Association*.
- Ma, R., Sun, E. D., Donoho, D., and Zou, J. (2024). Principled and interpretable alignability testing and integration of single-cell data. *Proceedings of the National Academy of Sciences*, 121(10):e2313719121.
- Mammen, E., Rothe, C., and Schienle, M. (2012). Nonparametric regression with nonparametrically generated covariates. *The Annals of Statistics*, pages 1132–1170.
- Miao, W., Geng, Z., and Tchetgen Tchetgen, E. J. (2018). Identifying causal effects with proxy variables of an unmeasured confounder. *Biometrika*, 105(4):987–993.
- Miao, W., Hu, W., Ogburn, E. L., and Zhou, X.-H. (2023). Identifying effects of multiple treatments in the presence of unmeasured confounding. *Journal of the American Statistical Association*, 118(543):1953–1967.
- Miao, W., Shi, X., Li, Y., and Tchetgen Tchetgen, E. J. (2024). A confounding bridge approach for double negative control inference on causal effects. *Statistical Theory and Related Fields*, pages 1–12.
- Moon, H., Du, J.-H., Lei, J., and Roeder, K. (2024). Augmented doubly robust post-imputation inference for proteomic data. *bioRxiv*, pages 2024–03.
- Newey, W. K. and Robins, J. R. (2018). Cross-fitting and fast remainder rates for semiparametric estimation. *arXiv preprint arXiv:1801.09138*.
- Paddock, S. M., Franco, C., Breidt, F. J., and Betancourt, B. (2024). Statistical data integration for health policy evidence-building. *Annual Review of Statistics and Its Application*, 12.
- Patil, P., Du, J.-H., and Kuchibhotla, A. K. (2023). Bagging in overparameterized learning: Risk characterization and risk monotonization. *Journal of Machine Learning Research*, 24(319):1–113.
- Robins, J., Li, L., Tchetgen, E., van der Vaart, A., et al. (2008). Higher order influence functions and minimax estimation of nonlinear functionals. In *Probability and statistics: essays in honor of David A. Freedman*, volume 2, pages 335–422. Institute of Mathematical Statistics.
- Robins, J. M., Mark, S. D., and Newey, W. K. (1992). Estimating exposure effects by modelling the expectation of exposure conditional on confounders. *Biometrics*, pages 479–495.
- Robinson, P. M. (1988). Root-n-consistent semiparametric regression. *Econometrica: Journal of the Econometric Society*, pages 931–954.
- Rosenbaum, P. R. (1989). The role of known effects in observational studies. *Biometrics*, pages 557–569.

- Scornet, E., Biau, G., and Vert, J.-P. (2015). Consistency of random forests. *The Annals of Statistics*, pages 1716–1741.
- Severini, T. A. and Staniswalis, J. G. (1994). Quasi-likelihood estimation in semiparametric models. *Journal of the American statistical Association*, 89(426):501–511.
- Shi, X., Miao, W., Nelson, J. C., and Tchetgen Tchetgen, E. J. (2020). Multiply robust causal inference with double-negative control adjustment for categorical unmeasured confounding. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82(2):521–540.
- Shi, X., Pan, Z., and Miao, W. (2023). Data integration in causal inference. *Wiley Interdisciplinary Reviews: Computational Statistics*, 15(1):e1581.
- Sun, Y., Zhang, N. R., and Owen, A. B. (2012). Multiple hypothesis testing adjusted for latent variables, with an application to the agemap gene expression data. *The Annals of Applied Statistics*, pages 1664–1688.
- Tran, H. T. N., Ang, K. S., Chevrier, M., Zhang, X., Lee, N. Y. S., Goh, M., and Chen, J. (2020). A benchmark of batch-effect correction methods for single-cell rna sequencing data. *Genome biology*, 21:1–32.
- Trefethen, L. N. and Bau, D. (2022). *Numerical linear algebra*. SIAM.
- Vansteelandt, S. and Dukes, O. (2022). Assumption-lean inference for generalised linear model parameters. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(3):657–685.
- Wang, J., Zhao, Q., Hastie, T., and Owen, A. B. (2017). Confounder adjustment in multiple hypothesis testing. *The Annals of Statistics*, 45(5):1863.
- Windmeijer, F., Liang, X., Hartwig, F. P., and Bowden, J. (2021). The confidence interval method for selecting valid instrumental variables. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 83(4):752–776.
- Yang, S., Kim, J. K., and Song, R. (2020). Doubly robust inference when combining probability and non-probability samples with high dimensional data. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82(2):445–465.
- Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterprofiler: an r package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology*, 16(5):284–287.
- Zeng, Z., Balakrishnan, S., Han, Y., and Kennedy, E. H. (2024). Causal inference with high-dimensional discrete covariates. *arXiv preprint arXiv:2405.00118*.
- Zhang, Z., Mathew, D., Lim, T., Mason, K., Martinez, C. M., Huang, S., Wherry, E. J., Susztak, K., Minn, A. J., Ma, Z., et al. (2023). Signal recovery in single cell batch integration. *bioRxiv*.
- Zhou, Y., Tang, D., Kong, D., and Wang, L. (2024). Promises of parallel outcomes. *Biometrika*, 111(2):537–550.

Supplementary Material

The appendix includes related work, comparisons with related deconfounding strategies, proof of all the theoretical results, and extra experiment results. The outline of the appendix and the summary of the notation are given below.

Outline. The structure of the appendix is listed below:

Appendix	Content
Appendix A	Related work.
Appendix B	B.1 Comparisons with design-based deconfounding strategies. B.2 Comparisons with sparsity-based deconfounding strategies.
Appendix C	Proof of Theorem 1.
Appendix D	D.1 Proof of Theorem 2. D.2 Proof of Lemma 3. D.3 Auxiliary lemmas: Lemma D.1, D.2, and D.3.
Appendix E	E.1 Proof of Theorem 4 and Corollary 5. E.2 Proof of Proposition 6. E.3 DR inference with nonlinear link function: Theorem E.1. E.4 Auxiliary lemmas: Lemma E.2, E.3, E.4 and E.5.
Appendix F	F.1 Extra experimental results on simulation. F.2 Extra experimental results on real data.

Notation. Throughout our exposition, we will use the following notational conventions. We use uppercase letters for random variables/vectors (e.g., Y, X, U) and lowercase for sample vectors, respectively (e.g., y, x, u). For a matrix $\beta \in \mathbb{R}^{d \times p}$, its j th column is denoted by $\beta_{\cdot j}$. Sets are denoted by calligraphic uppercase letters (\mathcal{A}, \mathcal{C}). Bold font is only used to denote design matrices and response matrices (e.g., $\mathbf{Y}, \mathbf{X}, \mathbf{U}$) whose first dimension equals the sample size. For $p \in \mathbb{N}$, $[p] := \{1, \dots, p\}$. For a set \mathcal{A} , let $|\mathcal{A}|$ be its cardinality.

For a random vector $X \in \mathbb{R}^p$, \mathcal{P}_X denotes the projection in L_2 . For any matrix $A \in \mathbb{R}^{n \times p}$ with full column rank, let $P_A = A(A^\top A)^{-1}A^\top$ and $P_A^\perp = I_p - P_A$ be the orthogonal projection matrices on the A 's column space and its orthogonal space, respectively. For any square matrix $A \in \mathbb{R}^{n \times n}$, $\lambda_i(A)$ denotes its i th eigenvalue. The Gram matrix of A^\top is denoted by $A^{\otimes 2} := AA^\top$. Matrix Hadamard product is denoted by \odot . For two symmetric matrices $A, B \in \mathbb{R}^{n \times n}$, we write that $A \preceq B$ ($A \succeq B$) if $B - A$ ($A - B$) is positive semi-definite. For $a \in \mathbb{R}^m$, $\|a\|_q$ denotes the ℓ_q -norm for $q = 1, \dots, \infty$. For $a \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$, $\|a\|$ and $\|A\|$ denote the ℓ_2 -norm and operator norm, respectively. The condition number of A is defined as $\kappa(A) = \|A\| \|A^{-1}\|$. For any random vector X , its L_q norm is defined as $\|X\|_{L_q} = \mathbb{E}[\|X\|_q^{q/1/q}]$ for $q = 1, \dots, \infty$.

For (potentially random) measurable functions f , we denote expectations with respect to Z alone by $\mathbb{P}f(Z) = \int f d\mathbb{P}$, and with respect to both Z and the observations where f is fitted on by $\mathbb{E}[f(Z)]$. The empirical expectation is denoted by $\mathbb{P}_nf(Z) = \frac{1}{n} \sum_{i=1}^n f(Z_i)$. Similarly, the population and empirical variances (or covariance) are denoted by \mathbb{V} and \mathbb{V}_n , respectively. The identity map is denoted by \mathbb{I} . We write the (conditional) L_p norm of f as $\|f\|_{L_p} = [\int f(z)^p d\mathbb{P}(z)]^{1/p}$ for $p \geq 1$.

We use “ o ” and “ \mathcal{O} ” to denote the little-o and big-O notations and let “ $o_{\mathbb{P}}$ ” and “ $\mathcal{O}_{\mathbb{P}}$ ” be their probabilistic counterparts. For sequences $\{a_n\}$ and $\{b_n\}$, we write $a_n \ll b_n$ or $b_n \gg a_n$ if $a_n = o(b_n)$; $a_n \lesssim b_n$ or $b_n \gtrsim a_n$ if $a_n = \mathcal{O}(b_n)$; and $a_n \asymp b_n$ if $a_n = \mathcal{O}(b_n)$ and $b_n = \mathcal{O}(a_n)$. Convergence in distribution and probability are denoted by “ \xrightarrow{d} ” and “ \xrightarrow{P} ”. For $a, b \in \mathbb{R}$, we write $a \vee b = \max\{a, b\}$ and $a \wedge b = \min\{a, b\}$.

A Related work

Batch correction and data integration. Large-scale single-cell transcriptomic datasets often include samples that span locations, laboratories, and conditions, leading to complex, nested batch effects in data (Tran et al., 2020; Luecken et al., 2022). Batch correction specifically targets the removal of unwanted variation due to differences in batches within a single study, ensuring that the remaining data is comparable and reflects true biological differences. On the other hand, data integration focuses on combining and harmonizing multiple datasets to enhance statistical power and provide a more comprehensive analysis, dealing with both batch effects and between-dataset heterogeneity. Despite these differences, batch correction and data integration share the common goal of removing unwanted variation and preserving biological variation (Zhang et al., 2023). The integrated data is then used for downstream analysis, such as dimension reduction, clustering, and differential expressed gene testing. Integrated cellular profiles are typically represented as an integrated graph, a joint embedding, or a corrected response matrix. The main focus of the current paper is on the last category.

Despite the efforts from the computational biology and machine learning community to achieve better predictive power and data alignment, most existing batch correction methods are shown to be poorly calibrated (Antonsson and Melsted, 2024; Ma et al., 2024). For statistical inference, many heuristic methods have been proposed to remove the batch effects and unwanted variations in the past decade. Leading examples include Remove Unwanted Variation (RUV) (Gagnon-Bartsch and Speed, 2012) and Surrogate Variable Analysis (SVA) (Leek et al., 2012). RUV/SVA uses estimated factors of unwanted variation from unadjusted data, which works even if the batch design is unknown. When the batch design is known, two-step procedures for batch correction have also been proposed under parametric or mixture models (Li et al., 2023; Luo and Wei, 2018).

Unmeasured confounders adjustment and negative control outcomes. Over the past decades, researchers have been exploring methods to address the issue of unmeasured confounders in statistical analysis. In the presence of multiple outcomes, deconfounding techniques primarily employ two strategies: incorporating known negative control outcomes or leveraging sparsity assumptions (Wang et al., 2017); while there is also another line of research on proximal causal inference, which uses both negative control outcomes and/or exposures for deconfounding (Miao et al., 2018). For a comprehensive review of the literature on sparsity-based methods, readers are directed to Du et al. (2023) and Zhou et al. (2024). This paper focuses on the negative control approach in the context of multiple outcomes.

Most existing works on confounder adjustment presume the knowledge of causal structure when the unobserved variable U is a mediator (Wang et al., 2017) and when U is a confounder Miao et al. (2023), corresponding to Figure 3(a) and Figure 3(b), respectively. Recently developed sparsity-based methods by Bing et al. (2023); Du et al. (2023) have tried to relax this assumption to allow for a more flexible relationship between X and U . In particular, each entry of U can belong to different cases in Figure 3.

Negative control outcomes are used in observational studies under the key assumption that exposure has no causal effect on these outcomes. Rosenbaum (1989) demonstrated that negative control outcomes can be employed to test for the presence of hidden confounding in observational studies. By introducing an additional variable known as a negative control exposure, Miao et al. (2018) further showed that the average causal effect can be identified nonparametrically. Building upon this work, Shi et al. (2020) developed a semiparametric inference procedure specifically for scenarios involving a categorical latent confounder and a binary exposure. Under linear latent models, Galbraith and Zinde-Walsh (2020) use principal components of a set of potential controls to adjust for unmeasured confounding effects. Under nonparametric models for a single outcome and multiple treatments Miao et al. (2023) derive nonparametric identification conditions.

Assumption-lean semiparametric inference. There is increasing interest in deriving assumption-lean inference by using projection-based estimators (Berk et al., 2021) or semiparametric estimators (Vansteelandt and Dukes, 2022). The inferential problems we considered are also related to two-stage inference problems, such as post-sufficient dimension reduction inference (Kim et al., 2020), post-imputation inference (Moon et al., 2024), and inference with substituted covariate (Adams and Hansen, 2024) or nonparametrically generated covariates (Mammen et al., 2012). While these related methods offer valuable insights into two-stage inference processes, they do not directly extend to address the challenges encountered in post-integrated inference problems.

B Comparisons with related deconfounding approaches

B.1 Design-based approaches

As mentioned in the introduction, our paper mainly focuses on design-free data integration approaches. However, it is possible to relate the design-based approaches to design-free approaches so that the proposed method can be applied, as we discuss below. Design-based data integration approaches, such as Combat (Johnson et al., 2007) and BUS (Luo and Wei, 2018), are usually based on a linear model:

$$Y_j = \alpha_j + X\beta_j + \gamma_{Bj} + \epsilon_{Bj}, \quad j = 1, \dots, p,$$

where $\alpha_j, \beta_j \in \mathbb{R}$ are coefficients for common variations while $\gamma_{Bj} \in \mathbb{R}$ is the location and $\epsilon_{Bj} \in \mathbb{R}$ is a mean-zero noise with scale differences across batches, respectively, for batch $B \in [n_B - 1]$ (with group $B = 0$ being the baseline and $\gamma_{0j} = 0$) and n_B is the total number of batches. This implies that

$$\mathbb{E}[Y_j | X, B] = \alpha_j + X\beta_j + \gamma_{Bj}, \quad j = 1, \dots, p.$$

Let $U_B \in \{0, 1\}^{n_B}$ be the one-hot vector with only the B -th entry being one and zero elsewhere. Then, we can rewrite the above as

$$\mathbb{E}[Y | X, U_B] = \alpha + \beta^\top X + f(U_B)$$

where $f(U_B) = \gamma^\top U_B$ and $\gamma = [\gamma_{bj}]_{b \in [n_B], j \in [p]}$. In other words, the location-and-scale model considered by Johnson et al. (2007) and Luo and Wei (2018) is a special case of partial linear models with heterogenous noises, though they have utilized empirical Bayes shrinkage to improve the estimates. For this reason, a generalized least square approach could be used to improve Combat, as suggested by Li et al. (2023).

In fact, when the additive noises are normal, we can decompose the noise as $\epsilon_{Bj} = U_\epsilon + Z_j$ for $B > 0$ such that $U_\epsilon \perp\!\!\!\perp \epsilon'_j$ and $\epsilon'_j \stackrel{d}{=} \epsilon_{0j}$. To see this, define $\tau^2 = \mathbb{V}(\epsilon_{Bj})$ and $\sigma^2 = \mathbb{V}(\epsilon_{0j})$. Without loss of generality, we assume $\epsilon_{0j} \leq \min_{b \in [n_B]} \epsilon_{bj}$ so that $\tau^2 \geq \sigma^2$. If we define $U_\epsilon := \frac{1}{\tau(\tau^2 - \sigma^2)} \epsilon_{Bj} + Z_j$ and $\epsilon'_j := \epsilon_{Bj} - U_\epsilon$, where $Z_j \sim \mathcal{N}(0, \frac{\sigma^2}{(\tau^2 - \sigma^2)^2})$ is independent of ϵ_{Bj} , then U_ϵ and $\epsilon_{Bj} - U_\epsilon$ are independent because $\text{Cov}(U_\epsilon, \epsilon_{Bj} - U_\epsilon) = (\tau^2(\tau^2 - \sigma^2)^2)^{-1} \mathbb{V}(\epsilon_{Bj}) - \mathbb{V}(Z_j) = 0$. Here, we use the fact that two jointly normal random variables are independent if they are uncorrelated. In other words, we can rewrite the above model as

$$\mathbb{E}[Y | X, U] = \alpha + \beta^\top X + f(U),$$

where $f(U) = [\gamma, \mathbf{1}\{B > 0\}]^\top U$ and $U = [U_B, U_\epsilon]$. By absorbing part of the randomness of the additive noises into U , we convert the problem with heterogeneous noises into one with homogeneous noises studied in the current paper.

B.2 Unknown negative control outcomes

In this paper, we have focused on negative control outcomes to remove unwanted variations. When the negative control outcomes are unknown in advance, there are still possibilities to estimate the latent embedding and provide valid inferences. However, this typically requires extra sparsity assumptions on the effects of the covariate on multiple outcomes rather than utilizing the negative control outcomes. To illustrate the idea, we consider the following partial linear model:

$$\mathbb{E}[Y | X, U] = \beta^\top X + h(U).$$

Many methods start from the projected model

$$\mathbb{E}[\mathcal{P}_X^\perp Y | X, U] = \mathcal{P}_X^\perp h(U). \quad (10)$$

If the function $\mathcal{P}_X^\perp h$ has a good structure, then one may be able to recover U from $\mathcal{P}_X^\perp Y$. Alternatively, we can linearize the problem and seek partial recovery of the effect, as demonstrated below.

Example 1 (Linear models). If h is a linear function such that $h : U \mapsto \eta^\top U$ for $\eta \in \mathbb{R}^{r \times p}$, then

$$Y = [\beta \ \eta]^\top \begin{bmatrix} X \\ U \end{bmatrix} + E.$$

With n i.i.d. samples, we obtain the following equation in matrix form:

$$\mathbf{Y} = \mathbf{X}\beta + \mathbf{U}\eta + \mathbf{E}. \quad (11)$$

Early methods in the literature rely on the assumption of the functional relationship between X and U . For example, Wang et al. (2017) assume U to be a linear function of X with an additive Gaussian noise:

$$U = X\alpha + W, \quad (12)$$

in which case the sample counterpart of (10) reduces to

$$\mathcal{P}_{\mathbf{X}}^\perp \mathbf{Y} = \mathcal{P}_{\mathbf{X}}^\perp (\mathbf{W}\eta + \mathbf{E}). \quad (13)$$

Because the orthogonal projection is rank-deficient, one can further eliminate d rows of the above system of equations by elementary matrix transformation. For this purpose, Wang et al. (2017) use QR decomposition by Householder rotation to derive a linear system of $n - d$ equations; e.g., Equation (2.5) and Equation (4.5) in Wang et al. (2017) for $d = 1$ and $d > 1$, respectively. From this, $\hat{\eta}$ is recovered from quasi-log-likelihood estimation. In the second step, the unknown coefficient (α, β) is estimated from (13) by plugging in the estimate $\hat{\eta}$.

Under more general confounding mechanism when (12) does not necessarily hold, Bing et al. (2022, 2023) rotate the original system to consistently estimate the marginal effect, under sparsity assumption on β and proper moment assumptions. They then use the residual from the lava fit to uncover the column space of η . Finally, the partial coefficient βP_η^\perp is recovered from the rotated system:

$$\mathbf{Y} P_\eta^\perp = \mathbf{X}\beta P_\eta^\perp + \mathbf{E} P_\eta^\perp.$$

These results have been extended to generalized linear models by Du et al. (2023) using joint maximum likelihood estimation. When β is sparse, then it can be recovered by some estimator of βP_η^\perp asymptotically. Note that the above approaches do not have too many restrictions on the observed covariate X and the latent embedding U , except for certain bounded moment assumptions.

Inspired by the success of methodologies development under linear models Example 1, one strategy for an extension to a nonlinear model is by linearizing the estimation problem. Specifically, suppose $h(U) = U\eta + R(U)$ for some remainder term R that depends on U , similarly we have a projection-based decomposition:

$$\begin{aligned} P_{\mathbf{X}}^{\perp}Y &= P_{\mathbf{X}}^{\perp}\mathbf{U}\eta + P_{\mathbf{X}}^{\perp}(R(\mathbf{U}) + \mathbf{E}) \\ \mathbf{Y}P_{\eta}^{\perp} &= \mathbf{X}\beta P_{\eta}^{\perp} + (R(\mathbf{U}) + \mathbf{E})P_{\eta}^{\perp}, \end{aligned}$$

from which one may seamlessly use the methods by Bing et al. (2022, 2023) and Du et al. (2023) when the remainder term can be well controlled.

Another possible strategy aligned with the angle of the current paper is to detect “weak” negative control outcomes and perform post-integrated inference based on such pseudo-negative control outcomes, as in Section 5. This approach is very similar to weak instrument detection and invalid instrumental variables selection; see, for example, Andrews et al. (2019) and Windmeijer et al. (2021). We expect the rich literature on these related problems could lead to new methodological advances in post-integrated inference problems.

C Nonparametric identification

Proof of Theorem 1. Under the equivalence assumption (Assumption 2 (ii)), for any admissible distribution $\tilde{f}(y_C, u)$ we must have some invertible function v such that $\tilde{f}(y_C, u) = f\{Y_C = y_C, v(U) = u\}$. Note that (2) has at least one solution $\tilde{f}(x | u) = f(x | v^{-1}(u))$; when this is the solution, define $\tilde{f}(y_{C^c} | x, u) := f(y_{C^c} | x, v^{-1}(u))$. Then, $\tilde{f}(y_{C^c} | x, u)$ is also one solution to (3).

Because $v(U)$ is invertible, the ignorability assumption (Assumption 1 (iii)) $Y(x) \perp\!\!\!\perp X | U$ implies that $Y(x) \perp\!\!\!\perp X | v(U)$; the completeness assumption Assumption 2 (iii) implies that $\tilde{f}(u) > 0$ on $u \in v(\mathcal{U})$ and $\tilde{f}(u | y_C, x; \alpha)$ is also complete in y_C . Further, from Assumption 1 (ii), the positivity condition $f_{X|v(U)}(x | v(u)) \in (0, 1)$ also holds for all $(x, u) \in \mathcal{X} \times \mathcal{U}$. Then, we have

$$\begin{aligned} f_{Y(x)}(y) &= \int f_{Y(x)|U}(y | u) f(u) du \\ &= \int f_{Y(x)|U,X}(y | u, x) f(u) du && (\text{Assumption 1 (ii)-(iii)}) \\ &= \int f(y | u, x) f(u) du && (\text{Assumption 1 (i)}) \\ &= \int f(y_{C^c} | u, x) f(y_C | u) f(u) du && (\text{Assumption 2 (i)}) \\ &= \int f(y_{C^c} | u, x) f(y_C, u) du \\ &= \int \tilde{f}(y_{C^c} | x, u) \tilde{f}(y_C, u) du, \end{aligned}$$

where the last equality follows from the same derivation of g-formula applied on random variables $(Y, X, v(U))$. This completes the proof for the second conclusion.

We next show the uniqueness of the solutions to (2) and (3). For any candidate solutions $\tilde{f}_1(x | u)$ and $\tilde{f}_2(x | u)$ to (2), we must have that

$$\int (\tilde{f}_1(x | u) - \tilde{f}_2(x | u)) \tilde{f}(u) du = 0,$$

which implies that $\tilde{f}_1(x | U) - \tilde{f}_2(x | U) = 0$ almost surely because of the completeness of $\tilde{f}(u)$. Note that for any candidate solutions $\tilde{f}_1(y_{C^c} | x, u)$ and $\tilde{f}_2(y_{C^c} | x, u)$ to (3), we must have that

$$\int (\tilde{f}_1(y_{C^c} | x, u) - \tilde{f}_2(y_{C^c} | x, u)) \tilde{f}(u | y_C, x) du \cdot f(y_C, x) = 0.$$

By the completeness property, this implies that $\tilde{f}_1(y_{C^c} | x, U) - \tilde{f}_2(y_{C^c} | x, U) = 0$ almost surely. Therefore, $\tilde{f}(y_{C^c} | x, u)$ is uniquely determined from (3). This completes the proof. \square

D Nonlinear main effects with estimated embeddings

D.1 Proof of Theorem 2

Proof of Theorem 2. Denote $A = \mathbb{E}[\text{Cov}(X | U)]$, $\hat{A} = \mathbb{E}[\text{Cov}(X | \hat{U})]$, $B = \mathbb{E}[\text{Cov}(X, \mathbb{E}[Y | X, U] | U)]$, and $\hat{B} = \mathbb{E}[\text{Cov}(X, \mathbb{E}[Y | X, U] | \hat{U})]$. From Lemma D.1, we know that the error of two linear regression coefficients $\|\beta_{\cdot j} - \hat{\beta}_{\cdot j}\|$ is governed by $\|A - \hat{A}\|$ and $\|B_{\cdot j} - \hat{B}_{\cdot j}\|$, where the subscript j indicates the j th column of the corresponding matrices.

Part (1) Covariance estimation errors. To apply Lemma D.1, we first derive the error bounds for the two quantities. Note that $\text{Cov}(X | U) = \mathbb{E}[X^{\otimes 2} | U] - \mathbb{E}[X | U]^{\otimes 2}$. We have

$$\begin{aligned} & \|A - \hat{A}\| \\ &= \|\mathbb{E}[\text{Cov}(X | U) - \text{Cov}(X | \hat{U})]\| \\ &= \|\mathbb{E}[\mathbb{E}[X | U]^{\otimes 2} - \mathbb{E}[X | \hat{U}]^{\otimes 2}]\| \\ &\leq \mathbb{E}[\|\mathbb{E}[X | U]^{\otimes 2} - \mathbb{E}[X | \hat{U}]^{\otimes 2}\|] && \text{(Jensen's inequality)} \\ &\leq \mathbb{E}[\|\mathbb{E}[X | U](\mathbb{E}[X | U] - \mathbb{E}[X | \hat{U}])^\top\| + \|(\mathbb{E}[X | U] - \mathbb{E}[X | \hat{U}])\mathbb{E}[X | \hat{U}]^\top\|] && \text{(triangle inequality)} \\ &= \mathbb{E}[\|\mathbb{E}[X | U]\|\|\mathbb{E}[X | U] - \mathbb{E}[X | \hat{U}]\| + \|\mathbb{E}[X | U] - \mathbb{E}[X | \hat{U}]\|\|\mathbb{E}[X | \hat{U}]\|] \\ &\leq (\|\mathbb{E}[X | U]\|_{L_2} + \|\mathbb{E}[X | \hat{U}]\|_{L_2})\|\mathbb{E}[X | \hat{U}] - \mathbb{E}[X | U]\|_{L_2} && \text{(Cauchy-Schwarz inequality)} \\ &\leq 2\|X\|_{L_2}\|\mathbb{E}[X | \hat{U}] - \mathbb{E}[X | U]\|_{L_2}. && \text{(Jensen's inequality)} \end{aligned}$$

Similarly, the second covariance estimation error can be upper bounded as

$$\begin{aligned} & \|B_{\cdot j} - \hat{B}_{\cdot j}\| \\ &= \|\mathbb{E}[\text{Cov}(X, \mathbb{E}[Y_j | X, U] | U) - \text{Cov}(X, \mathbb{E}[Y_j | X, \hat{U}] | \hat{U})]\| \\ &= \|\mathbb{E}[\mathbb{E}[X | U]\mathbb{E}[Y_j | X, U] - \mathbb{E}[X | \hat{U}]\mathbb{E}[Y_j | X, \hat{U}]]\| \\ &\leq \mathbb{E}[\|\mathbb{E}[X | U]\mathbb{E}[Y_j | X, U] - \mathbb{E}[X | \hat{U}]\mathbb{E}[Y_j | X, \hat{U}]\|] && \text{(Jensen's inequality)} \\ &\leq \mathbb{E}[\|\mathbb{E}[X | U](\mathbb{E}[Y_j | X, U] - \mathbb{E}[Y_j | X, \hat{U}])\| + \|(\mathbb{E}[X | U] - \mathbb{E}[X | \hat{U}])\mathbb{E}[Y_j | X, \hat{U}]\|] && \text{(triangle inequality)} \\ &= \mathbb{E}[\|\mathbb{E}[X | U]\|\|\mathbb{E}[Y_j | X, U] - \mathbb{E}[Y_j | X, \hat{U}]\| + \|\mathbb{E}[X | U] - \mathbb{E}[X | \hat{U}]\|\|\mathbb{E}[Y_j | X, \hat{U}]\|] \\ &\leq \|\mathbb{E}[X | U]\|_{L_2}\|\mathbb{E}[Y_j | X, \hat{U}] - \mathbb{E}[Y_j | X, U]\|_{L_2} + \|\mathbb{E}[Y_j | X, \hat{U}]\|_{L_2}\|\mathbb{E}[X | \hat{U}] - \mathbb{E}[X | U]\|_{L_2} && \text{(Cauchy-Schwarz inequality)} \\ &\leq \|X\|_{L_2}\|\mathbb{E}[Y_j | X, \hat{U}] - \mathbb{E}[Y_j | X, U]\|_{L_2} + \|Y_j\|_{L_2}\|\mathbb{E}[X | \hat{U}] - \mathbb{E}[X | U]\|_{L_2}. && \text{(Jensen's inequality)} \end{aligned}$$

Part (2) Coefficient estimation error in terms of covariance estimation errors. When $\|\mathbb{E}[X | \hat{U}] - \mathbb{E}[X | U]\|_{L_2} < \sigma/(2M)$, from Part (1) we have $\kappa(A)\|A - \hat{A}\|/\|A\| = \|A^{-1}\|\|A - \hat{A}\| < 1$. From Lemma D.1, we further have that

$$\begin{aligned} \max_{j \in \mathcal{C}^c} \|\tilde{\beta}_{\cdot j} - \beta_{\cdot j}\| &\leq \frac{\max_{j \in \mathcal{C}^c} \kappa(A) \left(\frac{\|\beta_{\cdot j}\|\|A - \hat{A}\|}{\|A\|} + \frac{\|B_{\cdot j} - \hat{B}_{\cdot j}\|}{\|A\|} \right)}{1 - \kappa(A)\frac{\|A - \hat{A}\|}{\|A\|}} \\ &\lesssim \|X\|_{L_2} \max_{j \in \mathcal{C}^c} \|\beta_{\cdot j}\| \|\mathbb{E}[Y_j | X, \hat{U}] - \mathbb{E}[Y_j | X, U]\|_{L_2} \\ &\quad + (\|X\|_{L_2} + \max_{j \in \mathcal{C}^c} \|Y_j\|_{L_2})\|\mathbb{E}[X | \hat{U}] - \mathbb{E}[X | U]\|_{L_2}, \end{aligned}$$

where in the last inequality, we use the boundedness of A 's spectrum from Assumption 4.

Part (3) Coefficient estimation error in terms of covariate estimation errors. From Lemma D.3, it follows that

$$\max_{j \in \mathcal{C}^c} \|\tilde{\beta}_{\cdot j} - \beta_{\cdot j}\| \lesssim \left(\|X\|_{L_2} (L_X^{\frac{1}{2}} + L_Y^{\frac{1}{2}}) + \max_{j \in \mathcal{C}^c} \|Y_j\|_{L_2} \right) \|v^{-1}(\hat{U}) - U\|_{L_2},$$

with $L_Y = \max_{j \in [p]} L_{Y_j}$. \square

D.2 Proof of Lemma 3 (linear models)

Proof of Lemma 3. For observations $(\mathbf{X}, \mathbf{Y}, \mathbf{U}) \in \mathbb{R}^{n \times d} \times \mathbb{R}^{n \times p} \times \mathbb{R}^{n \times r}$ and an estimate $\hat{\mathbf{U}} \in \mathbb{R}^{n \times \hat{r}}$ of \mathbf{U} , we have $S = \mathbf{X}^\top P_U^\perp \mathbf{X}$, $\tilde{S} = \mathbf{X}^\top P_{\hat{\mathbf{U}}}^\perp \mathbf{X}$, $\bar{\mathbf{Y}} = P_U^\perp \mathbf{Y}$, and $\tilde{\mathbf{Y}} = P_{\hat{\mathbf{U}}}^\perp \mathbf{Y}$. Furthermore, the regression coefficient on $(P_U^\perp \mathbf{X}, P_U^\perp \mathbf{Y})$ can be expressed as

$$b = (\mathbf{X}^\top P_U^\perp \mathbf{X})^{-1} \mathbf{X}^\top P_U^\perp \mathbf{Y} = S^{-1} \mathbf{X}^\top \bar{\mathbf{Y}} / n,$$

and the regression coefficient on $(P_{\hat{\mathbf{U}}}^\perp \mathbf{X}, P_{\hat{\mathbf{U}}}^\perp \mathbf{Y})$ can be expressed as

$$\tilde{b} = (\mathbf{X}^\top P_{\hat{\mathbf{U}}}^\perp \mathbf{X})^{-1} \mathbf{X}^\top P_{\hat{\mathbf{U}}}^\perp \mathbf{Y} = \tilde{S}^{-1} \mathbf{X}^\top \tilde{\mathbf{Y}} / n.$$

From Lemma D.1, we have

$$\max_{j \in \mathcal{C}^c} \|\tilde{b}_{\cdot j} - b_{\cdot j}\| \leq \frac{\kappa(S)}{\|S\|_{\text{op}}} \frac{\max_{j \in [p]} \|b_{\cdot j}\| \|\tilde{S} - S\| + \|\mathbf{X}^\top (\tilde{\mathbf{Y}}_{\cdot j} - \bar{\mathbf{Y}}_{\cdot j}) / n\|}{1 - \kappa(S) \frac{\|\tilde{S} - S\|}{\|S\|_{\text{op}}}}. \quad (14)$$

This requires verifying the assumptions therein. Specifically, we verify (17) below. Because

$$\begin{aligned} \|\tilde{S} - S\| &= \|\mathbf{X}^\top (P_{\hat{\mathbf{U}}}^\perp - P_U^\perp) \mathbf{X} / n\| \\ &\leq \|\mathbf{X}^\top \mathbf{X} / n\| \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\| \\ &= \|S\|_{\text{op}} \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\|, \end{aligned}$$

and $\kappa(S) \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\| < 1$ as assumed, we have

$$\frac{\|\tilde{S} - S\|}{\|S\|_{\text{op}}} < \frac{1}{\kappa(S)},$$

which verifies (17) of Lemma D.1. On the other hand, we also have

$$\|\mathbf{X}^\top (\tilde{\mathbf{Y}} - \bar{\mathbf{Y}})\|_{2,\infty} = \|\mathbf{X}^\top (P_{\hat{\mathbf{U}}}^\perp - P_U^\perp) \mathbf{Y}\|_{2,\infty} \leq \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\| \|\mathbf{X}\|_{\text{op}} \|\mathbf{Y}\|_{2,\infty}.$$

Therefore, (14) implies that

$$\begin{aligned} \max_{j \in \mathcal{C}^c} \|\tilde{b}_{\cdot j} - b_{\cdot j}\| &\leq \frac{\|S\|_{\text{op}} \|b\|_{2,\infty} + \|\mathbf{X}\|_{\text{op}} \|\mathbf{Y}\|_{2,\infty} / n}{\|S\|_{\text{op}}} \frac{\kappa(S) \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\|}{1 - \kappa(S) \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\|} \\ &\leq (\|b\|_{2,\infty} + \|S\|_{\text{op}}^{-\frac{1}{2}} \|\mathbf{Y}\|_{2,\infty} n^{-\frac{1}{2}}) \frac{\kappa(S) \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\|}{1 - \kappa(S) \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\|} \end{aligned}$$

whenever $\|S\|_{\text{op}} \neq 0$ and $\kappa(S) \|P_{\hat{\mathbf{U}}}^\perp - P_U^\perp\| < 1$. \square

D.3 Auxillary lemmas

Lemma D.1 (Backward error of perturbed linear systems). *Let $A \in \mathbb{R}^{n \times n}$ be nonsingular, $b \in \mathbb{R}^n$, and $x = A^{-1}b \in \mathbb{R}^n$. In the following, $\Delta A \in \mathbb{R}^{n \times n}$ and $\Delta b \in \mathbb{R}^n$ are some arbitrary matrix and vector. We assume that the norm on A satisfies $\|Ax\| \leq \|A\|\|x\|$ for all $A \in \mathbb{R}^{n \times n}$ and all $x \in \mathbb{R}^n$. Suppose $(A + \Delta A)\hat{x} = \hat{b}$ such that*

$$\hat{b} := b + \Delta b \neq 0 \quad (15)$$

$$\hat{x} := x + \Delta x \neq 0 \quad (16)$$

$$\frac{\|\Delta A\|}{\|A\|} < \frac{1}{\kappa(A)}, \quad (17)$$

where $\kappa(A) = \|A\|\|A^{-1}\|$ is the condition number of A . Then, it holds that

$$\|\Delta x\| \leq \frac{\|x\|\kappa(A)\frac{\|\Delta A\|}{\|A\|} + \kappa(A)\frac{\|\Delta b\|}{\|A\|}}{1 - \kappa(A)\frac{\|\Delta A\|}{\|A\|}}.$$

If further, $b \neq 0$ (or equivalently $x \neq 0$), then

$$\frac{\|\Delta x\|}{\|x\|} \leq \frac{\kappa(A)\left(\frac{\|\Delta A\|}{\|A\|} + \frac{\|\Delta b\|}{\|b\|}\right)}{1 - \kappa(A)\frac{\|\Delta A\|}{\|A\|}}.$$

Proof of Lemma D.1. We split the proof into two parts.

Part (1) We first show that when (17) is satisfied, $A + \Delta A$ must be nonsingular. If $A + \Delta A$ is singular, then exists nonzero v such that $(A + \Delta A)v = 0$. Since A is nonsingular, we have $A^{-1}\Delta Av = -v$. So

$$\|v\| = \|A^{-1}\Delta Av\| \leq \|A^{-1}\|\|\Delta A\|\|v\|,$$

which implies that

$$\|\Delta A\| \geq \frac{1}{\|A^{-1}\|}.$$

On the other hand, since $\kappa(A) = \|A\|\|A^{-1}\|$, from (17) we have

$$\frac{\|\Delta A\|}{\|A\|} < \frac{1}{\|A\|\|A^{-1}\|},$$

or equivalently,

$$\|\Delta A\| < \frac{1}{\|A^{-1}\|}.$$

This leads to contradictions. Therefore, $A + \Delta A$ must be nonsingular.

Part (2) Since $(A + \Delta A)\hat{x} = b + \Delta b$ and $Ax = b$, we have $A\Delta x + \Delta A\hat{x} = \Delta b$. So $\Delta x = A^{-1}(\Delta b - \Delta A\hat{x})$. Then we have

$$\begin{aligned} \frac{\|\Delta x\|}{\|\hat{x}\|} &= \frac{\|A^{-1}(\Delta b - \Delta A\hat{x})\|}{\|\hat{x}\|} \\ &\leq \frac{\|A^{-1}\|(\|\Delta A\|\|\hat{x}\| + \|\Delta b\|)}{\|\hat{x}\|} \end{aligned}$$

$$\begin{aligned}
&= \|A^{-1}\| \|A\| \left(\frac{\|\Delta A\|}{\|A\|} + \frac{\|\Delta b\|}{\|A\| \|\widehat{x}\|} \right) \\
&= \kappa(A) \left(\frac{\|\Delta A\|}{\|A\|} + \frac{\|\Delta b\|}{\|A\| \|\widehat{x}\|} \right),
\end{aligned}$$

and

$$\begin{aligned}
\|\Delta x\| &\leq \kappa(A) \left(\frac{\|\Delta A\|}{\|A\|} + \frac{\|\Delta b\|}{\|A\| \|\widehat{x}\|} \right) \|\widehat{x}\| \\
&= \kappa(A) \left(\frac{\|\Delta A\|}{\|A\|} \|\widehat{x}\| + \frac{\|\Delta b\|}{\|A\|} \right) \\
&\leq \kappa(A) \frac{\|\Delta A\|}{\|A\|} (\|x\| + \|\Delta x\|) + \kappa(A) \frac{\|\Delta b\|}{\|A\|}.
\end{aligned}$$

Rearrange the above inequality, we have

$$\begin{aligned}
\left(1 - \kappa(A) \frac{\|\Delta A\|}{\|A\|} \right) \|\Delta x\| &\leq \kappa(A) \frac{\|\Delta A\|}{\|A\|} \|x\| + \kappa(A) \frac{\|\Delta b\|}{\|A\|} \\
\|\Delta x\| &\leq \frac{\|x\| \kappa(A) \frac{\|\Delta A\|}{\|A\|} + \kappa(A) \frac{\|\Delta b\|}{\|A\|}}{1 - \kappa(A) \frac{\|\Delta A\|}{\|A\|}}.
\end{aligned}$$

When $x \neq 0$, we further have

$$\frac{\|\Delta x\|}{\|x\|} \leq \frac{\kappa(A) \frac{\|\Delta A\|}{\|A\|} + \kappa(A) \frac{\|\Delta b\|}{\|A\| \|\widehat{x}\|}}{1 - \kappa(A) \frac{\|\Delta A\|}{\|A\|}} \leq \frac{\kappa(A) \left(\frac{\|\Delta A\|}{\|A\|} + \frac{\|\Delta b\|}{\|A\|} \right)}{1 - \kappa(A) \frac{\|\Delta A\|}{\|A\|}},$$

where the last inequality holds since $\|b\| = \|Ax\| \leq \|A\| \|x\|$. \square

Lemma D.2. Suppose X, Y are two random vectors in \mathbb{R}^d defined on probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a \mathcal{F} -measurable and satisfies the L -Lipschitz condition (in ℓ_q -norm) almost surely. Then it holds that

$$\|f(X) - f(Y)\|_{L_q} \leq L^{1/q} \|X - Y\|_{L_q}.$$

Proof of Lemma D.2. Note that

$$\begin{aligned}
\|f(X) - f(Y)\|_{L_q}^q &= \int |f(X) - f(Y)|^q d\mathbb{P} \\
&\leq \int L \|X - Y\|_q^q d\mathbb{P} \quad (\text{Lipschitz condition}) \\
&= L \sum_{j=1}^d \int |X_j - Y_j|^q d\mathbb{P} \\
&= L \|X - Y\|_{L_q}^q
\end{aligned}$$

Then the conclusion follows by taking the q^{-1} -power on both sides. \square

Lemma D.3 (Error bound of regression function with estimated covaraites). On a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$, consider a random vector W and a sequence of random vectors $\{V_m\}_{m \in \mathbb{N}}$ adapted to a filtration $\{\mathcal{F}_m\}_{m \in \mathbb{N}}$ such that $\mathcal{F}_m \subseteq \mathcal{F}_{m+1}$. Suppose that (i) $\|W\|_{L_2} < \infty$, (ii)

$V_m \xrightarrow{\text{a.s.}} V$, and (iii) the function $h(v) = \mathbb{E}[W \mid V = v]$ satisfies the L -Lipschitz condition in ℓ_2 -norm almost surely. Then, under (i)-(ii), it holds

$$\mathbb{E}[W \mid V_m] \xrightarrow{L_2} \mathbb{E}[W \mid V],$$

and under (i)-(iii), it holds that

$$\|\mathbb{E}[W \mid V_m] - \mathbb{E}[W \mid V]\|_{L_2} \leq 2L^{\frac{1}{2}}\|V_m - V\|_{L_2}.$$

Proof of Lemma D.3. Define $\mathcal{F}_\infty = \sigma(\cup_m \mathcal{F}_m)$. There exists some \mathcal{F}_∞ -measurable function h and \mathcal{F}_m -measurable function h_m such that $h(V) = \mathbb{E}[W \mid V]$ and $h_m(V_m) = \mathbb{E}[W \mid V_m]$ almost surely. Notice that $(\mathbb{E}[W \mid V_m])_{m \in \mathbb{N}}$ is a Doob martingale (because $\|W\|_{L_1} < \infty$). From martingale convergence theorem, there exists $V_\infty = \mathbb{E}[W \mid \mathcal{F}_\infty]$ that is measurable with respect to \mathcal{F}_∞ such that $\|V_\infty\|_{L_1} < \infty$ and $\mathbb{E}[W \mid V_m] \xrightarrow{\text{a.s.}} V_\infty$. On the other hand, because $V_m \xrightarrow{\text{a.s.}} V$ from Assumption (ii), we know that $V \equiv V_\infty$ is \mathcal{F}_∞ -measurable. This implies that $\mathbb{E}[W \mid V_\infty] = h(V_\infty) = h(V) = \mathbb{E}[W \mid V]$ almost surely. Thus, we conclude that $\mathbb{E}[W \mid V_m] \xrightarrow{\text{a.s.}} \mathbb{E}[W \mid V]$. By Jensen's inequality and Assumption (ii), we have $\|\mathbb{E}[W \mid V_m]\|_{L_2} \leq \|W\|_{L_2} < \infty$, which implies that the set of functions $\{\mathbb{E}[W \mid V_m] : m \in \mathbb{N}\}$ is uniformly integrable. Thus, we further have $\mathbb{E}[W \mid V_m] \xrightarrow{L_2} \mathbb{E}[W \mid V]$ from dominated convergence theorem.

Next, we need to derive the convergence rate. We have that

$$\|h_m(V_m) - h(V)\|_{L_2} \leq \|h_m(V_m) - h(V_m)\|_{L_2} + \|h(V_m) - h(V)\|_{L_2}. \quad (18)$$

For the first term in (18), from the martingale property, the function representation $h_m(V_m) = \mathbb{E}[h(V) \mid \mathcal{F}_m]$ gives that

$$\|h_m(V_m) - h(V_m)\|_{L_2} = \|\mathbb{E}[h(V) \mid \mathcal{F}_m] - h(V_m)\|_{L_2} \leq \|h(V) - h(V_m)\|_{L_2} \quad (19)$$

where the last inequality is from Jensen's inequality.

Combining (18) and (19) yields that

$$\|h_m(V_m) - h(V)\|_{L_2} \leq 2\|h(V_m) - h(V)\|_{L_2} \leq 2L^{\frac{1}{2}}\|V_m - V\|_{L_2},$$

where the last inequality is from Lemma D.2 by noting that h satisfies the L -Lipschitz condition from Assumption (iii). \square

E Doubly robust semiparametric inference

E.1 Proof of Theorem 4 and Corollary 5

Proof of Theorem 4 and Corollary 5. Theorem 4 is a special case of Theorem E.1 with nonlinear link functions. The proof follows by applying Theorem E.1 with g being identity. Meanwhile, the assumption in Theorem E.1 can be relaxed under this special case by noting that $\mathbb{E}[Y | X, U]$ can be replaced by Y because the residual $Y - \mathbb{E}[Y | X, U]$ is orthogonal to mean-zero functions of (X, U) in the L_2 space so that $\eta(O) = Y - \mathbb{E}[Y | U]$ under identity link. \square

E.2 Proof of Proposition 6

Proof of Proposition 6. From Theorem 4, we have

$$\sqrt{n}(\tilde{b} - \tilde{\beta}) = \sqrt{n}\tilde{\Sigma}^{-1}(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}(O; \mathbb{P})\} + \xi,$$

where the remainder term ξ satisfies that $\|\xi\|_{2,\infty} = o_{\mathbb{P}}(1)$ under the rate conditions in Theorem 4. Recall that $t_j = \sqrt{n}\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\}^{\frac{1}{2}}\hat{\Sigma}^{-1}(\tilde{b}_j - \tilde{\beta}_j)$. We have that

$$\begin{aligned} v^\top t_j &= \sqrt{n}v^\top \mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}^{\frac{1}{2}}(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} \\ &\quad + (\sqrt{n}v^\top (\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\}^{\frac{1}{2}} - \mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}^{\frac{1}{2}})(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} + v^\top \mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\}^{\frac{1}{2}}\hat{\Sigma}^{-1}\xi_{\cdot j}) \\ &=: \vartheta_j + \varsigma_j. \end{aligned}$$

For the first component, note that ϑ_j for $j = 1, \dots, p$ are independent conditional on (X, U) 's. Furthermore, the self-normalizing term $\vartheta_j \xrightarrow{d} \mathcal{N}(0, 1)$ for $j \in \mathcal{N}_p$. By the strong law of large number, $\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} \xrightarrow{\text{a.s.}} \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}$, $\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\} \xrightarrow{\text{a.s.}} \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}$ and $\hat{\Sigma} \xrightarrow{\text{a.s.}} \tilde{\Sigma}$ when both $m, n, p \rightarrow \infty$. This implies that $\max_j \|\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\} - \mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}\|_{\text{op}} \xrightarrow{\text{a.s.}} 0$, $\max_j \|\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\} - \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}\|_{\text{op}} \xrightarrow{\text{a.s.}} 0$, $\|\hat{\Sigma} - \tilde{\Sigma}\|_{\text{op}} \xrightarrow{\text{a.s.}} 0$, which follows from Patil et al. (2023, Lemma S.8.6 (1)) by noting that the variables O 's are iid in the triangular array. For the second component, we have that

$$\begin{aligned} \max_{1 \leq j \leq p} |\varsigma_j| &\leq \max_{1 \leq j \leq p} \|\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\}^{\frac{1}{2}} - \mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}^{\frac{1}{2}}\|_{\text{op}} \cdot \|\sqrt{n}(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\}\|_2 \\ &\quad + \max_{1 \leq j \leq p} \|\mathbb{V}_n\{\tilde{\varphi}_{\cdot j}(O; \hat{\mathbb{P}})\}^{\frac{1}{2}}\hat{\Sigma}^{-1}\|_{\text{op}} \|\xi_{\cdot j}\|_2 \\ &= o_{\mathbb{P}}(1)\mathcal{O}_{\mathbb{P}}(1) + \mathcal{O}_{\mathbb{P}}(1)o_{\mathbb{P}}(1) \\ &= o_{\mathbb{P}}(1). \end{aligned}$$

Let $\varrho = |\mathcal{N}_p|^{-1} \sum_{j \in \mathcal{N}_p} \mathbb{1}\{|v^\top t_j| > z_{\frac{\alpha}{2}}\}$. To prove the overall Type-I error control, we will show the expectation that ϱ tends to α , and its variance tends to zero. For the expectation, for any $\epsilon > 0$, we have

$$\begin{aligned} \mathbb{E}[\varrho] &= \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}\left(|v^\top t_j| > z_{\frac{\alpha}{2}}\right) \\ &\leq \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \left[\mathbb{P}\left(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon\right) + \mathbb{P}(|\varsigma_j| > \epsilon) \right] \\ &= \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}\left(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon\right) + \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}(|\varsigma_j| > \epsilon) \\ &\leq \frac{1}{|\mathcal{N}_p|} \sum_{j \in \mathcal{N}_p} \mathbb{P}\left(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon\right) + \mathbb{P}\left(\max_{1 \leq j \leq p} |\varsigma_j| > \epsilon\right) \rightarrow 2\left(1 - \Phi\left(z_{\frac{\alpha}{2}} - \epsilon\right)\right), \end{aligned}$$

where the last convergence holds because the Cesaro mean converges to the same limit as

$$\lim_{n,p} \mathbb{P} \left(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon \right) = 2 \left(1 - \Phi \left(z_{\frac{\alpha}{2}} - \epsilon \right) \right),$$

while the term $\mathbb{P}(\max_{1 \leq j \leq p} |\varsigma_j| > \epsilon)$ vanishes. Similarly, we can show that $\liminf_{n,p \rightarrow \infty} \mathbb{E}[\varrho] \geq 2 \left(1 - \Phi \left(z_{\frac{\alpha}{2}} - \epsilon \right) \right)$ for all $\epsilon > 0$. Let $\epsilon \rightarrow 0^+$, it follows that $\mathbb{E}[\varrho] \rightarrow \alpha$ as $n, p \rightarrow \infty$.

For any $\epsilon > 0$, the second moment can be upper bounded as below:

$$\begin{aligned} \mathbb{E}[\varrho^2] &= \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p} \mathbb{P} \left(|v^\top t_j| > z_{\frac{\alpha}{2}}, |v^\top t_k| > z_{\frac{\alpha}{2}} \right) \\ &= \frac{1}{|\mathcal{N}_p|^2} \sum_{j \in \mathcal{N}_p} \mathbb{P} \left(|v^\top t_j| > z_{\frac{\alpha}{2}} \right) + \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P} \left(|v^\top t_j| > z_{\frac{\alpha}{2}}, |v^\top t_k| > z_{\frac{\alpha}{2}} \right) \\ &\leq \frac{1}{|\mathcal{N}_p|^2} \sum_{j \in \mathcal{N}_p} \mathbb{P} \left(|v^\top t_j| > z_{\frac{\alpha}{2}} \right) \\ &\quad + \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P} \left(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon, |\vartheta_k| > z_{\frac{\alpha}{2}} - \epsilon \right) + \mathbb{P}(|\varsigma_j| > \epsilon) + \mathbb{P}(|\varsigma_k| > \epsilon) \\ &= \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P} \left(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon, |\vartheta_k| > z_{\frac{\alpha}{2}} - \epsilon \right) + o(1) \\ &= \frac{1}{|\mathcal{N}_p|^2} \sum_{j,k \in \mathcal{N}_p, j \neq k} \mathbb{P}(|\vartheta_j| > z_{\frac{\alpha}{2}} - \epsilon \mid X, \widehat{U}) \mathbb{P}(|\vartheta_k| > z_{\frac{\alpha}{2}} - \epsilon \mid X, \widehat{U}) + o(1) \\ &\rightarrow 4 \left(1 - \Phi \left(z_{\frac{\alpha}{2}} - \epsilon \right) \right)^2, \end{aligned}$$

where the last equality is from the independence of ϑ_j and ϑ_k . We can similarly obtain the lower bound. Let $\epsilon \rightarrow 0^+$, it follows that $\mathbb{E}[\varrho^2] \rightarrow \alpha^2$ and $\mathbb{V}(\varrho) \rightarrow 0$ as $n, p \rightarrow \infty$. Combining the previous results yields that $\varrho \xrightarrow{P} \alpha$. \square

E.3 Nonlinear modeling

The natural extension of partial linear models to the nonlinear cases is the generalized partially linear models (Severini and Staniswalis, 1994; Härdle et al., 1998):

$$g(\mathbb{E}[Y \mid X, U]) = \beta^\top X + h(U), \quad (20)$$

by introducing a proper link function g , applied element-wisely on the conditional mean of the outcomes. Similar to the results in the previous sections, a nonlinear counterpart of the main effect estimand (5) is given by

$$\beta(\mathbb{P}) = \mathbb{E}[\text{Cov}(X \mid U)]^{-1} \mathbb{E}[\text{Cov}[X, g(\mathbb{E}(Y \mid X, U)) \mid U]]. \quad (21)$$

Such an estimand has been considered in Robins et al. (2008); Newey and Robins (2018) with the identity link and in Vansteelandt and Dukes (2022) with a single treatment. When the model (20) is correctly specified, (21) is equivalent to the regression coefficient under model (20). On the other hand, when the model (20) is misspecified, estimand (21) still represents a meaningful statistical quantity.

With a differentiable link function g , the influence function (for $\tilde{\Sigma}\tilde{\beta}$) analogous to (9) is given by:

$$\tilde{\varphi}(O; \mathbb{P}) := (X - \mathbb{E}[X \mid \widehat{U}])(\eta(O) - \tilde{\beta}^\top(X - \mathbb{E}[X \mid \widehat{U}]))^\top,$$

where the main effect estimand with estimated embedding is defined as:

$$\tilde{\beta} = \mathbb{E}[\text{Cov}(X | \hat{U})]^{-1} \mathbb{E}[\text{Cov}[X, g(\mathbb{E}(Y|X, \hat{U})) | \hat{U}]], \quad (22)$$

and the function η is defined as:

$$\eta(O) = g'(\mathbb{E}[Y | X, \hat{U}]) \odot (Y - \mathbb{E}[Y | X, \hat{U}]) + g(\mathbb{E}[Y | X, \hat{U}]) - \mathbb{E}[g(\mathbb{E}[Y | X, \hat{U}]) | \hat{U}].$$

The doubly robust semiparametric inference results in Theorem 4 and Corollary 5 can be extended to accommodate nonlinear link functions, as shown in the next theorem.

Theorem E.1 (Doubly robust inference with nonlinear link functions). *Under a nonparametric model and a differentiable link function g , define the estimator of β in (21) as:*

$$\hat{\beta} = \mathbb{P}_n\{(X - \hat{\mathbb{E}}(X | \hat{U}))^{\otimes 2}\}^{-1} \mathbb{P}_n\{(X - \hat{\mathbb{E}}(X | \hat{U})) \cdot (\mathbb{I} - \mathbb{P}_n)\{g(\hat{\mathbb{E}}[Y | X, \hat{U}])\}^\top\}, \quad (23)$$

which depends on empirical measure \mathbb{P}_n and two nuisance functions $\hat{\mathbb{E}}[X | \hat{U}]$ and $\hat{\mathbb{E}}[Y | X, \hat{U}]$ estimated from independent samples of \mathbb{P}_n . Under Assumptions 4 and 6 and assume that

- (i) *Local Lipschitzness: There exists $L > 0$ such that $\|g(\mathbb{E}[Y | X, \hat{U}]) - g(\hat{\mathbb{E}}[Y | X, \hat{U}]) - g'(\hat{\mathbb{E}}[Y | X, \hat{U}]) \odot (\mathbb{E}[Y | X, \hat{U}] - \hat{\mathbb{E}}[Y | X, \hat{U}])\|_\infty \leq L \|\mathbb{E}[Y | X, \hat{U}] - \hat{\mathbb{E}}[Y | X, \hat{U}]\|_\infty^2$.*
- (ii) *Boundeness and consistency: Assumptions 4 and 6 hold with additionally, $\|\eta(O)\|_{L_{2(1+\delta^{-1})}} < M$ and $\|\hat{\eta}(O) - \eta(O)\|_\infty\|_{L_{2(1+\delta)}} = o_{\mathbb{P}}(1)$.*
- (iii) *Rate condition: $\|\mathbb{E}[X | \hat{U}] - \hat{\mathbb{E}}[X | \hat{U}]\|_{L_2}^2$, $\|\mathbb{E}[Y | X, \hat{U}] - \hat{\mathbb{E}}[Y | X, \hat{U}]\|_{L_2, \infty}^2$, and $\|\mathbb{E}[g(\mathbb{E}[Y | X, \hat{U}]) | \hat{U}] - \hat{\mathbb{E}}[g(\hat{\mathbb{E}}[Y | X, \hat{U}]) | \hat{U}]\|_{L_2, \infty} \|\mathbb{E}[X | \hat{U}] - \hat{\mathbb{E}}[X | \hat{U}]\|_{L_2}$ are of order $o_{\mathbb{P}}(n^{-\frac{1}{2}})$.*

Then, the estimator \tilde{b} is asymptotically normal:

$$\sqrt{n}(\tilde{b}_{\cdot j} - \tilde{\beta}_{\cdot j}) \xrightarrow{d} \mathcal{N}_d(0, \tilde{\Sigma}^{-1} \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} \tilde{\Sigma}^{-1}), \quad j = 1, \dots, p.$$

Furthermore, if the conditions of Theorem 2 hold with $\ell_m = o(n^{-\frac{1}{2}})$, then we have

$$\sqrt{n}(\tilde{b}_{\cdot j} - \beta_{\cdot j}) \xrightarrow{d} \mathcal{N}_d(0, \tilde{\Sigma}^{-1} \mathbb{V}\{\tilde{\varphi}_{\cdot j}(O; \mathbb{P})\} \tilde{\Sigma}^{-1}), \quad j = 1, \dots, p.$$

Compared to Theorem 4, Theorem E.1 requires additional assumptions regarding the Lipschitzness of the link function around the true regression function, as noted by Vansteelandt and Dukes (2022). It also requires boundedness and consistency assumptions on the first-order expansion term η . Nevertheless, the overall conclusion is similar when both the estimators and the influence functions have a different link function for a different target estimand. The double robustness still allows efficient semiparametric inference with data-adaptive estimation procedures.

Proof of Theorem E.1. From Lemma E.3, we have

$$\sqrt{n}(\tilde{b} - \tilde{\beta}) = \sqrt{n}\mathbb{P}\{(X - \mathbb{E}(X | \hat{U}))^{\otimes 2}\}^{-1}(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}(O; \mathbb{P})\} + \tilde{\xi}$$

where $\tilde{\varphi}$ is defined as

$$\tilde{\varphi}(O; \mathbb{P}) = (X - \mathbb{E}[X | \hat{U}])(\eta(O) - \tilde{\beta}^\top(X - \mathbb{E}[X | \hat{U}]))^\top \quad (24)$$

and the remainder term $\tilde{\xi}$ satisfies that $\|\tilde{\xi}\|_{2, \infty} = o_{\mathbb{P}}(1)$. This proves the first statement.

Algorithm E.3 Semiparametric inference for main effects with nonlinear link functions

Input: Responses Y , covariate X , estimated latent embedding \widehat{U} , and link function g .

- 1: Use machine learning methods to obtain nuisance estimates $\widehat{\mathbb{E}}[Y | X, \widehat{U}]$ and $\widehat{\mathbb{E}}[X | \widehat{U}]$.
- 2: Use a data-adaptive fit $g(\widehat{\mathbb{E}}[Y | X, \widehat{U}]) \sim \widehat{U}$ to obtain estimated regression function $\widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, \widehat{U}]) | \widehat{U}]$. If X is categorical with finite support $|\mathcal{X}| < \infty$, this simply reduces to $\widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, \widehat{U}]) | \widehat{U}] = \sum_{x \in \mathcal{X}} g(\widehat{\mathbb{E}}[Y | X = x, \widehat{U}]) \widehat{\mathbb{E}}[X = x | \widehat{U}]$.
- 3: Fit a linear regression of $\widehat{\eta}(O) \sim X - \widehat{\mathbb{E}}[X | \widehat{U}]$ without an intercept to obtain an estimate \tilde{b} as defined in (23) of $\tilde{\beta}$ as defined in (22).
- 4: Estimate the variance of $\tilde{b}_{.j}$ by \widehat{S}_j/n based on Theorem E.1, where $\widehat{S}_j = \widehat{\Sigma}^{-1} \mathbb{V}_n \{\tilde{\varphi}_{.j}(O; \widehat{\mathbb{P}})\} \widehat{\Sigma}^{-1}$.

Output: Confidence intervals and p-values based on asymptotic null distribution $\tilde{b}_{.j} \stackrel{d}{\sim} \mathcal{N}_d(\tilde{\beta}_{.j}, \frac{\widehat{S}_j}{n})$.

When $\|\widehat{U} - U\|_{L_2} = o_{\mathbb{P}}(n^{-\frac{1}{2}})$, from Theorem 2 we have $\|\tilde{\beta} - \beta\|_{2,\infty} = o_{\mathbb{P}}(n^{-\frac{1}{2}})$. Therefore, we further have

$$\sqrt{n}(\tilde{b} - \beta) = \sqrt{n}(\tilde{b} - \tilde{\beta}) + \sqrt{n}(\tilde{\beta} - \beta) = \sqrt{n}\mathbb{P}\{(X - \mathbb{E}(X | \widehat{U}))^{\otimes 2}\}^{-1}(\mathbb{P}_n - \mathbb{P})\{\tilde{\varphi}(O; \mathbb{P})\} + \xi,$$

with $\|\xi\|_{2,\infty} = o_{\mathbb{P}}(1)$.

To establish the asymptotic normality, we apply the triangle-array CLT in Lemma E.4. This requires verifying the sufficient condition of the Lindeberg condition. Because $\mathbb{V}\{\widetilde{\Sigma}^{-1}\tilde{\varphi}_{.j}(O; \mathbb{P})\} = \widetilde{\Sigma}^{-1}\mathbb{V}\{\tilde{\varphi}_{.j}(O; \mathbb{P})\}\widetilde{\Sigma}^{-1}$, we have

$$\begin{aligned} & \mathbb{E}[\|\mathbb{V}\{\widetilde{\Sigma}^{-1}\tilde{\varphi}_{.j}(O; \mathbb{P})\}^{-\frac{1}{2}}(\widetilde{\Sigma}^{-1}\tilde{\varphi}_{.j}(O; \mathbb{P}))\|^{2+\frac{2}{\delta}}] \\ &= \mathbb{E}[\|\mathbb{V}\{\tilde{\varphi}_{.j}(O; \mathbb{P})\}^{-\frac{1}{2}}\tilde{\varphi}_{.j}(O; \mathbb{P})\|^{2+\frac{2}{\delta}}] \\ &\leq \|\mathbb{V}\{\tilde{\varphi}_{.j}(O; \mathbb{P})\}^{-\frac{1}{2}}\|_{op}^{2+\frac{2}{\delta}} \cdot \mathbb{E}[\|\tilde{\varphi}_{.j}(O; \mathbb{P})\|^{2+\frac{2}{\delta}}] \\ &\leq \|\mathbb{V}\{\tilde{\varphi}_{.j}(O; \mathbb{P})\}^{-\frac{1}{2}}\|_{op}^{2+\frac{2}{\delta}} \cdot (\mathbb{E}[\|(X - \mathbb{E}[X | \widehat{U}])\eta_j(O)\|^{2+\frac{2}{\delta}}] + \mathbb{E}[\|(X - \mathbb{E}[X | \widehat{U}])^{\otimes 2}\tilde{\beta}_{.j}\|^{2+\frac{2}{\delta}}]) \\ &\leq \|\mathbb{V}\{\tilde{\varphi}_{.j}(O; \mathbb{P})\}^{-\frac{1}{2}}\|_{op}^{2+\frac{2}{\delta}} \cdot \mathbb{E}[\|X - \mathbb{E}[X | \widehat{U}]\|^{1+\frac{1}{\delta}}\|\eta(O)\|^{1+\frac{1}{\delta}}] + \|\tilde{\beta}_{.j}\|^{2+\frac{2}{\delta}} \\ &\leq \sigma^{-1-\frac{1}{\delta}} M^{2+\frac{2}{\delta}} + \|\tilde{\beta}_{.j}\|^{2+\frac{2}{\delta}}. \end{aligned}$$

Now applying Lemma E.4 finishes the proof. \square

E.4 Auxillary lemmas

Lemma E.2 (Efficient influence function). *Consider a random variable $O = (X, U, Y) \in \mathbb{R}^d \times \mathbb{R}^r \times \mathbb{R}^p$ under a nonparameteric model and a differentiable function g , the main effect estimand in $\mathbb{R}^{d \times p}$:*

$$\beta = \mathbb{E}[\text{Cov}(X | U)]^{-1} \mathbb{E}[\text{Cov}(X, g(\mathbb{E}[Y | X, U]) | U)],$$

(where g is applied entry-wisely) has an efficient influence function $\mu : \mathbb{R}^r \times \mathbb{R}^d \times \mathbb{R}^p \rightarrow \mathbb{R}^{d \times p}$ given by:

$$\varphi(O) = \mathbb{E}[\text{Cov}(X | U)]^{-1}(X - \mathbb{E}[X | U])(\eta(O) - \beta^\top(X - \mathbb{E}[X | U]))^\top,$$

where $\eta : \mathbb{R}^r \times \mathbb{R}^d \times \mathbb{R}^p \rightarrow \mathbb{R}^p$ is defined as:

$$\eta(O) = g'(\mathbb{E}[Y | X, U]) \odot (Y - \mathbb{E}[Y | X, U]) + g(\mathbb{E}[Y | X, U]) - \mathbb{E}[g(\mathbb{E}[Y | X, U]) | U].$$

Proof of Lemma E.2. The proof follows similarly as in Vansteelandt and Dukes (2022, Theorem 1) for a univariate treatment and a univariate outcome, and extends the previous results to the multivariate cases. Below, we present a simplified derivation of the influence function.

Under the nonparametric model for the observed data $O = (X, U, Y)$. We first calculate the efficient influence function of

$$\begin{aligned}\theta(\beta) &= \mathbb{E}[(X - \mathbb{E}[X | U])(g(\mathbb{E}[Y | X, U]) - \beta^\top(X - \mathbb{E}[X | U]))^\top] \\ &= \int(X - \mathbb{E}[X | U])(g(\mathbb{E}[Y | X, U]) - \beta^\top X) dP(O),\end{aligned}$$

where $P(O)$ is the joint distribution of data. Note that by the definition of β , we have $\theta(\beta) = 0$.

Consider a one-dimensional submodel of $p(O)$ indexed by a scalar parameter t , and let $S_t(o) = \partial \log dP_t(o)/\partial t|_{t=0}$ denote the score function of the submodel. Similarly, let $S_t(Y | X, U)$, $S_t(X | U)$ and $S_t(U)$ be the scores w.r.t. t in that parametric submodel, corresponding to the distributions $p(Y | X, U)$, $p(X | U)$ and $p(U)$, respectively. Taking the derivative of θ w.r.t. t , we obtain

$$\begin{aligned}\frac{\partial \theta(\beta)}{\partial t}|_{t=0} &= \int \frac{\partial(X - \mathbb{E}_t[X | U])}{\partial t}|_{t=0}(g(\mathbb{E}[Y | X, U]) - \beta^\top X)^\top dP(O) \\ &\quad + \int(X - \mathbb{E}[X | U])(g'(\mathbb{E}[Y | X, U]) \odot \frac{\partial \mathbb{E}[Y | X, U]}{\partial t}|_{t=0})^\top dP(O) \\ &\quad + \int(X - \mathbb{E}[X | U])(g(\mathbb{E}[Y | X, U]) - \beta^\top X)^\top \frac{\partial p_t(X, U)}{\partial t}|_{t=0} dO \\ &= - \int(X - \mathbb{E}[X | U])\mathbb{E}[g(\mathbb{E}[Y | X, U]) - \beta^\top X | U]^\top S_t(X | U) dP(O) \\ &\quad + \int(X - \mathbb{E}[X | U])(g'(\mathbb{E}[Y | X, U]) \odot (Y - \mathbb{E}[Y | X, U]))^\top S_t(Y | X, U) dP(O) \\ &\quad + \int(X - \mathbb{E}[X | U])(g(\mathbb{E}[Y | X, U]) - \beta^\top X)^\top S_t(X, U) dP(O),\end{aligned}$$

where in the first equality, we apply the product and chain rules (Kennedy, 2022, Section 3.4.3); and in the second equality, we use the identity $S_t(Z) = \partial \log p_t(Z)/\partial t = (\partial p_t(Z)/\partial t)/p_t(Z)$ for score functions.

Note that

$$S_t(O) = S_t(Y | X, U) + S_t(X | U) + S_t(U).$$

From the zero mean properties of scores and $\theta(\beta) = 0$, we further have

$$\begin{aligned}\frac{\partial \theta(\beta)}{\partial t}|_{t=0} &= - \int(X - \mathbb{E}[X | U])\mathbb{E}[g(\mathbb{E}[Y | X, U]) - \beta^\top X | U]^\top S_t(O) dP(O) \\ &\quad + \int(X - \mathbb{E}[X | U])(g'(\mathbb{E}[Y | X, U]) \odot (Y - \mathbb{E}[Y | X, U]))^\top S_t(O) dP(O) \\ &\quad + \int(X - \mathbb{E}[X | U])(g(\mathbb{E}[Y | X, U]) - \beta^\top X)^\top S_t(O) dP(O) \\ &= \int(X - \mathbb{E}[X | U])(\eta(O) - \beta^\top(X - \mathbb{E}[X | U]))^\top S_t(O) dP(O),\end{aligned}$$

which implies that $(X - \mathbb{E}[X | U])(\eta(O) - \beta^\top(X - \mathbb{E}[X | U]))^\top$ is an influence function for θ . From a similar argument in the proof of Theorem 1 in Vansteelandt and Dukes (2022), it is also the efficient influence function of $\theta(\beta)$ under the nonparametric model. Consequently, by chain rule $\partial \theta / \partial t = (\partial \theta / \partial \beta)(\partial \beta / \partial t)$, the conclusion follows by taking the inverse of $\partial \theta / \partial \beta$. \square

Remark 7 (Alternative expression of the estimand). Note that the first part of the influence function also gives an alternative expression for β :

$$\beta = \mathbb{E}[\text{Cov}(X | U)]^{-1} \mathbb{E}[(X - \mathbb{E}[X | U])\eta(O)^\top] \tag{25}$$

because

$$\mathbb{E}[(X - \mathbb{E}[X | U])(g'(\mathbb{E}[Y | X, U]) \odot (Y - \mathbb{E}[Y | X, U]))^\top] = 0, \quad (26)$$

by the law of iterated expectation.

Lemma E.3 (Doubly robust estimation). *Consider the setting in Lemma E.2. Define a plug-in estimator of β :*

$$\widehat{\beta} = \mathbb{P}_n\{(X - \widehat{\mathbb{E}}(X | U))^2\}^{-1}\mathbb{P}_n\{(X - \widehat{\mathbb{E}}(X | U)) \cdot (\mathbb{I} - \mathbb{P}_n)\{g(\widehat{\mathbb{E}}[Y | X, U])\}^\top\}$$

which depends on empirical measure \mathbb{P}_n and two nuisance functions $\widehat{\mathbb{E}}[X | U]$ and $\widehat{\mathbb{E}}[Y | X, U]$ estimated from independent samples of \mathbb{P}_n . Define the population and empirical variance by

$$\begin{aligned}\Sigma &:= \mathbb{P}\{(X - \mathbb{E}(X | U))^{\otimes 2}\} \\ \widehat{\Sigma} &:= \mathbb{P}_n\{(X - \widehat{\mathbb{E}}(X | U))^{\otimes 2}\},\end{aligned}$$

the empirical influence function (for $\Sigma\beta$) by:

$$\varphi(O; \widehat{\mathbb{P}}) := (X - \widehat{\mathbb{E}}[X | U])(\widehat{\eta}(O) - \widehat{\beta}^\top(X - \widehat{\mathbb{E}}[X | U]))^\top.$$

Suppose the following conditions hold:

- (Regularity conditions) There exists $\sigma > 0$ such that $\Sigma \succeq \sigma I_d$, $\widehat{\Sigma} \succeq \sigma I_d$.
- (Bounded moments and consistency) There exists $\delta \in (0, 1]$ and $M > 0$, such that

$$\begin{aligned}\|\beta\|_{2,\infty} \vee \|X - \mathbb{E}[X | U]\|_{L_{2(1+\delta^{-1})}} \vee \|X - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta^{-1})}} \vee \|\eta(O)\|_{L_{2(1+\delta^{-1})}} &< M \\ \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} \vee \|\widehat{\eta}(O) - \eta(O)\|_\infty &= o_{\mathbb{P}}(1)\end{aligned}$$

- (Local Lipschitzness) There exists $L > 0$ such that

$$\begin{aligned}\|g(\mathbb{E}[Y | X, U]) - g(\widehat{\mathbb{E}}[Y | X, U]) - g'(\widehat{\mathbb{E}}[Y | X, U]) \odot (\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U])\|_\infty \\ \leq L\|\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U]\|_\infty^2.\end{aligned} \quad (27)$$

Then, it holds that

$$\sqrt{n}(\widehat{\beta} - \beta) = \sqrt{n}\Sigma^{-1}(\mathbb{P}_n - \mathbb{P})\{\varphi(O; \mathbb{P})\} + \xi,$$

where for any $\epsilon > 0$, there exists a constant $C = C(\epsilon, \sigma, M, L)$, such that with probability at least $1 - 3\epsilon$, the remainder term satisfies that

$$\begin{aligned}\|\xi\|_{2,\infty} &\leq C\{ \|(\mathbb{P}_n - \mathbb{P})\{(X - \mathbb{E}[X | U])^{\otimes 2}\}\|_{\text{op}} + \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} + \|\|\eta(O) - \widehat{\eta}(O)\|_\infty\|_{L_{2(1+\delta)}} \} \\ &\quad + C\sqrt{n}\{\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2 \\ &\quad + ML\|\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U]\|_{L_2,\infty}^2 \\ &\quad + \|\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U]\|_{L_2,\infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}\},\end{aligned}$$

When $\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2$, $\|\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U]\|_{L_2,\infty}^2$, and $\|\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U]\|_{L_2,\infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}$ are of order $o_{\mathbb{P}}(n^{-\frac{1}{2}})$, we further have that $\|\xi\|_{2,\infty} = o_{\mathbb{P}}(1)$ and hence

$$\sqrt{n}(\widehat{\beta}_{\cdot j} - \beta_{\cdot j}) \xrightarrow{d} \mathcal{N}_d(0, \Sigma^{-1}\mathbb{V}\{\varphi_{\cdot j}(O; \mathbb{P})\}), \quad j = 1, \dots, p.$$

Proof of Lemma E.3. From the definition of $\widehat{\beta}$, we have $\mathbb{P}_n\{\varphi(O; \widehat{\mathbb{P}})\} = 0$. Therefore, $\widehat{\beta}$ is also a one-step estimator. We begin with a three-term decomposition of the estimation error (see, for example, Du et al. (2024b, Equation (2.2)) and Kennedy (2022, Equation (10))):

$$\begin{aligned}\widehat{\Sigma}\sqrt{n}(\widehat{\beta} - \beta) &= \sqrt{n}(\mathbb{P}_n - \mathbb{P})\{\varphi(O; \mathbb{P})\} \\ &\quad + \sqrt{n}(\mathbb{P}_n - \mathbb{P})\{\varphi(O; \widehat{\mathbb{P}}) - \varphi(O; \mathbb{P})\} + \sqrt{n}(\widehat{\Sigma} - \widetilde{\Sigma})(\widehat{\beta} - \beta) \\ &\quad + \sqrt{n}\widetilde{\Sigma}(\widehat{\beta} - \beta) + \sqrt{n}\mathbb{P}\{\varphi(O; \widehat{\mathbb{P}})\} \\ &=: C + T_1 + T_2,\end{aligned}\tag{28}$$

where $\widetilde{\Sigma} := \mathbb{P}\{(X - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\}$. By the central limit theorem, each entry of the first term C is $\mathcal{O}_{\mathbb{P}}(1)$. We next derive finite-sample deviation bounds for the other terms and show that they are $o_{\mathbb{P}}(1)$ under the extra rate conditions as assumed.

Part (1) Controlling the empirical process term T_1 . We begin by decomposing T_1 :

$$\begin{aligned}\varphi(O; \widehat{\mathbb{P}}) - \varphi(O; \mathbb{P}) + (X - \widehat{\mathbb{E}}[X | U])^{\otimes 2}(\widehat{\beta} - \beta) \\ &= (X - \widehat{\mathbb{E}}[X | U])(\widehat{\eta}(O) - \widehat{\beta}^\top(X - \widehat{\mathbb{E}}[X | U]))^\top - (X - \mathbb{E}[X | U])(\eta(O) - \beta^\top(X - \mathbb{E}[X | U]))^\top \\ &\quad + (X - \widehat{\mathbb{E}}[X | U])^{\otimes 2}(\widehat{\beta} - \beta) \\ &= [(X - \mathbb{E}[X | U])^{\otimes 2} - (X - \widehat{\mathbb{E}}[X | U])^{\otimes 2}]\beta + [(X - \widehat{\mathbb{E}}[X | U])\widehat{\eta}(O)^\top - (X - \mathbb{E}[X | U])\eta(O)^\top] \\ &=: S_1 + S_2.\end{aligned}$$

Note that each term above takes the form of $\widehat{ab} - ab = \widehat{a}(\widehat{b} - b) + (\widehat{a} - a)b$, which we will next use to derive the upper bound.

For the first term, we have

$$\begin{aligned}\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})S_1\|_{2,\infty} \\ &= \sqrt{n}\|(\mathbb{P}_n - \mathbb{P})[(X - \widehat{\mathbb{E}}[X | U])^{\otimes 2} - (X - \mathbb{E}[X | U])^{\otimes 2}]\beta\|_{2,\infty} \\ &= \sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{A_1\}\beta\|_{2,\infty},\end{aligned}$$

where

$$A_1 = (X - \widehat{\mathbb{E}}[X | U])^{\otimes 2} - (X - \mathbb{E}[X | U])^{\otimes 2}.$$

From Lemma E.5, we have

$$\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})S_1\|_{2,\infty} \leq \epsilon^{-\frac{1}{2}}\mathbb{E}[\|A_1\|_{\text{op}}^2\|\beta\|_{2,\infty}^2]^{\frac{1}{2}} \leq \epsilon^{-\frac{1}{2}}\mathbb{E}[\|A_1\|_{\text{op}}^2]^{\frac{1}{2}}\|\beta\|_{2,\infty},$$

with probability at least $1 - \epsilon$. Now, it remains to derive the upper bound of the expected squared operator norm:

$$\begin{aligned}\mathbb{E}[\|A_1\|_{\text{op}}^2]^{\frac{1}{2}} &\leq \mathbb{E}[\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_2^2(\|X - \mathbb{E}[X | U]\|_2 + \|X - \widehat{\mathbb{E}}[X | U]\|_2)^2]^{\frac{1}{2}} \\ &\leq \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}}(\|\mathbb{E}[X | U]\|_{L_{2(1+\delta^{-1})}} + \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta^{-1})}}).\end{aligned}$$

Therefore, we have

$$\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})S_1\|_{2,\infty} \leq 2M^2\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}}$$

with probability at least $1 - \epsilon$.

For the second term, similarly, we have

$$\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})S_2\|_{2,\infty}$$

$$\begin{aligned}
&\leq \epsilon^{-\frac{1}{2}} \max_{j \in [p]} \mathbb{E}[\|(X - \widehat{\mathbb{E}}[X | U])([\widehat{\eta}(O) - \eta(O)]^\top)_{\cdot j} + (\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])[\eta(O)^\top]_{\cdot j}\|^2]^{\frac{1}{2}} \\
&\leq \epsilon^{-\frac{1}{2}} (\|X - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta^{-1})}} \|(\widehat{\eta}(O) - \eta(O))^\top\|_{2,\infty}^2 \|_{L_{1+\delta}}^{\frac{1}{2}} + \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} \|\eta(O)^\top\|_{L_{2(1+\delta^{-1}),\infty}}) \\
&\leq \epsilon^{-\frac{1}{2}} M (\|\|\widehat{\eta}(O) - \eta(O)\|_\infty\|_{L_{2(1+\delta)}} + \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}})
\end{aligned}$$

with probability at least $1 - \epsilon$.

Combining the above results, with probability at least $1 - 2\epsilon$, we have

$$\begin{aligned}
\|T_1\|_{2,\infty} &\leq 2M^2 \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} \\
&\quad + M (\|\|\widehat{\eta}(O) - \eta(O)\|_\infty\|_{L_{2(1+\delta)}} + \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}}) \\
&\leq 2M(M \vee 1) \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} + M \|\|\widehat{\eta}(O) - \eta(O)\|_\infty\|_{L_{2(1+\delta)}}. \quad (29)
\end{aligned}$$

Part (2) Controlling the bias term T_2 . For the third term T_2 in (28), we have

$$\begin{aligned}
T_2 &= \sqrt{n} \widetilde{\Sigma}(\widehat{\beta} - \beta) + \sqrt{n} \mathbb{P}\{\varphi(O; \widehat{\mathbb{P}})\} \\
&= \sqrt{n} \mathbb{P}\{(X - \widehat{\mathbb{E}}[X | U])\widehat{\eta}(O)^\top\} - \sqrt{n} \widetilde{\Sigma} \beta \\
&= \sqrt{n} \mathbb{P}\{(X - \widehat{\mathbb{E}}[X | U])(g'(\widehat{\mathbb{E}}[Y | X, U]) \odot (Y - \widehat{\mathbb{E}}[Y | X, U]) + g(\widehat{\mathbb{E}}[Y | X, U]) - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U])^\top\} \\
&\quad - \sqrt{n} \widetilde{\Sigma} \Sigma^{-1} \Sigma \beta, \quad (30)
\end{aligned}$$

where the last equality is because of Equations (25) and (26). Denote the second-order remaining term by $Q = g(\mathbb{E}[Y | X, U]) - g(\widehat{\mathbb{E}}[Y | X, U]) - g'(\widehat{\mathbb{E}}[Y | X, U]) \odot (\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U])$. Then, we further have

$$\begin{aligned}
T_2 &= -\sqrt{n} \mathbb{P}\{(X - \widehat{\mathbb{E}}[X | U])(g(\mathbb{E}[Y | X, U]) + \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U] + Q)^\top\} \\
&\quad + \sqrt{n} \mathbb{P}\{(X - \mathbb{E}[X | U])(g(\mathbb{E}[Y | X, U]) - \mathbb{E}[g(\mathbb{E}[Y | X, U]) | U])^\top\} \\
&\quad + \sqrt{n} (I_d - \widetilde{\Sigma} \Sigma^{-1}) \Sigma \beta \\
&= \sqrt{n} \mathbb{P}\{(X - \widehat{\mathbb{E}}[X | U])Q^\top\} \\
&\quad + \sqrt{n} \mathbb{P}\{(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])(\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U])^\top\} \\
&\quad + \sqrt{n} (\Sigma - \widetilde{\Sigma}) \beta. \quad (31)
\end{aligned}$$

Because by the law of iterative expectation,

$$\mathbb{P}\{(X - \mathbb{E}[X | U])(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^\top\} = 0, \quad (32)$$

we have

$$\begin{aligned}
1 - \widetilde{\Sigma} \Sigma^{-1} &= 1 - \mathbb{P}\{(X - \mathbb{E}[X | U] + \mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\} \Sigma^{-1} \\
&= -\mathbb{P}\{(X - \mathbb{E}[X | U])(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^\top\} - \mathbb{P}\{(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])(X - \mathbb{E}[X | U])^\top\} \\
&\quad + \mathbb{P}\{(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\} \Sigma^{-1} \\
&= \mathbb{P}\{(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\} \Sigma^{-1}
\end{aligned}$$

and

$$\begin{aligned}
\|\Sigma - \widetilde{\Sigma}\|_{\text{op}} &= \|(I_d - \widetilde{\Sigma} \Sigma^{-1})\Sigma\|_{\text{op}} \quad (33) \\
&= \|\mathbb{P}\{(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\}\|_{\text{op}} \\
&\leq \|\mathbb{P}\{(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\}\|_{\text{op}} \quad (\text{Jensen's inequality})
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{P}\{\|(\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U])^{\otimes 2}\|_{\text{op}}\} \\
&= \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2
\end{aligned} \tag{34}$$

Combining Equations (27), (31) and (34) yields that

$$\begin{aligned}
\|T_2\|_{2,\infty} &\leq \sqrt{n}\|Q\|_{L_2,\infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2} \\
&\quad + \sqrt{n}\|\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U]\|_{L_2,\infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2} \\
&\quad + \sqrt{n}\|\Sigma - \widetilde{\Sigma}\|_{\text{op}}\|\beta\|_{2,\infty} \\
&\leq \sqrt{n}ML\|\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U]\|_{L_2,\infty}^2 \\
&\quad + \sqrt{n}\|\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U]\|_{L_2,\infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2} \\
&\quad + M\sqrt{n}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2.
\end{aligned} \tag{35}$$

Part (3) Combining the above results. Finally, from Equations (28), (29) and (35)

$$\widehat{\Sigma}\sqrt{n}(\widehat{\beta} - \beta) = \sqrt{n}(\mathbb{P}_n - \mathbb{P})\{\varphi(O; \mathbb{P})\} + \varsigma$$

for some $\varsigma \in \mathbb{R}^{d \times p}$ with

$$\begin{aligned}
\|\varsigma\|_{2,\infty} &\leq 2M(M \vee 1)\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} + \sqrt{n}M\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2 \\
&\quad + M\|\|\widehat{\eta}(O) - \eta(O)\|_{\infty}\|_{L_{2(1+\delta)}} \\
&\quad + \sqrt{n}ML\|\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U]\|_{L_2,\infty}^2 \\
&\quad + \sqrt{n}\|\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U]\|_{L_2,\infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}.
\end{aligned}$$

Note that

$$\begin{aligned}
\|\Sigma^{-1} - \widehat{\Sigma}^{-1}\|_{\text{op}} &= \|\widehat{\Sigma}^{-1}(\widehat{\Sigma} - \Sigma)\Sigma^{-1}\|_{\text{op}} \\
&\leq \|\widehat{\Sigma}^{-1}\|_{\text{op}}\|\widehat{\Sigma} - \Sigma\|_{\text{op}}\|\Sigma^{-1}\|_{\text{op}} \\
&\leq \sigma^2\|\widehat{\Sigma} - \Sigma\|_{\text{op}} \\
&\leq \sigma^2\|(\mathbb{P}_n - \mathbb{P})\{(X - \mathbb{E}[X | U])^{\otimes 2}\}\|_{\text{op}} + \sigma^2\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2,
\end{aligned}$$

where the first equality is from $\widehat{\Sigma}^{-1}(\widehat{\Sigma} - \Sigma)\Sigma^{-1} = \Sigma^{-1} - \widehat{\Sigma}^{-1}$, the second inequality is from the positivity assumption that $\|\widehat{\Sigma}^{-1}\|_{\text{op}} \leq \sigma$, $\|\Sigma^{-1}\|_{\text{op}} \leq \sigma$, and the last inequality is from (34). We further have

$$\sqrt{n}(\widehat{\beta} - \beta) = \sqrt{n}\Sigma^{-1}(\mathbb{P}_n - \mathbb{P})\{\varphi(O; \mathbb{P})\} + \xi,$$

with

$$\xi = \sqrt{n}(\widehat{\Sigma}^{-1} - \Sigma^{-1})(\mathbb{P}_n - \mathbb{P})\{\varphi(O; \mathbb{P})\} + \widehat{\Sigma}^{-1}\varsigma.$$

By multidimensional Chebyshev inequality and union bound, with probability at least $1 - 3\epsilon$,

$$\begin{aligned}
\|\xi\|_{2,\infty} &\leq \sigma^2(\|(\mathbb{P}_n - \mathbb{P})\{(X - \mathbb{E}[X | U])^{\otimes 2}\}\|_{\text{op}} + \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2) \\
&\quad \cdot \|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)^{-1}}}(\|\eta(O)\|_{L_{2(1+\delta)},\infty} + \|\beta\|_{2,\infty}) + \sigma\|\varsigma\|_{2,\infty} \\
&\leq 2\sigma^2M^2\|(\mathbb{P}_n - \mathbb{P})\{(X - \mathbb{E}[X | U])^{\otimes 2}\}\|_{\text{op}} \\
&\quad + 2M(M \vee 1)\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_{2(1+\delta)}} + M\|\|\widehat{\eta}(O) - \eta(O)\|_{\infty}\|_{L_{2(1+\delta)}}
\end{aligned}$$

$$\begin{aligned}
& + \sqrt{n}2(\sigma^2 \vee 1)M(M \vee 1)\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}^2 \\
& + \sqrt{n}ML\|\mathbb{E}[Y | X, U] - \widehat{\mathbb{E}}[Y | X, U]\|_{L_2, \infty}^2 \\
& + \sqrt{n}\|\mathbb{E}[g(\mathbb{E}[Y | X, U]) | U] - \widehat{\mathbb{E}}[g(\widehat{\mathbb{E}}[Y | X, U]) | U]\|_{L_2, \infty}\|\mathbb{E}[X | U] - \widehat{\mathbb{E}}[X | U]\|_{L_2}.
\end{aligned}$$

Under the extra rate conditions as assumed, we further have $\|\xi\|_{2, \infty} = o_{\mathbb{P}}(1)$. This completes the proof. \square

Lemma E.4 (Multivariate Lindeberg CLT for triangular array). *Let $m = m_n$ and $p = p_n$ be two sequences indexed by n . Consider the influence-function-based linear expansion for estimator $\widehat{\tau}_j$ of $\tau_j \in \mathbb{R}^d$:*

$$\sqrt{n}(\widehat{\tau}_j - \tau_j) = \sqrt{n}(\mathbb{P}_n - \mathbb{P})\{\varphi_{m_n j}\} + \varsigma_{m_n j}, \quad j = 1, \dots, p$$

where $\varphi_{m_n j}$ is the influence function that depends on m and the residual ς_j 's satisfy that $\|\varsigma_{m_n}\|_{2, \infty} = o_{\mathbb{P}}(1)$ as $n \rightarrow \infty$. Further assume that (i) there exists a constant $c > 0$, such that $\mathbb{V}(\varphi_{m_n j}(O_1)) \geq c$, and (ii) $\max_{k \in [n]} \mathbb{E}[\|\mathbb{V}\{\varphi_{m_n j}(O_k)\}^{-\frac{1}{2}}\varphi_{m_n j}(O_k)\|^{2+\frac{2}{\delta}}] \leq M$, then

$$\sqrt{n}\mathbb{V}\{\varphi_{m_n j}\}^{-1/2}(\widehat{\tau}_j - \tau_j) \xrightarrow{d} \mathcal{N}_d(0, I_d)$$

Proof of Lemma E.4. Note that $\varphi_{m_n j}$ is the centered influence function such that $\mathbb{E}[\varphi_{m_n j}(O)] = 0$. Let $X_{nk} = \mathbb{V}\{\varphi_{m_n j}(O_k)\}^{-\frac{1}{2}}\varphi_{m_n j}(O_k)$. From assumption (ii) that $\max_{k \in [n]} \mathbb{E}[\|X_{nk}\|^{2+\frac{2}{\delta}}] \leq M$, we have that, for any $\xi > 0$,

$$\frac{1}{n} \sum_{k=1}^n \mathbb{E} [\|X_{nk}\|^2 \mathbf{1}\{\|X_{nk}\| \geq \xi\sqrt{n}\}] \leq \frac{1}{n^{1+\frac{1}{\delta}}} \sum_{k=1}^n \mathbb{E} [\|X_{nk}\|^{2+\frac{2}{\delta}}] \leq M \frac{n}{n^{1+\frac{1}{\delta}}} \rightarrow 0.$$

This verifies Lindeberg's condition for a triangular array of random variables. From the multivariate Lindeberg's theorem (e.g., Billingsley (1995, Theorem 29.5)), it follows that

$$\sqrt{n}\mathbb{V}\{\varphi_{m_n j}\}^{-\frac{1}{2}}(\mathbb{P}_n - \mathbb{P})\{\varphi_{m_n j}\} \xrightarrow{d} \mathcal{N}(0, 1),$$

as $n \rightarrow \infty$. From assumption (i) that $\mathbb{V}(\varphi_{m_n j}) \geq c > 0$ and $\max_{j \in p_n} \|\varsigma_{m_n j}\| = o_{\mathbb{P}}(1)$, we further have

$$\max_{j \in p_n} \|\mathbb{V}\{\varphi_{m_n j}\}^{-\frac{1}{2}}\varsigma_{m_n j}\| = o_{\mathbb{P}}(1)$$

as $n \rightarrow \infty$. Consequently, the conclusion follows. \square

Lemma E.5 (Matrix Chebyshev inequality). *Let A denote a random matrix in $\mathbb{R}^{d \times r}$ and $\beta \in \mathbb{R}^{r \times p}$ such that $\mathbb{E}[A\beta] = 0_{d \times p}$. Then with probability at least $1 - \epsilon$, it holds that*

$$\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{A\beta\}\|_{2, \infty} \leq \epsilon^{-\frac{1}{2}}\mathbb{E}[\|A\|_{\text{op}}^2\|\beta\|_{2, \infty}^2]^{\frac{1}{2}},$$

and

$$\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{A\}\|_{\text{op}} \leq \epsilon^{-\frac{1}{2}}\mathbb{E}[\|A\|_{\text{op}}^2]^{\frac{1}{2}}.$$

Proof of Lemma E.5. By Chebyshev inequality, we have

$$\begin{aligned}
\mathbb{P}(\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{A\beta\}\|_{2, \infty} > t) & \leq \frac{n\mathbb{E}[\|(\mathbb{P}_n - \mathbb{P})\{A\beta\}\|_{2, \infty}^2]}{t^2} \\
& \leq \frac{n\mathbb{E}[\|A\beta - \mathbb{E}[A\beta]\|_{2, \infty}^2]}{nt^2} \\
& = \frac{\mathbb{E}[\|A\beta\|_{2, \infty}^2]}{t^2}.
\end{aligned}$$

Choosing $t = \mathbb{E}[\|A\beta\|_{2,\infty}^2]^{\frac{1}{2}}\epsilon^{-\frac{1}{2}}$ yields that, with probability at least $1 - \epsilon$,

$$\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{A\beta\}\|_{2,\infty} \leq \epsilon^{-\frac{1}{2}}\mathbb{E}[\|A\beta\|_{2,\infty}^2]^{\frac{1}{2}} \leq \epsilon^{-\frac{1}{2}}\mathbb{E}[\|A\|_{\text{op}}^2\|\beta\|_{2,\infty}^2]^{\frac{1}{2}},$$

which finishes the proof of the first statement.

Similarly, considering all unit vectors in the unit sphere \mathbb{S}^{r-1} (i.e., the set of vector $v \in \mathbb{R}^r$ such that $\|v\|_2 = 1$), it holds that

$$\mathbb{P}(\sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{A\}\|_{\text{op}} > t) = \mathbb{P}\left(\sup_{v \in \mathbb{S}^{r-1}} \sqrt{n}\|(\mathbb{P}_n - \mathbb{P})\{Av\}\|_2 > t\right) \leq \frac{\mathbb{E}[\|A\|_{\text{op}}^2]}{t^2}.$$

The second conclusion follows by choosing $t = (\mathbb{E}[\|A\|_{\text{op}}^2]/\epsilon)^{\frac{1}{2}}$. \square

F Extra experimental results

F.1 Simulation

As in Theorem E.1, the nuisance functions need to be estimated fast enough such that valid inference can be guaranteed. We first examine the convergence rate of the nuisance estimations. The L_2 consistency of random forests has been examined in various studies; see, for example, (Biau, 2012; Scornet et al., 2015). The rate of convergence is closely related to the minimax rate of $\mathcal{O}_{\mathbb{P}}(n^{-2/(q+2)})$ for nonparametric estimation involving q features. In a simplified setting, Biau (2012) demonstrated that this rate can be improved to $\mathcal{O}_{\mathbb{P}}(n^{-0.75/(s+0.75)})$, where s represents the intrinsic dimension, which can be substantially smaller than the total feature dimension q . By numerical examination of the convergence rate for nuisance estimation, our findings indicate a L_2 convergence rate of approximately $n^{-1/4}$ for both nuisance functions on the simulated data, as illustrated in Figure F1. This supports the appropriate use of doubly robust estimators in our experiments.




Figure F1: Estimation error of the nuisance regression function on simulated data using random forests. The axes are shown in the logarithm scale and the slope represents the estimated rate of convergence. The data-generating process is given in Section 4, and we use the true latent embedding U so that the ground truth regression function is computable. The errors are computed based on 1000 test observations without irreducible additive noises.

F.2 Real data

Extended background In a recent single-cell CRISPR perturbation study, Lalli et al. (2020) investigated the molecular mechanisms of genes associated with neurodevelopmental disorders, particularly Autism Spectrum Disorder (ASD). Using a modified CRISPR-Cas9 system, they performed gene suppression experiments on 13 ASD-linked genes in the Lund Human Mesencephalic (LUHMES) neural progenitor cells. The experiment comprised 14 groups: 13 treatment groups with individual gene knockdowns and one control group. Single-cell RNA sequencing was employed to assess gene expression changes resulting from each knockdown. The authors estimated a pseudotime trajectory, which approximates the progression of neuronal differentiation. The analysis of Lalli et al. (2020) suggests that some perturbations cause changes in pseudotime (slow or speed development); see Figure F2. A scientific question of interest not answered by Lalli et al. (2020) is whether some perturbation explains anything beyond the changes in expression levels caused by cell development.

In single-cell CRISPR perturbation experiments, confounding factors can significantly impact the interpretation of results. Unlike controlled experiments, these studies often resemble observational data, where confounding variables such as cell size, cell cycle stage, or microenvironment heterogeneity may influence gene expression patterns. These confounders can mask or mimic the effects of the intended genetic perturbations, potentially leading to erroneous conclusions about gene function or regulatory networks. Addressing these confounding issues is crucial for the accurate interpretation of CRISPR perturbation data and for distinguishing true biological effects from technical artifacts.

To adjust for possible confounding effects, we may take advantage of the multiple negative control genes. Even though tens of thousands of genes are measured, one typically restricts the differential expression analysis to the top thousands of highly variable genes. For the remaining genes with low variations, it is believed that there will not be sufficient power to differentiate the response from the null distribution. But even with low power, it is likely that, in total, one can detect the impact of confounding. For this reason, we use such genes as “pseudo-negative control”; even if this choice is incorrect, we still target meaningful statistical estimands, provided that the estimated embedding captures the common variability of all cells under control. Alternatively, we can also use housekeeping genes as negative control outcomes. The main goal here is to demonstrate a practical procedure for post-integrated inference and show that our asymptotic results are reasonably accurate in real data.

Data. After filtering out low-quality cells and genes that expressed in less than 10 cells, we retained 8320 cells and 13086 genes under 14 perturbation conditions (including control) from Lalli et al. (2020). Following the routine selection procedure of highly variable genes in genomics (Hao et al., 2021), we select 4163 genes whose standardized variance is larger than 1, and the last 4000 genes with the lowest standardized variances are treated as negative control outcomes. The covariates we measured include the logarithm of library sizes, cell cycle scores ('S.Score' and 'G2M.Score'), batches (3 categories), and pseudotime states (normalized to range from 0 to 1). After one-hot encoding of the categorical features, we have 19 covariates (including 13 perturbation indicators), and 4163 genes for model fitting. For each highly variable gene, we aim to test whether its gene expressions vary along the pseudotime state under perturbation conditions.




Figure F2: Expression levels of marker genes in different estimated pseudotime states. Genes *MAP2* and *DCX* are neuronal markers (expressed in more differentiated cells) while genes *TP53* and *CDK4* are progenitor markers (expressed in less differentiated cells).




Figure F3: Histogram of test statistics for main effects of pseudotime states on the expressions of 4163 genes. Many genes are significant because the expression levels are expected to change during neural differentiation.




Figure F4: Histogram of test statistics on 4163 genes for 12 different perturbation conditions. Different rows represent the results of different methods: GLM: Score tests by generalized linear models with Negative Binomial likelihood and log link function. The covariance matrix is estimated using the HC3-type robust estimator. PII: The proposed post-integrated inference with 50 principal components as the estimated embeddings.




Figure F5: Gene expressions of significant genes in the control group and the *PTEN* knockdown group. Four genes with positive estimated effect sizes are selected with a p-value threshold of 0.01 for both pseudotime states and *PTEN* knockdown for three PII methods in Figure 8(b) and a median expression level larger than zero.

When restricted to a small subset of significant genes discovered by PII, their expression levels are visualized as a function of pseudotime states and perturbation conditions in Figure F5. We observe an increasing trend of the expression and the overexpression in the perturbed group at the very late stage of pseudotime. The significance suggests that these genes could be affected by not only the cell development but also the *PTEN* repression. *NEFM* is involved in neurite outgrowth and axon caliber (Cheung et al., 2023), *TUBB2B* and *TUBA1A* encode critical structural subunits of microtubules that are enriched during brain development (Jaglin et al., 2009), *HN1* is related to cancer and senescence (Jia et al., 2019). Given the role of *PTEN* on neural differentiation and related processes, these genes could be affected. Further research would be needed to establish any direct links between *PTEN* repression and the expression or function of these specific genes during neural differentiation.