

1 Distrbutions

Bernoulli:  $\begin{cases} 1-p & \text{if } x = 0 \\ p & \text{if } x = 1 \end{cases}$

$$\mu = p, Var = pq$$

$$MGF = q + pe^t$$

Binomial:  $\binom{n}{k} p^k q^{n-k}$

$$\mu = np, Var = npq$$
$$MGF = (q + pe^t)^n$$

Geometric:  $p(1-p)^{k-1}$

$$\mu = \frac{1}{p}, Var = \frac{1-p}{p^2}$$
$$MGF = \frac{pe^t}{1-(1-p)e^t}$$

Poisson:  $\frac{e^{-\lambda} \lambda^k}{k!}$

$$\mu = \lambda, Var = \lambda$$
$$MGF = e^{\lambda(e^t-1)}$$

Normal:  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(k-\mu)^2}{2\sigma^2}}$

$$\mu = \mu, Var = \sigma^2$$
$$MGF = e^{\mu t + \frac{1}{2}\sigma^2 t^2}$$

Exponential:  $\lambda e^{-\lambda k}$

$$\mu = \frac{1}{\lambda}, Var = \frac{1}{\lambda^2}$$

$$MGF = \frac{\lambda}{\lambda - t}, \quad t < \lambda$$

Uniform:  $\frac{1}{b-a}, \quad a \leq k \leq b$

$$\mu = \frac{a+b}{2}, Var = \frac{(b-a)^2}{12}$$
$$MGF = \frac{e^{bt} - e^{at}}{t(b-a)}$$

Beta:  $\frac{k^{\alpha-1}(1-k)^{\beta-1}}{B(\alpha,\beta)}, \quad 0 \leq k \leq 1$

$$\mu = \frac{\alpha}{\alpha + \beta}, Var = \frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)}$$
$$MGF = {}_2F_1(\alpha, \alpha + \beta; \alpha + \beta + 1; t)$$

$$\beta(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$
$$\Gamma(\alpha) = (\alpha - 1)!, \quad \alpha \in \mathbb{N}$$

Gamma:  $\frac{\beta^\alpha k^{\alpha-1} e^{-\beta k}}{\Gamma(\alpha)}$

$$\mu = \frac{\alpha}{\beta}, Var = \frac{\alpha}{\beta^2}$$
$$MGF = \left(\frac{\beta}{\beta - t}\right)^\alpha, \quad t < \beta$$

2 Formulas

2.1 Probability Formulas

$$A \perp B \Leftrightarrow \mathbb{P}(A\|B) = \mathbb{P}(A) \& \mathbb{P}(B\|A) = \mathbb{P}(B) \Leftrightarrow \mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Union Bound:  $\mathbb{P}(\cup_{i=1}^n A_i) \leq \sum_{i=1}^n \mathbb{P}(A_i)$

Bayes' Rule:  $\mathbb{P}(A\|B) = \frac{\mathbb{P}(B\|A) \cdot \mathbb{P}(A)}{\mathbb{P}(B)}$

Law of Total Probability:  $\mathbb{P}(A) = \sum_{i=1}^n \mathbb{P}(A\|B_i) \cdot \mathbb{P}(B_i)$

Chain Rule:  $\mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n) = \mathbb{P}(A_1) \cdot \mathbb{P}(A_2\|A_1) \cdot \dots \cdot \mathbb{P}(A_n\|A_1 \cap \dots \cap A_{n-1})$

Conditional Independence:  $A \perp B\|C \Leftrightarrow \mathbb{P}(A\|B \cap C) = \mathbb{P}(A\|C)$

2.2 Expected Value and Variance

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$
$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \mathbb{E}((X - \mathbb{E}(X))^2)$$
$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) \text{ if } X \perp Y$$
$$\text{Var}(aX) = a^2 \text{Var}(X)$$

2.3 Moments and MGF

$$\mu_k = \mathbb{E}(X^k)$$
$$\bar{\mu}_k = \mathbb{E}((X - \mathbb{E}(X))^k)$$
$$M_X(t) = \mathbb{E}(e^{tX})$$
$$M_{X+Y}(t) = M_X(t) \cdot M_Y(t) \text{ if } X \perp Y$$
$$\mu_k = M_X^{(k)}(0)$$
$$\mathbb{E}[\mathbb{E}[X\|Y]] = \mathbb{E}[X]$$

2.4 function of random variables

$$\mathbb{E}[g(X)] = \sum_x g(x) \cdot \mathbb{P}(X = x)$$
$$f_Y(y) = f_X(g^{-1}(y)) \cdot \left| \frac{d}{dy} g^{-1}(y) \right|$$

2.5 Common Distributions

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y)$$
$$f_{X,Y}(x, y) = \frac{\partial^2}{\partial x \partial y} F_{X,Y}(x, y)$$
$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$$
$$X \perp Y \Leftrightarrow f_{X,Y}(x, y) = f_X(x) \cdot f_Y(y)$$

Jointly Gaussian:

$$f_{X,Y}(x, y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left(-\frac{1}{2(1-\rho^2)}\left(\left(\frac{x-\mu_X}{\sigma_X}\right)^2 - 2\rho\left(\frac{x-\mu_X}{\sigma_X}\right)\left(\frac{y-\mu_Y}{\sigma_Y}\right) + \left(\frac{y-\mu_Y}{\sigma_Y}\right)^2\right)\right)$$

$$f_{X\|Y}(x\|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$
$$\mathbb{E}[X\|Y = y] = \int_{-\infty}^{\infty} x \cdot f_{X\|Y}(x\|y) dx$$

2.6 Normal Distribution

if  $X \sim \mathcal{N}(\mu_X, \sigma_X^2)$  and  $Y \sim \mathcal{N}(\mu_Y, \sigma_Y^2)$ , then  $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \sigma_X^2 + \sigma_Y^2)$

if X and Y are jointly Gaussian, then  $X\|Y = y \sim \mathcal{N}(\mu_X + \rho \frac{\sigma_X}{\sigma_Y}(y - \mu_Y), \sigma_X^2(1 - \rho^2))$

2.7 Covaariance and Correlation

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}(X))(Y - \mathbb{E}(Y))] = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)$$
$$\text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X\sigma_Y}$$

if two random variables are linearly independent then  $\text{Cov}(X, Y) = 0$

$$\text{Cov}(aX + b, cY + d) = ac\text{Cov}(X, Y) \text{ (independent of a,b,c,d)}$$
$$\mu_{i,j} = \mathbb{E}(X^i Y^j)$$
$$\bar{\mu}_{i,j} = \mathbb{E}((X - \mathbb{E}(X))^i (Y - \mathbb{E}(Y))^j)$$

2.8 CLT and LLN

Central Limit Theorem:  $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \rightarrow \mathcal{N}(0, 1)$  as  $n \rightarrow \infty$

Law of Large Numbers:  $\bar{X} \rightarrow \mu$  as  $n \rightarrow \infty$

2.9 Functions of multiple Random Variables

$$J = \begin{bmatrix} \frac{\partial g_1}{\partial x_1} & \dots & \frac{\partial g_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial g_m}{\partial x_1} & \dots & \frac{\partial g_m}{\partial x_n} \end{bmatrix}$$
$$f_{Y_1, \dots, Y_m}(y_1, \dots, y_m) = f_{X_1, \dots, X_n}(g_1^{-1}(y_1, \dots, y_m), \dots, g_m^{-1}(y_1, \dots, y_m)) \cdot \det(J)$$

3 Probability Inequalities

3.1 Markov's Inequality

For any non-negative random variable X and any  $a > 0$ :

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$$

3.2 Chebyshev's Inequality

For any random variable X and any  $a > 0$ :

$$\mathbb{P}(|X - \mathbb{E}[X]| \geq a) \leq \frac{\text{Var}(X)}{a^2}$$

3.3 Chernoff Bound

For any random variable X and any  $t > 0$ :

$$\mathbb{P}(X \geq (1 + \delta)\mathbb{E}[X]) \leq \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}}\right)^{\mathbb{E}[X]}$$

$$\mathbb{P}(X \leq (1 - \delta)\mathbb{E}[X]) \leq \left(\frac{e^{-\delta}}{(1 - \delta)^{1-\delta}}\right)^{\mathbb{E}[X]}$$

3.4 Jensen's Inequality

For any random variable X and any convex function f:

$$f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)]$$

3.5 Cauchy-Schwarz Inequality

given two vectors x and y:

$$\left(\sum_{i=1}^n x_i y_i\right)^2 \leq \left(\sum_{i=1}^n x_i^2\right) \left(\sum_{i=1}^n y_i^2\right)$$

For any random variables X and Y:

$$\mathbb{E}[XY]^2 \leq \mathbb{E}[X^2] \cdot \mathbb{E}[Y^2]$$

and the equality holds if and only if  $X = aY$  for some  $a \in \mathbb{R}$ .

## 4 Sample Statistics

### 4.1 Properties:

Bias:  $\mathbb{E}[\hat{\theta}] - \theta$   
Variance:  $\text{Var}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \mathbb{E}[\hat{\theta}])^2]$   
Mean Squared Error:  $\text{MSE}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \theta)^2] = \text{Var}(\hat{\theta}) + \text{Bias}^2$   
Consistency:  $\lim_{n \rightarrow \infty} \hat{\theta} = \theta$

### 4.2 Estimators:

Sample Mean:  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$   
Sample Variance:  $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$

### 4.3 Maximum Likelihood Estimation:

$\hat{\theta}_{\text{MLE}} = \arg \max_{\theta} \prod_{i=1}^n f(X_i | \theta) = \arg \max_{\theta} \sum_{i=1}^n \log f(X_i | \theta)$   
 $\Rightarrow \frac{\partial \ell(\theta)}{\partial \theta}$

### 4.4 MMSE:

linear MMSE:  $h(a, b) = \mathbb{E}[(X - aY - b)^2]$   
objective: minimize  $h(a, b)$   
solution:  $a = \frac{\text{Cov}(X, Y)}{\text{Var}(Y)} = \rho \frac{\sigma_X}{\sigma_Y}$ ,  $b = \mathbb{E}[X] - a\mathbb{E}[Y]$   
multiple MMSE:

$$\begin{bmatrix} \sigma_{X_1}^2 & \sigma_{X_1 X_2} \\ \sigma_{X_1 X_2} & \sigma_{X_2}^2 \end{bmatrix} \begin{bmatrix} a_1^* \\ a_2^* \end{bmatrix} = \begin{bmatrix} \text{Cov}(X_1, Y) \\ \text{Cov}(X_2, Y) \end{bmatrix}$$

Covariance Matrix

$$\hat{Y}(X_1, X_2) = a_1^*(X_1 - \mathbb{E}[X_1]) + a_2^*(X_2 - \mathbb{E}[X_2]) + \mathbb{E}[Y]$$

### 4.5 Hypothesis Testing:

Null Hypothesis:  $H_0 : \theta = \theta_0$   
Alternative Hypothesis:  $H_1 : \theta \neq \theta_0$   
Type I Error: Reject  $H_0$  when it is true  
Type II Error: Accept  $H_0$  when it is false  
P\_value: Probability of observing the data or more extreme data given  $H_0$  is true

### 4.6 Z-Test:

$Z = \frac{\bar{X} - \mu}{\sigma / \sqrt{n}}$   
 $\Rightarrow$  p-value =  $2 \cdot (1 - \Phi(|Z|))$

Confidence Interval for Mean:  $\bar{X} \pm Z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$

### 4.7 T-test

: Single Sample T-test:  
 $T = \frac{\bar{X} - \mu}{S / \sqrt{n}}$   
 $\Rightarrow$  p-value =  $2 \cdot (1 - t_{n-1}(|T|))$   
2-Sample T-test:  
 $T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}}$   
 $\Rightarrow$  p-value =  $2 \cdot (1 - t_{n_1+n_2-2}(|T|))$   
2-Sample T-test (Unequal Variance):  
 $T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}}$   
 $\Rightarrow$  p-value =  $2 \cdot (1 - t_{\nu}(|T|))$

where  $\nu = \frac{\left(\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}\right)^2}{\left(\frac{S_1^2}{n_1}\right)^2 + \left(\frac{S_2^2}{n_2}\right)^2}$

### 4.8 Chi-Square Test:

Goodness of Fit Test:  
 $\chi^2 = \sum_{i=1}^k \frac{(O_i - E_i)^2}{E_i}$   
 $\Rightarrow$  p-value =  $1 - \chi_{k-1}^2(\chi^2)$

### 4.9 Fisher's Exact Test:

$H_0$  : The two samples are from the same distribution  
 $H_1$  : The two samples are from different distributions  
 $P = \frac{\binom{n_1}{x} \binom{n_2}{y}}{\binom{n_1+n_2}{x+y}}$   
p\_value: getting the observed data or more extreme data

### 4.10 Permutation Test:

$H_0$  : The two samples are from the same distribution  
 $H_1$  : The two samples are from different distributions  
consider an arbitrary test statistic  $T$   
permute the labels of the samples to get the distribution of  $T$  under  $H_0$   
then calculate the p-value as the probability of observing the data or more extreme data under  $H_0$

### 4.11 Bayesian Inference:

MAP:  
 $\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} f(\theta | X) = \arg \max_{\theta} f(X | \theta) \cdot f(\theta)$

### 4.12 Linear Regression:

Simple Linear Regression:  $Y = \beta_0 + \beta_1 X + \epsilon$   
where  $\epsilon \sim \mathcal{N}(0, \sigma^2)$   
 $\beta_1$  and  $\beta_0$  can be estimated using the Maximum Likelihood Estimation

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2} = \frac{\text{Cov}(X, Y)}{\text{Var}(X)}$$
$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X}$$
$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{\beta}_0 - \hat{\beta}_1 X_i)^2$$