

Linear Regression / LDA

Linear Methods for Classification

2019.1.30 임소현

Contents

Introduction

Linear Regression of an Indicator Matrix

Linear Discriminant Analysis

Reference

Classification ?

: 입력 벡터 x 로부터 이에 대응되는 타겟 클래스 K 에 대해 어떤 하나의 클래스에 속하도록 선정하는 작업

- 타겟 t 에 대해 K 의 크기를 가지는 이진 벡터로 정의

ex > $K=5$, 클래스가 2에 속하는 경우

$$t = (0, 1, 0, 0, 0)^T$$

4.1 Introduction




First

Second

Third

Fourth



- 클래스는 상호 배타적인 관계
- 이렇게 나누어진 지역을 *decision region* 라고 부른다
- 이를 나누는 경계면을 *decision boundaries* 라고 부른다

4.1 Introduction

• • •

First

Second

Third

Fourth

Model

- discriminant functions $\delta_k(x)$ for each class
- Posterior probabilities $\Pr(G = k|X = x)$

$$\Pr(G = 1|X = x) = \frac{\exp(\beta_0 + \beta^T x)}{1 + \exp(\beta_0 + \beta^T x)},$$

$$\Pr(G = 2|X = x) = \frac{1}{1 + \exp(\beta_0 + \beta^T x)}.$$

logit transformation



$$\log \frac{\Pr(G = 1|X = x)}{\Pr(G = 2|X = x)} = \beta_0 + \beta^T x$$

4.2 Linear Regression of an Indicator Matrix

First

Second

Third

Fourth

• • •

- there will be K such indicators Y_k , $k = 1, \dots, K$, with $Y_k = 1$ if $G = k$ else 0
- $Y = (Y_1, \dots, Y_K)$, *indicator response matrix* ($N \times K$)
- Y is a matrix of 0's and 1's, with each row having a single 1
- Fit a linear regression model

$$\hat{\mathbf{Y}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$$

- Coefficient matrix

$$\hat{\mathbf{B}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$$

4.2 Linear Regression of an Indicator Matrix

First

Second

Third

Fourth

• • •

- A new observation with input x is classified as follows:
 - compute the fitted output $\hat{f}(x)^T = (1, x^T)\hat{\mathbf{B}}$, a K vector;
 - identify the largest component and classify accordingly: $\hat{G}(x) = \operatorname{argmax}_{k \in \mathcal{G}} \hat{f}_k(x)$
- The response vector y_i (i th row of \mathbf{Y}) for observation i has the value $y_i = t_k$ if $g_i = k$
- Fit the linear model by least squares $\min_{\mathbf{B}} \sum_{i=1}^N \|y_i - [(1, x_i^T)\mathbf{B}]^T\|^2$
- The criterion is a sum-of-squared Euclidean distances of the fitted vectors from their targets

$$\hat{G}(x) = \operatorname{argmin}_k \|\hat{f}(x) - t_k\|^2$$

4.2 Linear Regression of an Indicator Matrix

First


Second

Third

Fourth

• • •

- There is a serious problem with the regression approach when the number of classes $K \geq 3$
- Figure illustrates an extreme situation when $K = 3$.
- The three classes are perfectly separated by linear decision boundaries, yet linear regression misses the middle class completely.



4.3 Linear Discriminant Analysis

First

Second

Third

Fourth

• • •

- Suppose $f_k(x)$ is the class conditional density of X in class $G=k$

multivariate Gaussian



$$f_k(x) = \frac{1}{(2\pi)^{p/2} |\Sigma_k|^{1/2}} e^{-\frac{1}{2} (x - \mu_k)^T \Sigma_k^{-1} (x - \mu_k)}$$

- Let π_k be the prior probability of class k , with $\sum_{k=1}^K \pi_k = 1$
- By Bayes theorem,

$$\Pr(G = k | X = x) = \frac{f_k(x) \pi_k}{\sum_{\ell=1}^K f_{\ell}(x) \pi_{\ell}}$$

4.3 Linear Discriminant Analysis

First

Second

Third

Fourth

• • •

LDA

- assume that the classes have a common covariance matrix
- Comparing two classes k and ℓ , log-ratio

$$\begin{aligned}\log \frac{\Pr(G = k|X = x)}{\Pr(G = \ell|X = x)} &= \log \frac{f_k(x)}{f_\ell(x)} + \log \frac{\pi_k}{\pi_\ell} \\ &= \log \frac{\pi_k}{\pi_\ell} - \frac{1}{2}(\mu_k + \mu_\ell)^T \Sigma^{-1}(\mu_k - \mu_\ell) \\ &\quad + x^T \Sigma^{-1}(\mu_k - \mu_\ell),\end{aligned}$$

4.3 Linear Discriminant Analysis

First

Second

Third

Fourth

• • •

LDA

- *linear discriminant functions*

$$\delta_k(x) = x^T \Sigma^{-1} \mu_k - \frac{1}{2} \mu_k^T \Sigma^{-1} \mu_k + \log \pi_k$$

- We don't know the parameters of the Gaussian distributions, so we need to estimate them
 - $\hat{\pi}_k = N_k/N$, where N_k is the number of class- k observations;
 - $\hat{\mu}_k = \sum_{g_i=k} x_i / N_k$;
 - $\hat{\Sigma} = \sum_{k=1}^K \sum_{g_i=k} (x_i - \hat{\mu}_k)(x_i - \hat{\mu}_k)^T / (N - K)$.

4.3 Linear Discriminant Analysis



• • •

First

Second

Third

Fourth



- shows three Gaussian distributions, with the same covariance and different means
- included are the contours of constant density enclosing 95% of the probability in each case
- the Bayes decision boundaries separating all three classes are the thicker solid lines

4.3 Linear Discriminant Analysis

First

Second

Third

Fourth

• • •

QDA (*quadratic discriminant functions*)

- if the Σ_k are not assumed to be equal
- *linear discriminant functions*

$$\delta_k(x) = -\frac{1}{2} \log |\Sigma_k| - \frac{1}{2} (x - \mu_k)^T \Sigma_k^{-1} (x - \mu_k) + \log \pi_k$$

- The decision boundary between each pair of classes k and ℓ is described by a quadratic equation $\{x : \delta_k(x) = \delta_\ell(x)\}$

4.3 Linear Discriminant Analysis

First

Second

Third

Fourth

...

RDA (*Regularized Discriminant Analysis*)

- to shrink the separate covariances of QDA toward a common covariance as in LDA
- very similar in flavor to ridge regression
- $\alpha \in [0, 1]$ allows a continuum of models between LDA and QDA

$$\hat{\Sigma}_k(\alpha) = \alpha \hat{\Sigma}_k + (1 - \alpha) \hat{\Sigma} \quad (4.13)$$

- Similar modifications to be shrunk toward the scalar covariance

$$\hat{\Sigma}(\gamma) = \gamma \hat{\Sigma} + (1 - \gamma) \hat{\sigma}^2 \mathbf{I}$$

- Replacing $\hat{\Sigma}$ in (4.13) by $\hat{\Sigma}(\gamma)$ leads to a more general family

Reference



First

Second

Third

Fourth

- <http://norman3.github.io/prml/docs/chapter04/1>
- <https://m.blog.naver.com/PostView.nhn?blogId=sw4r&logNo=221033110991&proxyReferer=https%3A%2F%2Fwww.google.com%2F>

Thank you :-)