






50–30 kya · s=0.01
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
28/100 with ≥1 significant test
48/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



50–30 kya · s=0.01
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
15/100 with ≥1 significant test
30/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



50–30 kya · s=0.01
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
6/100 with ≥1 significant test
6/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.01
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
65/100 with ≥1 significant test
73/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.01
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
22/100 with ≥1 significant test
42/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



10 kya–present · s=0.01
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
5/100 with ≥1 significant test
5/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.01
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
99/100 with ≥1 significant test
100/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.01
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
88/100 with ≥1 significant test
95/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.

50 kya–present · s=0.01

Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb




82/100 with ≥1 significant test


82/100 with ≥1 significant test


Order: first significant time, then longer spans · based on p < 0.05 in both panels

Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101

Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
82/100 with ≥1 significant test
92/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
48/100 with ≥1 significant test
74/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
34/100 with ≥1 significant test
35/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.