



50–30 kya · s=0.01
True TMRCA · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
69/100 with ≥1 significant test
97/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50–30 kya · s=0.01
Posterior median · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
36/100 with ≥1 significant test
68/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.




19/100 with ≥1 significant test


18/100 with ≥1 significant test


Order: first significant time, then longer spans · based on p < 0.05 in both panels


Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101


Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.01
True TMRCA · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
86/100 with ≥1 significant test
97/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.01
Posterior median · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
58/100 with ≥1 significant test
78/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



10 kya–present · s=0.01
Posterior-weighted mass · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
26/100 with ≥1 significant test
23/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.01
True TMRCA · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
99/100 with ≥1 significant test
100/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.01
Posterior median · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
99/100 with ≥1 significant test
100/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.

50 kya–present · s=0.01

Posterior-weighted mass · pointwise p < 0.05 · selected site: 500 kb




96/100 with ≥1 significant test


96/100 with ≥1 significant test


Order: first significant time, then longer spans · based on p < 0.05 in both panels

Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101

Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
True TMRCA · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
94/100 with ≥1 significant test
100/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
Posterior median · pointwise p < 0.05 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
81/100 with ≥1 significant test
91/100 with ≥1 significant test
Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



65/100 with ≥1 significant test

65/100 with ≥1 significant test

Order: first significant time, then longer spans · based on p < 0.05 in both panels

Same 100 neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101

Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.