






50–30 kya · s=0.01
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
25/100 with ≥1 significant test
49/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



50–30 kya · s=0.01
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
9/100 with ≥1 significant test
31/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



50–30 kya · s=0.01
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
6/100 with ≥1 significant test
6/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



10 kya–present · s=0.01
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
62/100 with ≥1 significant test
71/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



10 kya–present · s=0.01
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
21/100 with ≥1 significant test
35/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



10 kya–present · s=0.01
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
5/100 with ≥1 significant test
4/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



50 kya–present · s=0.01
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
99/100 with ≥1 significant test
99/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.01
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
79/100 with ≥1 significant test
94/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.01
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
79/100 with ≥1 significant test
84/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
81/100 with ≥1 significant test
88/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
35/100 with ≥1 significant test
65/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


50 kya–present · s=0.005
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
31/100 with ≥1 significant test
34/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.