

Dots: tested cumulative cutoffs. Bars: significant 1-kyr bins (1–2 through 49–50 kya).

p = (1 + number of neutrals ≥ selected) / 101; p < 0.05 requires at most 4 neutrals.

Pointwise comparisons; no correction across times or loci. These are not inferred selection durations.


Pointwise p < 0.05: posterior median
Selected site: 500 kb · 100 neutrals · same 1,000 pairs for all methods
Cumulative cutoff
1-kyr interval
Selection active
50–30 kya
s=0.01
10 kya–present
s=0.01
50 kya–present
s=0.01
50 kya–present
s=0.005
0
5
10
15
20
25
30
35
40
45
50
Time before present (kya)
Dots: tested cumulative cutoffs. Bars: significant 1-kyr bins (1–2 through 49–50 kya).
p = (1 + number of neutrals ≥ selected) / 101; p < 0.05 requires at most 4 neutrals.
Pointwise comparisons; no correction across times or loci. These are not inferred selection durations.



Dots: tested cumulative cutoffs. Bars: significant 1-kyr bins (1–2 through 49–50 kya).

p = (1 + number of neutrals ≥ selected) / 101; p < 0.05 requires at most 4 neutrals.

Pointwise comparisons; no correction across times or loci. These are not inferred selection durations.