





10 kya–present · s=0.05
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
100/100 with ≥1 significant test
100/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.05
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
100/100 with ≥1 significant test
100/100 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.



10 kya–present · s=0.05
Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (sorted)
1
25
50
75
100
0
10
20
30
40
50
Time before present (kya)
Time before present (kya)
98/100 with ≥1 significant test
99/100 with ≥1 significant test
Order: first significant time, then longer spans · based on p < 0.05 in both panels
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.1
True TMRCA · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
2
3
0
10
20
30
40
50
Time before present (kya)
3/3 with ≥1 significant test
3/3 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.


10 kya–present · s=0.1
Posterior median · pointwise p < 0.01 · selected site: 500 kb
Significant cutoff
Significant bin
Selection active
Cumulative cutoff
1-kyr interval
Selected simulation (proposal order)
1
2
3
0
10
20
30
40
50
Time before present (kya)
3/3 with ≥1 significant test
3/3 with ≥1 significant test
Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101
Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.

10 kya–present · s=0.1

Posterior-weighted mass · pointwise p < 0.01 · selected site: 500 kb



3/3 with ≥1 significant test

3/3 with ≥1 significant test

Order: first significant time, then longer spans · based on p < 0.05 in both panels

Same 100 segregating neutrals · 1,000 pairs · p = (1 + neutral exceedances) / 101

Blank rows have no pointwise call. No correction across times/loci; bars do not estimate selection duration.