





Whole-population AF; conditioned on segregation today. Boxes: IQR; whiskers: 10th–90th percentiles.



Lines: means. Bands: 10th–90th percentiles across regions, not confidence intervals.
Right-hand y-scales vary. Mass is not renormalized to 1–50 kya; focal coordinate is 500 kb.


Introgression, s=0.01 · selected-minus-neutral coalescence mass
Dashed: selection boundaries (10 kya–present). Dotted: ±100 kb. n=100 selected, 50 neutral.
True TMRCAs · 19,900 pairs
Posterior median · 1,000 pairs
Posterior mass · 1,000 pairs
Distance from selected coordinate (kb)
500
0
−500
1
10
20
30
40
50
Coalescence time (kya)
−0.80% −0.60% −0.40% −0.20% 0.00% 0.20% 0.40% 0.60% 0.80%
Excess mass per 1-kyr bin


Introgression, s=0.01 · temporal calls across 100 survivors
Significant support
True selection: 10 kya–present
True TMRCAs
19,900 pairs
1 overlap; 13 mistimed
Posterior median
1,000 pairs
0 overlap; 0 mistimed
Posterior mass
1,000 pairs
4 overlap; 0 mistimed
Replicates ordered by present-day AF (low → high)
100
75
50
25
1
1
10
30
50
Coalescence time (kya)
Support is the union over positions within ±100 kb; threshold is calibrated over the entire 1-Mb region.
“Mistimed” means no near-site temporal overlap. Rows are identical across panels; every survivor is shown.