




100 neutral allele-frequency trajectories
EAS demography | 1% Neanderthal introduction at 50 kya | s = 0
Original focal allele segregating in the present-day population
Focal allele frequency
100%
80%
60%
40%
20%
0%
50
40
30
20
10
0
Time before present (kya)
0%
10%
20%
30%
40%
50%
60%
Present-day focal allele frequency (same color in every panel)


100 neutrals: True TMRCA
Focal site: 500 kb | 1,000 fixed pairs | Present-day segregation required
100 individual neutrals
Cohort median
Pointwise p < 0.05 boundary
Cumulative: TMRCA below cutoff
Pair fraction
0%
20%
40%
10
20
30
40
50
TMRCA cutoff (kya)
Interval: TMRCA within each 1-kya bin
Pair fraction
0.0%
5.0%
10.0%
10
20
30
40
50
TMRCA bin midpoint (kya)
0%
10%
20%
30%
40%
50%
60%
Present-day focal allele frequency (same color in every panel)


100 neutrals: Posterior median
Focal site: 500 kb | 1,000 fixed pairs | Present-day segregation required
100 individual neutrals
Cohort median
Pointwise p < 0.05 boundary
Cumulative: TMRCA below cutoff
Pair fraction
0%
20%
40%
60%
10
20
30
40
50
TMRCA cutoff (kya)
Interval: TMRCA within each 1-kya bin
Pair fraction
0.0%
2.0%
4.0%
10
20
30
40
50
TMRCA bin midpoint (kya)
0%
10%
20%
30%
40%
50%
60%
Present-day focal allele frequency (same color in every panel)


100 neutrals: Posterior-weighted mass
Focal site: 500 kb | 1,000 fixed pairs | Present-day segregation required
100 individual neutrals
Cohort median
Pointwise p < 0.05 boundary
Cumulative: TMRCA below cutoff
Pair fraction
60%
40%
20%
0%
10
20
30
40
50
TMRCA cutoff (kya)
Interval: TMRCA within each 1-kya bin
Pair fraction
2.00%
1.00%
0.00%
10
20
30
40
50
TMRCA bin midpoint (kya)
0%
10%
20%
30%
40%
50%
60%
Present-day focal allele frequency (same color in every panel)