

Derivation of gradient descent training rule for a single unit with output O

where $O = w_0 + w_1 x_1 + x_2 x_2 + x_1 x_1^2 + x_2 x_2^2 + \dots + x_n x_n + x_n x_n^2 \quad (*)$

Derivation

We first need to note that
 \Rightarrow The training Error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where

Derivation of gradient descent training rule for a single unit with output O

where $O = w_0 + w_1 x_1 + x_2 x_2 + x_1 x_1^2 + x_2 x_2^2 + \dots + x_n x_n + x_n x_n^2 \quad (*)$

Derivation

We first need to note that
 \Rightarrow The training Error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where $t_d \Rightarrow$ is our training value
 $O_d \Rightarrow$ is output function value

\Rightarrow Note that in our given output function with bias and weight given as in $(*)$, It is represented as some input unit input and some polynomial function which

Derivation of gradient descent training rule for a single unit with output O

where $O = w_0 + w_1 x_1 + x_1 x_1^2 + x_2 x_2^2 + \dots + x_n x_n + x_n x_n^2 \quad (*)$

Derivation

We first need to note that
 \Rightarrow The training Error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where $t_d \Rightarrow$ is our training value
 $O_d \Rightarrow$ is output function value

\Rightarrow Note that in our given output function with bias and weight given as in $(*)$, It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values x ;

\Rightarrow Now we need to differentiate the ~~error~~ training error function

(1) with respect to each individual weights

$$\frac{\partial E}{\partial w} = \left[\frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

and with output 0

where $O = w_0 + w_1x_1 + x_1x_2 + x_1x_1^2 + x_2x_2^2 + \dots + x_nx_n + x_nx_n^2 \dots (x)$

Derivation

We first need to note that

\Rightarrow The training error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where $t_d \Rightarrow$ is our training value

$O_d \Rightarrow$ is output function value

\Rightarrow Note that in our given output function with bias and weights given as in (x), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values x_i ;

\Rightarrow Now we need to differentiate the ~~error~~ training error function (1) with respect to each individual weights

$$\Rightarrow \frac{\partial E}{\partial w_i} = \left[\frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

\Rightarrow Differentiating eqn (1) we have

$$\frac{\partial E}{\partial w_i} = \frac{1}{2} \sum_{d \in D} 2[t_d - O_d] \frac{\partial (t_d - O_d)}{\partial w_i} \quad \text{--- (3)}$$

we first need to note that
 \Rightarrow The training error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where $t_d \Rightarrow$ is our training value
 $O_d \Rightarrow$ is output function value

\Rightarrow Note that in our given output function with bias and weights given as in (x), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values, x_i

\Rightarrow Now we need to differentiate the ~~error~~ training error function (1) with respect to each individual weights

$$\Rightarrow \frac{\partial E}{\partial w_i} = \left[\frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

\Rightarrow Differentiating eqn (1) we have

$$\frac{\partial E}{\partial w_i} = \frac{1}{2} \sum_{d \in D} 2[t_d - O_d] \frac{\partial}{\partial w_i} (t_d - O_d) \quad \text{--- (3)}$$

\Rightarrow Since everything is a vector representation, whether it is weights or inputs, so we can basically represent (3) as

$$\Delta \vec{w} = (t_d - O_d) \sum_{d \in D} \frac{\partial}{\partial w_i} (t_d - O_d) \quad \text{--- (4)}$$

where $t_d \Rightarrow$ is our training value
 $O_d \Rightarrow$ is output function value
 \Rightarrow Note that in our given output function with bias and weight given as in (*), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector value, x_i

\Rightarrow Now we need to differentiate the error training error function (1) with respect to each individual weights

$$\Rightarrow \frac{\partial E}{\partial w_i} = \left[\frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad (2)$$

\Rightarrow Differentiating eqn (1) we have

$$\frac{\partial E}{\partial w_i} = \frac{1}{2} \sum_{d \in D} 2 [t_d - O_d] \frac{\partial (t_d - O_d)}{\partial w_i} \quad (3)$$

\Rightarrow Since everything is a vector representation, whether it is weights or inputs, so we can basically represent (3) as

$$\Delta \vec{w} = (t_d - O_d) \sum_{d \in D} \frac{\partial (t_d - O_d)}{\partial w_i} \quad (4)$$

\Rightarrow Question is what is O_d is our case according the given output function (*)

\Rightarrow We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2 \Rightarrow O_d(\vec{X}) = \vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2 \quad \text{--- (5)}$$

where \vec{w} is the weight vector that is general for all weights regardless of whether it is w_0, w_1, w_2, \dots so on
 \vec{X}_d is the input vector at each instance

Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2)] \quad \text{--- (6)}$$

Now, differentiating eqn (6) we have

$$\sum_{d \in D} [t_d - O_d] [0 - \vec{X}_{i,d} - \vec{X}_{i,d}^2]$$

in x we have

⇒ Question is what is O_d is our case according the given output function (*)

⇒ We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2 \Rightarrow O_d(\vec{x}) = \vec{w} \vec{x}_d + \vec{w} \vec{x}_d^2 \quad \text{--- (5)}$$

where \vec{w} is the weight vector that is general for all weights regardless of whether it is w_0, w_1, w_2, \dots so on
 \vec{x}_d is the input vector at each instance

⇒ Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{x}_d + \vec{w} \vec{x}_d^2)] \quad \text{--- (6)}$$

Now, differentiating eqn (6) we have

$$\sum_{d \in D} [t_d - O_d] [0 - \vec{x}_{i,d} - \vec{x}_{i,d}^2]$$

Factoring out negative in x_i we have

$$= \sum_{d \in D} [t_d - O_d] - (\vec{x}_{i,d} + \vec{x}_{i,d}^2)$$

\Rightarrow Question is what is O_d is our case according to the given output function (*)

\Rightarrow We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2 \Rightarrow O_d(\vec{x}) = \vec{w} \vec{x}_d + \vec{w} \vec{x}_d^2 \quad \text{--- (5)}$$

where \vec{w} is the weight vector that is general for all weights regardless of whether it is w_0, w_1, w_2, \dots so on
 \vec{X}_d is the input vector at each instance

\Rightarrow Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2)] \quad \text{--- (6)}$$

\Rightarrow Now, differentiating eqn (6) we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] [0 - \vec{X}_{i,d} - \vec{X}_{i,d}^2]$$

\Rightarrow Factoring out negative in x_i we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] - (\vec{X}_{i,d} + \vec{X}_{i,d}^2)$$

$$\text{Error function} \Rightarrow - \sum_{d \in D} [t_d - O_d] [\vec{X}_{i,d} + \vec{X}_{i,d}^2] = \nabla E(\vec{w})$$

output function (*)

\Rightarrow We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2 \Rightarrow O_d(\vec{X}) = \vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2 \quad \text{--- (5)}$$

where \vec{w} is the weight vector that is general for all weights regardless of whether it is w_0, w_1, w_2, \dots so on
 \vec{X}_d is the input vector at each instance

\Rightarrow Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{X}_d + \vec{w} \vec{X}_d^2)] \quad \text{--- (6)}$$

\Rightarrow Now, differentiating eqn (6) we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] [0 - \vec{X}_{i,d} - \vec{X}_{i,d}^2]$$

\Rightarrow Factoring out negative in x_i we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] - (\vec{X}_{i,d} + \vec{X}_{i,d}^2)$$

$$\text{Error function} \Rightarrow - \sum_{d \in D} [t_d - O_d] [\vec{X}_{i,d} + \vec{X}_{i,d}^2] = \nabla E(\vec{w})$$

- let's have a learning rate given by η

$$\Rightarrow \Delta \vec{w} = -\eta \nabla E(\vec{w})$$

\Rightarrow This can be represented in other notation as

$$\Delta w_i = \eta \frac{\partial E}{\partial w_i}$$

\Rightarrow Thus the training rule for gradient descent is

$$\vec{w}_i \leftarrow \vec{w}_i + \Delta w_i$$

\Rightarrow Therefore the modified weight equation is given by

$$w_i \leftarrow w_i - \eta \sum_{d \in D} [t_d - O_d] [\vec{x}_{i,d} + \vec{x}_{i,d}^2]$$

$$\Rightarrow \Delta \vec{w} = -\eta \nabla E(\vec{w})$$

\Rightarrow This can be represented in other notation as

$$\Delta w_i = \eta \frac{\partial E}{\partial w_i}$$

\Rightarrow Thus the training rule for gradient descent is

$$\vec{w}_i \leftarrow \vec{w}_i + \Delta w_i$$

\Rightarrow Therefore the modified weight equation is given by

$$w_i \leftarrow w_i - \eta \sum_{d \in D} [t_d - O_d] [\vec{x}_{i,d} + \vec{x}_{i,d}^2]$$