

Derivation of gradient descent training rule for a single unit with output  $O$

where  $O = w_0 + w_1x_1 + x_2x_2 + x_1x_1^2 + x_2x_2^2 + \dots + x_nx_n + x_nx_n^2 \dots (*)$

Derivation

We first need to note that

$\Rightarrow$  The training Error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where

Derivation of gradient descent training rule for a single unit with output  $O$

where  $O = w_0 + w_1x_1 + x_2x_2 + x_1x_1^2 + x_2x_2^2 + \dots + x_nx_n + x_nx_n^2 \dots (*)$

Derivation

We first need to note that

$\Rightarrow$  The training Error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where  $t_d \Rightarrow$  is our training value  
 $O_d \Rightarrow$  is output function value

$\Rightarrow$  Note that in our given output function with bias and weight given as in  $(*)$ , It is represented as some linear unit input and some polynomial function which

Derivation of gradient descent training rule for a single unit with output  $O$

where  $O = w_0 + w_1x_1 + w_2x_2 + w_3x_3 + w_4x_4 + \dots + w_nx_n + w_{n+1}x_{n+1}^2 \dots$  (1)

Derivation

We first need to note that

$\Rightarrow$  The training error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where  $t_d \Rightarrow$  is our training value  
 $O_d \Rightarrow$  is output function value

$\Rightarrow$  Note that in our given output function with bias and weight given as in (\*), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values  $x$ ;

$\Rightarrow$  Now we need to differentiate the ~~error~~ training error function (1) with respect to each individual weights

$$\frac{\partial E}{\partial w_i} = \left[ \frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

and with output  $O$

where  $O = w_0 + w_1x_1 + w_2x_2 + w_3x_3 + w_4x_4 + \dots + w_nx_n + w_{n+1}x_{n+1}^2 \dots$  (1)

Derivation

We first need to note that

$\Rightarrow$  The training error function depending on weights and bias is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - O_d)^2 \quad \text{--- (1)}$$

where  $t_d \Rightarrow$  is our training value  
 $O_d \Rightarrow$  is output function value

$\Rightarrow$  Note that in our given output function with bias and weight given as in (\*), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values  $x$ ;

$\Rightarrow$  Now we need to differentiate the ~~error~~ training error function (1) with respect to each individual weights

$$\Rightarrow \frac{\partial E}{\partial w_i} = \left[ \frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

$\Rightarrow$  Differentiating eqn (1) we have

$$\frac{\partial E}{\partial w_i} = \frac{1}{2} \sum_{d \in D} 2[t_d - O_d] \frac{\partial (t_d - O_d)}{\partial w_i} \quad \text{--- (3)}$$

we first need to note that  
 $\Rightarrow$  The training error function depending on weights and bias  
 is given by

$$E(\vec{w}) = \frac{1}{2} \sum_{d \in D} (t_d - o_d)^2 \quad \text{--- (1)}$$

where  $t_d \Rightarrow$  is our training value  
 $o_d \Rightarrow$  is output function value

$\Rightarrow$  Note that in our given output function with bias and weights given as in (\*), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values  $x_i$ ;

$\Rightarrow$  Now we need to differentiate the ~~error~~ training error function (1) with respect to each individual weights

$$\Rightarrow \frac{\partial E}{\partial w_i} = \left[ \frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

$\Rightarrow$  Differentiating eqn (1) we have

$$\frac{\partial E}{\partial w_i} = \frac{1}{2} \sum_{d \in D} 2[t_d - o_d] \frac{\partial}{\partial w_i} (t_d - o_d) \quad \text{--- (3)}$$

$\Rightarrow$  Since everything is a vector representation, whether it is weights or inputs, so we can basically represent (3) as

$$\Delta \vec{w} = (t_d - o_d) \sum_{d \in D} \frac{\partial}{\partial w_i} (t_d - o_d) \quad \text{--- (4)}$$

where  $t_d \Rightarrow$  is our training value  
 $o_d \Rightarrow$  is output function value

$\Rightarrow$  Note that in our given output function with bias and weights given as in (\*), It is represented as some linear unit input and some polynomial function which is not completely linear but only linear in weights but non-linear in input vector values  $x_i$ ;

$\Rightarrow$  Now we need to differentiate the ~~error~~ training error function (1) with respect to each individual weights

$$\Rightarrow \frac{\partial E}{\partial w_i} = \left[ \frac{\partial E}{\partial w_0}, \frac{\partial E}{\partial w_1}, \frac{\partial E}{\partial w_2}, \frac{\partial E}{\partial w_3}, \dots, \frac{\partial E}{\partial w_n}, \frac{\partial E}{\partial w_{n+1}} \right] \quad \text{--- (2)}$$

$\Rightarrow$  Differentiating eqn (1) we have

$$\frac{\partial E}{\partial w_i} = \frac{1}{2} \sum_{d \in D} 2[t_d - o_d] \frac{\partial}{\partial w_i} (t_d - o_d) \quad \text{--- (3)}$$

$\Rightarrow$  Since everything is a vector representation, whether it is weights or inputs, so we can basically represent (3) as

$$\Delta \vec{w} = (t_d - o_d) \sum_{d \in D} \frac{\partial}{\partial w_i} (t_d - o_d) \quad \text{--- (4)}$$



=> Question is what is  $O_d$  is our case according the given output function (\*)

=> We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} X_d^2 \Rightarrow O_d(\vec{x}) = \vec{w} \vec{X}_d + \vec{w} X_d^2 \quad \text{--- (5)}$$

where  $\vec{w}$  is the weight vector that is general for all weights regardless of whether it is  $w_0, w_1, w_2, \dots$  so on  
 $\vec{X}_d$  is the input vector at each instance

Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{X}_d + \vec{w} X_d^2)] \quad \text{--- (6)}$$

Now, differentiating eqn (6) we have

$$\sum_{d \in D} [t_d - O_d] [0 - \vec{X}_{i,d} - 2\vec{X}_{i,d}]$$

in x: we have

=> Question is what is  $O_d$  is our case according the given output function (\*)

=> We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} X_d^2 \Rightarrow O_d(\vec{x}) = \vec{w} \vec{X}_d + \vec{w} X_d^2 \quad \text{--- (5)}$$

where  $\vec{w}$  is the weight vector that is general for all weights regardless of whether it is  $w_0, w_1, w_2, \dots$  so on  
 $\vec{X}_d$  is the input vector at each instance

=> Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{X}_d + \vec{w} X_d^2)] \quad \text{--- (6)}$$

Now, differentiating eqn (6) we have

$$\sum_{d \in D} [t_d - O_d] [0 - \vec{X}_{i,d} - 2\vec{X}_{i,d}]$$

Factoring out negative in x: we have

$$\sum_{d \in D} [t_d - O_d] - (\vec{X}_{i,d} + 2\vec{X}_{i,d})$$

$\Rightarrow$  Question is what is  $O_d$  is our case according to given output function (x)

$\Rightarrow$  We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} X_d^2 \Rightarrow O_d(\vec{x}) = \vec{w} \vec{x}_d + \vec{w} x_d^2 \quad \text{--- (5)}$$

where  $\vec{w}$  is the weight vector that is general for all weights regardless of whether it is  $w_0, w_1, w_2, \dots$  so on  
 $\vec{x}_d$  is the input vector at each instance

$\Rightarrow$  Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{x}_d + \vec{w} x_d^2)] \quad \text{--- (6)}$$

$\Rightarrow$  Now, differentiating eqn (6) we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] [0 - \vec{x}_{i,d} - \vec{x}_{i,d}^2]$$

$\Rightarrow$  Factoring out negative in  $x_i$  we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] - (\vec{x}_{i,d} + \vec{x}_{i,d}^2)$$

$$\text{Error function} \Rightarrow - \sum_{d \in D} [t_d - O_d] [\vec{x}_{i,d} + \vec{x}_{i,d}^2] = \nabla E(\vec{w})$$

output function (x)

$\Rightarrow$  We need to write a general representation of our output function as below

$$O_d = \vec{w} \vec{X}_d + \vec{w} X_d^2 \Rightarrow O_d(\vec{x}) = \vec{w} \vec{x}_d + \vec{w} x_d^2 \quad \text{--- (5)}$$

where  $\vec{w}$  is the weight vector that is general for all weights regardless of whether it is  $w_0, w_1, w_2, \dots$  so on  
 $\vec{x}_d$  is the input vector at each instance

$\Rightarrow$  Substituting (5) into (4) we have

$$= \sum_{d \in D} [t_d - O_d] \frac{\partial}{\partial w_i} [t_d - (\vec{w} \vec{x}_d + \vec{w} x_d^2)] \quad \text{--- (6)}$$

$\Rightarrow$  Now, differentiating eqn (6) we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] [0 - \vec{x}_{i,d} - \vec{x}_{i,d}^2]$$

$\Rightarrow$  Factoring out negative in  $x_i$  we have

$$\Rightarrow \sum_{d \in D} [t_d - O_d] - (\vec{x}_{i,d} + \vec{x}_{i,d}^2)$$

$$\text{Error function} \Rightarrow - \sum_{d \in D} [t_d - O_d] [\vec{x}_{i,d} + \vec{x}_{i,d}^2] = \nabla E(\vec{w})$$

- let's have a learning rate given by  $\eta$

$$\Rightarrow \Delta \vec{w} = -\eta \nabla E(\vec{w})$$

$\Rightarrow$  This can be represented in other notation as

$$\Delta w_i = \eta \frac{\partial E}{\partial w_i}$$

$\Rightarrow$  Thus the training rule for gradient descent is

$$\vec{w}_i \leftarrow \vec{w}_i + \Delta w_i$$

$\Rightarrow$  Therefore the modified weight equation is given by

$$w_i \leftarrow w_i - \eta \sum_{d \in D} [t_d - O_d] [\vec{x}_{i,d} + \vec{x}_{i,d}^2]$$

$$\Rightarrow \Delta \vec{w} = -\eta \nabla E(\vec{w})$$

$\Rightarrow$  This can be represented in other notation as

$$\Delta w_i = \eta \frac{\partial E}{\partial w_i}$$

$\Rightarrow$  Thus the training rule for gradient descent is

$$\vec{w}_i \leftarrow \vec{w}_i + \Delta w_i$$

$\Rightarrow$  Therefore the modified weight equation is given by

$$w_i \leftarrow w_i - \eta \sum_{d \in D} [t_d - O_d] [\vec{x}_{i,d} + \vec{x}_{i,d}^2]$$