

# An unconditional bound for a Möbius-weighted correlation sum in fixed residue classes, with two consequences for the Huang–Li reduction of Goldbach to Elliott–Halberstam

Youngrok Lee

Independent researcher, Republic of Korea

## Abstract

Huang and Li [2] show that the binary Goldbach conjecture for large even  $N$  follows from  $EH$  together with a Möbius-twisted variant  $EH_\mu$  whose levels sum to more than 1; by Bombieri–Vinogradov their Corollary 1 reduces this to  $EH_\mu(N^{\theta'})$  alone for a single  $\theta' > 1/2$ . Their proof consumes  $EH_\mu$  at exactly two places, the functionals they call  $E_3(\alpha)$  and  $E_4(\alpha)$ . We prove unconditionally (Theorem 1) that the Möbius-weighted correlation sum underlying  $E_4$  is  $\ll_A N(\log N)^{-A}$  for every  $A > 0$ , uniformly in the truncation point; consequently (Corollary 2) the  $E_4$  consumption of  $EH_\mu$  is unnecessary and the entire  $EH_\mu$  demand of the Huang–Li argument collapses to the single scalar  $E_3(\alpha)$ . We then show (Theorem 3) that the corresponding statement for  $E_3$  — the same sum with the weight  $\log k$  — is not merely hard but, by an unconditional identity, *equivalent* to Huang–Li’s own equation (22): it therefore already yields binary Goldbach for large even  $N$ , and yields the asymptotic  $\tilde{r}(N) \sim \mathfrak{S}(N)N$  exactly when  $\sum_{n < N} \Lambda(n)\mu(N-n) = o(N)$ . The root cause is  $\mu * \log = \Lambda$ . We then locate the true strength of the surviving demand: Goldbach needs only a one-sided bound on  $E_3$  (Proposition 5), at a threshold of order  $N$  for almost all  $N$ , and hence only a constant-factor bound on exactly the object  $EH_\mu$  bounds (Proposition 6). Finally we record a defect in [2]: equation (18) drops an  $n$ -dependent constraint present in the definition of  $S_2(\alpha)$ ; we give the missing term and show it is harmless under the hypotheses already assumed (Section 7).

We state plainly what is not claimed. Theorem 1 removes only the part of the demand that carries no Goldbach content, and the net progress toward the Goldbach conjecture is zero. We also state plainly what is standard: the ingredients of Theorem 1 are classical and its mechanism — a divisor switch that moves the work onto a short variable, followed by Bombieri–Vinogradov — is ordinary practice. What is offered is the application, together with the corrections it turns out to require.

## 1 Introduction

### 1.1 The Huang–Li reduction

Let  $N$  be a large even integer,  $\Lambda$  the von Mangoldt function,  $\mu$  the Möbius function, and

$$\tilde{r}(N) = \sum_{n < N} \Lambda(n)\Lambda(N-n), \quad \mathfrak{S}(N) = 2 \prod_{p > 2} \left(1 - \frac{1}{(p-1)^2}\right) \prod_{\substack{p|N \\ p > 2}} \left(1 + \frac{1}{p-2}\right).$$

Write  $EH_\mu(Q)$  for the assertion that for every  $A > 0$

$$\sum_{q \leq Q} \max_{y < N} \max_{(a,q)=1} \left| \sum_{\substack{n \leq y \\ n \equiv a \pmod{q}}} \Lambda(n)\mu(N-n) - \frac{1}{\varphi(q)} \sum_{n \leq y} \Lambda(n)\mu(N-n) \right| \ll_A \frac{N}{(\log N)^A}. \quad (1)$$

Huang and Li [2], continuing Pan [8], prove that  $EH(N^\theta(\log N)^{2A+8})$  together with  $EH_\mu(N^{1-\theta})$  implies

$$\tilde{r}(N) \geq \mathfrak{S}(N)(1 - \mathfrak{A}(N))N + O\left(\frac{N}{(\log N)^A}\right), \quad (2)$$

and deduce (their Corollary 1) that, in view of Bombieri–Vinogradov, binary Goldbach for all sufficiently large even integers follows from  $EH_\mu(N^{\theta'})$  alone for any single  $\theta' > 1/2$ . Here

$$\mathfrak{A}(N) = \prod_{\substack{q \nmid N \\ q > 2}} \left(1 - \frac{1}{q(q-1)}\right) \quad (3)$$

is their equation (7);  $N$  is even, so  $q > 2$  is implied by  $q \nmid N$  and is written here only to match their display. We write  $\mathfrak{A}$  rather than their  $A$  because  $A$  is already the free exponent in (1), and one symbol must not carry two meanings.

This is the sharpest conditional frame we are aware of: one sentence stands between classical technology and Goldbach. This paper asks what that sentence costs on the side where it is consumed.

## 1.2 Where $EH_\mu$ is consumed

The hypothesis (1) enters the Huang–Li proof at exactly two places. With  $\alpha$  the Vaughan-type parameter of their §3 and  $K = (N-1)/\alpha$ , these are (their notation; note that the residue class is the *fixed* class  $a \equiv N$ ):

$$E_3(\alpha) = \sum_{\substack{k < K \\ (k, N)=1}} \mu(k) \log k \left[ \sum_{\substack{n < N \\ n \equiv N \pmod{k}}} \Lambda(n) \mu(N-n) - \frac{1}{\varphi(k)} \sum_{n < N} \Lambda(n) \mu(N-n) \right], \quad (4)$$

$$E_4(\alpha) = \sum_{\substack{k < K \\ (k, N)=1}} \mu(k) \left[ \sum_{\substack{n < N \\ n \equiv N \pmod{k}}} \Lambda(n) \mu(N-n) \log(N-n) - \frac{1}{\varphi(k)} \sum_{n < N} \Lambda(n) \mu(N-n) \log(N-n) \right]. \quad (5)$$

In both cases Huang–Li discard the signs  $\mu(k)$  by the triangle inequality on the first line and then appeal to (1) (their Lemma 4 for  $E_4$ ). The two functionals differ in exactly one respect:  $E_3$  carries the weight  $w_k = \log k$  and  $E_4$  carries  $w_k = 1$  (the factor  $\log(N-n)$  inside  $E_4$  is  $k$ -independent and is removed by partial summation). This paper shows that this one difference decides everything.

## 1.3 Results

For  $(k, N) = 1$  and  $1 \leq t < N$  put

$$E_\mu(t; k) := \sum_{\substack{n \leq t \\ n \equiv N \pmod{k}}} \Lambda(n) \mu(N-n) - \frac{1}{\varphi(k)} \sum_{n \leq t} \Lambda(n) \mu(N-n), \quad C(t) := \sum_{n \leq t} \Lambda(n) \mu(N-n), \quad (6)$$

and, for a weight  $w$ ,

$$T_w(t) := \sum_{\substack{k < K \\ (k, N)=1}} \mu(k) w(k) E_\mu(t; k).$$

Thus  $E_4$  is obtained from  $T_1$  and  $E_3$  from  $T_{\log}$ .

**Theorem 1.** Fix  $\theta' \in (1/2, 1)$  and put  $K = N^{\theta'}$ . Then for every  $A > 0$ ,

$$\sup_{1 \leq t < N} |T_1(t)| = \sup_{1 \leq t < N} \left| \sum_{\substack{k < K \\ (k, N)=1}} \mu(k) E_\mu(t; k) \right| \ll_{A, \theta'} \frac{N}{(\log N)^A}.$$

Moreover every ingredient of the proof except Bombieri–Vinogradov contributes only  $O\left(Ne^{-c\sqrt{\log N}}\right)$ : the power of  $\log$  in the statement is imposed by Bombieri–Vinogradov alone.

Theorem 1 is a statement about a *signed* sum in a *fixed* residue class. It is not an instance of, and does not imply,  $EH_\mu(N^{\theta'})$ , which asks for absolute values and a maximum over residue classes. The mechanism is described in Section 2: after divisor switching one *completes* the divisor sum, and on the complementary range  $k \geq K$  the Möbius factor on the long variable squares itself away while the surviving Möbius sits on a variable of size  $\leq N^{1/2-\delta}$ . The completion is not a formality — without it the cofactor is not short and the configuration is the one with no known machine.

**Corollary 2.** In the Corollary-1 regime,  $E_4(\alpha) \ll_A N(\log N)^{-A}$  for every  $A > 0$ , unconditionally. Consequently Huang–Li’s estimate  $S_4(\alpha) = O(N(\log N)^{-A})$  requires no hypothesis, their Lemma 4 is not needed, and the entire  $EH_\mu$  demand of their argument collapses to the single scalar  $E_3(\alpha)$  — together with the correction  $\Delta$  of Section 7, which is likewise unconditional there.

**Theorem 3.** In the Corollary-1 regime one has the unconditional identity

$$E_3(\alpha) = \sum_{n < N} \Lambda(n) \Lambda(N - n) - \mathfrak{S}(N) \left( N - \sum_{n < N} \Lambda(n) \mu(N - n) \right) + O_A \left( \frac{N}{(\log N)^A} \right).$$

In particular the bound  $E_3(\alpha) \ll_A N(\log N)^{-A}$  — the form of  $EH_\mu$  that the Huang–Li argument consumes, though not the weakest that suffices; see Proposition 5 — is equivalent to Huang–Li’s equation (22). It therefore yields binary Goldbach for large even  $N$  through their Theorem 1; and, granted it, the asymptotic  $\tilde{r}(N) \sim \mathfrak{S}(N)N$  holds if and only if  $C(N) = o(N)$ .

Note 4 (three statements, not two). The identity says that the displayed difference is  $E_3(\alpha)$ , so the hypothesis  $E_3 \ll_A N(\log N)^{-A}$  and Huang–Li’s (22) are the same assertion. What [2] record immediately after their (22) is a different equivalence, between  $\tilde{r}(N) \sim \mathfrak{S}(N)N$  and  $C(N) = o(N)$ . Chaining the two,  $E_3 \ll_A N(\log N)^{-A}$  delivers binary Goldbach — Huang–Li’s Theorem 1 supplies  $|C(N)| \leq \mathfrak{A}(N)N(1 + o(1))$  by the triangle inequality, and  $\mathfrak{A}(N) < 1$  — but it does not on its own deliver the asymptotic, which needs  $C(N) = o(N)$  in addition. The three statements must be kept apart.

Theorem 3 is the more consequential half. It says that the demand side is closed at the level of *identities* rather than of *estimates*: no choice of  $\theta'$ , of truncation, or of smoothing can evade it, because the root cause is the identity  $\mu * \log = \Lambda$  from which Huang–Li start. The weight  $\log k$  is the carrier of the Goldbach content, so every divisor switch must hand it back.

Closure at the level of identities is not closure at the level of *strength*, and the next two propositions separate the two. Every condition on  $E_3$  unwinds, through Theorem 3, to a condition on  $\tilde{r}(N)$  and  $C(N)$  — that is the identity part, and it is permanent. But Huang–Li consume  $E_3$  two-sidedly and at a saving of every power of  $\log$ , and Goldbach needs neither.

**Proposition 5** (the demand is one-sided, and its threshold is not a log power). In the Corollary-1 regime, binary Goldbach for all sufficiently large even  $N$  follows from

$$E_3(\alpha) > -\mathfrak{S}(N)(1 - \mathfrak{A}(N))N(1 + o(1)), \quad (7)$$

with  $\mathfrak{A}(N)$  as in (3). This is strictly weaker than  $E_3 \ll_A N(\log N)^{-A}$  in two independent ways: it constrains  $E_3$  from one side only, and its threshold is  $\mathfrak{S}(N)(1 - \mathfrak{A}(N))N$ , which is  $\asymp N$  for almost all even  $N$  and never smaller than  $cN/(\log N \log \log N)$  for an absolute constant  $c > 0$ .

**Proposition 6** (the demand needs no saving in  $\log N$ ). *Put*

$$B(N) := \sum_{\substack{k < K \\ (k, N) = 1}} \mu^2(k) (\log k) |E_\mu(N; k)|, \quad (8)$$

the restriction to squarefree  $k$  being what the triangle inequality on (4) leaves, since  $\mu(k) = 0$  elsewhere. This is sharper than the quantity [2] actually reach before appealing to  $EH_\mu$ , namely  $\log N \sum_{k < K, (k, N) = 1} |E_\mu(N; k)|$ , in which  $\log k$  has been raised to  $\log N$ ; the proposition below holds for either, and a fortiori for theirs. Then binary Goldbach for all sufficiently large even  $N$  follows from

$$B(N) \leq (1 - \varepsilon) \mathfrak{S}(N)(1 - \mathfrak{A}(N))N \quad (9)$$

for a single fixed  $\varepsilon > 0$ . Since  $\mathfrak{S}(N)(1 - \mathfrak{A}(N)) \asymp 1$  for almost all even  $N$ , (9) is a constant-factor bound on exactly the object  $EH_\mu$  bounds — absolute values kept, and no saving in  $\log N$  whatsoever — where  $EH_\mu(N^{\theta'})$  asks for  $\ll_A N(\log N)^{-A}$  for every  $A$ .

## 1.4 What is not claimed

- **No progress toward Goldbach.** Theorem 1 is unconditional but Goldbach-neutral: it removes exactly the half of the demand that carries no Goldbach content. The net progress toward the Goldbach conjecture is zero.
- **No new estimate for  $C(N)$ .** Theorem 3 places the whole surviving demand on  $C(N) = \sum_{n < N} \Lambda(n) \mu(N - n)$ ; nothing here bounds that quantity.
- **No claim of novelty of mechanism.** The ingredients of Theorem 1 are classical (Section 8). Its difficulty is bookkeeping, and the two places where the bookkeeping bites are isolated in Section 2 and Remark 9.
- **The weakenings do not reopen the design space.** Propositions 5 and 6 lower the required strength but not the required level; the divisor-switch route remains closed over its whole weight space [7].
- **The defect report is not a retraction.** [2]’s Theorem 1 and Corollary 1 stand as stated; what Section 7 supplies is a missing term and its treatment.

## 2 Notation and the mechanism

$p$  and  $q$  always denote primes.  $\varphi$  is Euler’s function,  $\tau_3$  the 3-fold divisor function,  $\text{rad}(u) = \prod_{p|u} p$ . Constants  $c, c_1, \dots$  are positive and absolute unless subscripted; implied constants may depend on  $A$  and  $\theta'$ . The letter  $A$  is reserved for the free exponent of (1); the local factor of (3) is  $\mathfrak{A}$ .

The mechanism of Theorem 1 is the following. The residue class in (6) is the fixed class  $n \equiv N \pmod{k}$ , i.e.

$$n \equiv N \pmod{k} \iff k \mid N - n.$$

So the  $k$ -sum is a divisor sum over  $u = N - n$ , namely the incomplete sum  $\sigma_K(u) = \sum_{k|u, k < K, (k, N) = 1} \mu(k)$ , and the mechanism has three steps rather than one.

*First one completes it.* By Lemma 7, for squarefree  $u$  the complete sum  $\sum_{k|u, (k, N) = 1} \mu(k)$  equals  $\mathbf{1}_{\text{rad}(u)|N}$ , which forces  $u \mid \text{rad}(N)$  and so leaves only  $N^{o(1)}$  terms. *Then the work sits*

entirely in the complementary sum, over  $k \geq K$  — and it is only on that range that writing  $u = mk$  gives

$$m = u/k < N/K = N^{1-\theta'} \leq N^{1/2-\delta}.$$

Finally, for squarefree  $u = mk$  the factorisation forces  $(m, k) = 1$  and  $\mu(u)\mu(k) = \mu(m)\mu^2(k)$ : the Möbius factor on the *long* variable  $k$  cancels against itself and leaves the nonnegative  $\mu^2$ , while the surviving Möbius sits on the *short* variable  $m$ . That is the classical “Möbius on the short variable plus Bombieri–Vinogradov” configuration.

The completion step is load-bearing and easy to omit. Over the range  $k < K$  as it stands the cofactor  $u/k$  runs up to  $u$  itself, so the bound  $m < N^{1-\theta'}$  is simply false there and the surviving Möbius would sit on the *long* variable — the configuration for which no machine is known. Steps 1–3 of Section 3 carry out the argument in that order.

### 3 Proof of Theorem 1

Fix  $A > 0$  and set

$$D_0 := (\log N)^{A+2}, \quad E_0 := (\log N)^{2A+4}. \quad (10)$$

The truncations must be allowed to depend on  $A$ ; a fixed choice covers only bounded  $A$ .

#### 3.1 Step 0: the subtracted mean term

By (6),

$$T_1(t) = D(t) - C(t)B(K), \quad D(t) := \sum_{\substack{k < K \\ (k, N)=1}} \mu(k) \sum_{\substack{n \leq t \\ n \equiv N \pmod{k}}} \Lambda(n) \mu(N - n), \quad (11)$$

where  $B(K) = \sum_{k < K, (k, N)=1} \mu(k)/\varphi(k)$ . Huang–Li’s Lemma 1 (a form of Goldston–Yıldırım [1]) gives  $B(K) \ll e^{-c_1 \sqrt{\log K}}$ , and trivially  $|C(t)| \leq \sum_{n < N} \Lambda(n) \ll N$ . Hence

$$|C(t)B(K)| \ll Ne^{-c\sqrt{\log N}} \quad (12)$$

uniformly in  $t$ .

#### 3.2 Step 1: divisor switching (an exact identity)

Since  $n \equiv N \pmod{k}$  with  $n \leq t < N$  is the same as  $u := N - n$  being a multiple of  $k$  in  $[N - t, N)$ ,

$$D(t) = \sum_{N-t \leq u < N} \Lambda(N - u) \mu(u) \sigma_K(u), \quad \sigma_K(u) := \sum_{\substack{k|u, k < K \\ (k, N)=1}} \mu(k). \quad (13)$$

This is a finite rearrangement, hence exact. It is verified independently by a script that accumulates the two sides in genuinely different index orders — the  $k$ -side by modular slices, the  $u$ -side against a divisor-sum array — and finds them equal to  $10^{-16}$  relative to  $N$  (`audit_switch_identity.py`).

#### 3.3 Step 2: the complete divisor sum

**Lemma 7.** For squarefree  $u \geq 1$ ,  $\sum_{k|u, (k, N)=1} \mu(k) = \mathbf{1}_{\text{rad}(u)|N}$ .

*Proof.* The sum is multiplicative in  $u$  with local factor 1 at  $p \mid N$  and  $1 - 1 = 0$  at  $p \nmid N$ . Hence it vanishes unless every prime of  $u$  divides  $N$ .  $\square$

Splitting  $\sigma_K(u)$  into the complete sum minus the tail  $k \geq K$ , (13) becomes  $D(t) = P(t) - R(t)$  with

$$P(t) = \sum_{\substack{N-t \leq u < N \\ \text{rad}(u) | N}} \Lambda(N-u) \mu(u), \quad R(t) = \sum_{N-t \leq u < N} \Lambda(N-u) \mu(u) \sum_{\substack{k|u, k \geq K \\ (k, N)=1}} \mu(k). \quad (14)$$

In  $P(t)$  the conditions  $\mu(u) \neq 0$  and  $\text{rad}(u) | N$  force  $u | \text{rad}(N)$ , so there are at most  $2^{\omega(N)} = N^{o(1)}$  terms, each  $\ll \log N$ :

$$P(t) \ll N^{o(1)}. \quad (15)$$

### 3.4 Step 3: the residual, and the degeneracy lemma

Write  $u = mk$  with  $k \geq K$ , so  $m = u/k < N/K =: M$ . For squarefree  $u$  the factorisation forces  $(m, k) = 1$  and  $\mu(u)\mu(k) = \mu(m)\mu^2(k)$ , so

$$R(t) = \sum_{m < M} \mu(m) \sum_{\substack{k \geq K, (k, m)=1, (k, N)=1 \\ N-t \leq mk < N}} \mu^2(k) \Lambda(N - mk). \quad (16)$$

**Lemma 8** (degeneracy). *The contribution to (16) of the terms with  $(k, N) > 1$  is  $O(N^{o(1)})$ . The same bound holds for the terms in which the modulus constructed in Step 4 is not coprime to  $N$ .*

*Proof.* Suppose  $p | (k, N)$  and  $\Lambda(N - mk) \neq 0$ , say  $N - mk = q^\ell$ . Since  $p | k$  and  $p | N$  we get  $p | N - mk$ , hence  $q = p$  and  $n := N - mk = p^\ell$  with  $p | N$ . The number of such  $n < N$  is  $\leq \sum_{p|N} \log N / \log p \ll (\log N)^2$ ; for each, the pairs  $(m, k)$  with  $mk = N - n$  number  $\leq \tau(N - n) \ll N^{o(1)}$ , and each term is  $\ll \log N$ .  $\square$

By Lemma 8 we may drop the condition  $(k, N) = 1$  from (16) at a cost of  $O(N^{o(1)})$ . This is essential: expanding  $\mathbf{1}_{(k, N)=1}$  by Möbius over  $e | N$  would multiply the number of Bombieri–Vinogradov calls by  $2^{\omega(N)} = N^{o(1)}$  and destroy the budget.

### 3.5 Step 4: unfolding $\mu^2$ and the coprimality

Insert  $\mu^2(k) = \sum_{d^2 | k} \mu(d)$  and  $\mathbf{1}_{(k, m)=1} = \sum_{e | (k, m)} \mu(e)$  and truncate at  $D_0, E_0$  of (10).

*Tail  $d > D_0$ .* Such terms have  $d^2 | k$ , hence  $n = N - mk \equiv N \pmod{md^2}$ ; the number of  $n \leq N$  in that progression is  $\ll N/(md^2) + 1$  and each  $\Lambda \ll \log N$ . Summing,

$$\ll \log N \sum_{m < M} \sum_{d > D_0} \left( \frac{N}{md^2} + 1 \right) \ll \frac{N(\log N)^2}{D_0} + \sqrt{NM} \ll \frac{N}{(\log N)^A} + N^{1-\theta'/2}.$$

*Tail  $e > E_0$ .* Such terms have  $e | k$  and  $e | m$ , hence  $n \equiv N \pmod{me}$ , and

$$\ll \log N \sum_{m < M} \sum_{\substack{e | m \\ e > E_0}} \left( \frac{N}{me} + 1 \right) \ll \frac{N \log N}{E_0} \sum_{m < M} \frac{\tau(m)}{m} + M \ll \frac{N(\log N)^3}{E_0} + M.$$

Both are  $\ll N(\log N)^{-A}$ . In the remaining range put  $L := \text{lcm}(d^2, e)$  and  $q := mL$ ; the conditions  $d^2 | k$ ,  $e | k$  and  $n = N - mk$  give  $n \equiv N \pmod{q}$ , and  $mk < N$ ,  $mk \geq \max(N - t, mK)$  give  $1 \leq n \leq \mathcal{T}_m(t)$  where

$$\mathcal{T}_m(t) := \min(t, N - mK). \quad (17)$$

Therefore, with  $\psi(y; q, a) = \sum_{n \leq y, n \equiv a \pmod{q}} \Lambda(n)$ ,

$$R(t) = \sum_{m < M} \mu(m) \sum_{d \leq D_0} \mu(d) \sum_{\substack{e|m \\ e \leq E_0}} \mu(e) \psi(\mathcal{T}_m(t); q, N) + O\left(\frac{N}{(\log N)^A}\right). \quad (18)$$

By Lemma 8 we may further restrict to  $(q, N) = 1$ : if  $g = (q, N) > 1$  then  $g \mid n$  forces  $n$  to be a power of a prime dividing  $N$ , and the total over the  $\ll MD_0 E_0 \tau$ -many triples is  $\ll N^{1-\theta'+o(1)}$ . *Note 9* (the false-positive trap). Restricting the *main terms* to classes with  $(q, N) = 1$  is not cosmetic. Assigning a main term  $\mathcal{T}_m/\varphi(q)$  to a degenerate class shifts the density of the whole computation by the factor  $N/\varphi(N)$ . Carried into the  $\log k$  branch of Section 5, that error produces an apparent *refutation* of  $EH_\mu$ . It is recorded here because it is the most plausible false positive in this circle of ideas.

Note the size of the modulus:  $q = m \operatorname{lcm}(d^2, e) \leq MD_0^2 E_0$ , so by (10)

$$q \leq N^{1-\theta'} (\log N)^{4A+8} = N^{1/2-\delta} (\log N)^{4A+8} \leq N^{1/2-\delta/2} \quad (19)$$

for  $N$  large. This is the level at which Bombieri–Vinogradov is applied.

### 3.6 Step 5: Bombieri–Vinogradov

**Lemma 10** ( $\tau$ -weighted Bombieri–Vinogradov). *For all  $A, B > 0$  there is  $C = C(A, B)$  such that for  $Q \leq N^{1/2} (\log N)^{-C}$ ,*

$$\sum_{q \leq Q} \tau_3(q)^B \max_{y \leq N} \max_{(a, q)=1} \left| \psi(y; q, a) - \frac{y}{\varphi(q)} \right| \ll_{A, B} \frac{N}{(\log N)^A}.$$

*Proof.* Write  $\mathcal{E}_q$  for the inner maximum. By Cauchy–Schwarz,

$$\sum_q \tau_3^B \mathcal{E}_q \leq \left( \sum_q \tau_3^{2B} \mathcal{E}_q \right)^{1/2} \left( \sum_q \mathcal{E}_q \right)^{1/2}.$$

For the first factor use the trivial bound  $\mathcal{E}_q \ll (N/\varphi(q)) \log N$ , giving  $\ll N (\log N)^{C_1(B)}$ ; for the second use Bombieri–Vinogradov [6] with exponent  $2A + C_1(B)$ .  $\square$

A modulus  $q$  arises in (18) from at most  $\sum_{m|q} \tau(q/m)^2 \leq \tau(q)^3 \leq \tau_3(q)^3$  triples  $(m, d, e)$ : given  $m \mid q$  there are at most  $\tau(q/m)$  choices of  $d$  with  $d^2 \mid q/m$  and at most  $\tau(q/m)$  of  $e$  with  $\operatorname{lcm}(d^2, e) = q/m$ . Writing  $\psi(\mathcal{T}_m; q, N) = \mathcal{T}_m/\varphi(q) + \mathcal{E}$  and applying Lemma 10 with  $B = 3$  and (19),

$$R(t) = \text{MT}(t) + O\left(\frac{N}{(\log N)^A}\right), \quad \text{MT}(t) = \sum_{\substack{m < M \\ (m, N)=1}} \mu(m) \mathcal{T}_m(t) c_{D_0, E_0}(m), \quad (20)$$

where  $c_{D_0, E_0}(m) = \sum_{d \leq D_0} \sum_{e|m, e \leq E_0} \mu(d) \mu(e) \mathbf{1}_{(d, N)=1} / \varphi(m \operatorname{lcm}(d^2, e))$ .

### 3.7 Step 6: the density

**Lemma 11.** *Let  $m$  be squarefree with  $(m, N) = 1$  and put*

$$c(m) := \sum_{d \geq 1} \sum_{e|m} \frac{\mu(d) \mu(e) \mathbf{1}_{(d, N)=1}}{\varphi(m \operatorname{lcm}(d^2, e))}.$$

*Then  $c(m) = \mathfrak{A}(N) \lambda(m)/m$  with  $\mathfrak{A}(N)$  as in (3) and  $\lambda(m) = \prod_{p|m} (1 - \frac{1}{p(p-1)})^{-1}$ . Moreover  $c_{D_0, E_0}(m) = c(m) + O(1/(mD_0))$ .*



*Proof.* The sum is multiplicative. If  $p \nmid m$  then  $p \nmid e$  and the local factor is  $1 - 1/\varphi(p^2) = 1 - 1/(p(p-1))$  for  $p \nmid N$ , and 1 for  $p \mid N$ . If  $p \mid m$  (so  $p \nmid N$ ) the  $p$ -part of  $m \operatorname{lcm}(d^2, e)$  is  $p^{1+\max(2v_p(d), v_p(e))}$ , and the four choices  $(v_p(d), v_p(e)) \in \{0, 1\}^2$  contribute

$$\frac{1}{p-1} - \frac{1}{p(p-1)} - \frac{1}{p^2(p-1)} + \frac{1}{p^2(p-1)} = \frac{1}{p}.$$

Hence  $c(m) = \prod_{p \mid m} p^{-1} \cdot \prod_{p \nmid m, p \mid N} (1 - \frac{1}{p(p-1)}) = m^{-1} \mathfrak{A}(N) \lambda(m)$ . The truncation error is the tail  $\sum_{d > D_0} 1/\varphi(md^2) \ll 1/(mD_0)$  and similarly for  $e$ .  $\square$

The load-bearing line is the local factor. Equivalently, for squarefree  $m$ ,

$$\sum_{g \mid m} \frac{\mu(g)}{\varphi(m/g) g \varphi(g)} = \frac{1}{m} :$$

the local factor is exactly  $p^{-1}$ , so the density exponent is exactly 1 and  $1/\zeta$  occurs to the *first* power in Lemma 12 below. A non-integral exponent would give, by Selberg–Delange, only  $(\log x)^{-c}$  for a fixed  $c$  in Lemma 12, and Theorem 1 would be false. The identity is verified in exact rational arithmetic for all squarefree  $m < 400$ , with zero mismatches (`audit.density_identity.py`).

**Lemma 12.** *Let  $f(m) := \mu(m) \lambda(m) \mathbf{1}_{(m, N)=1}$  and  $G(x) := \sum_{m \leq x} f(m)/m$ . Then*

$$G(x) \ll \frac{N}{\varphi(N)} e^{-c\sqrt{\log x}} + x^{-1/4} e^{C\sqrt{\log N}} \quad (x \geq 2).$$

*Proof.* Write  $F(s) = \sum_m f(m) m^{-s} = \prod_{p \mid N} (1 - \lambda(p) p^{-s})$  and  $F(s) = \zeta(s)^{-1} H(s)$ , so  $f = \mu * h$  with  $h$  multiplicative,  $h(p^j) = 1$  for  $p \mid N$  and  $h(p^j) = 1 - \lambda(p) = O(p^{-2})$  for  $p \nmid N$ . Then  $G(x) = \sum_{b \leq x} \frac{h(b)}{b} \sum_{a \leq x/b} \frac{\mu(a)}{a}$ . For  $b \leq \sqrt{x}$  use  $\sum_{a \leq y} \mu(a)/a \ll e^{-c\sqrt{\log y}}$  (the classical zero-free region) together with  $\sum_b |h(b)|/b \ll \prod_{p \mid N} (1 - 1/p)^{-1} \ll N/\varphi(N)$ . For  $b > \sqrt{x}$  bound  $\sum_{b > y} |h(b)|/b \leq y^{-1/2} \sum_b |h(b)| b^{-1/2} \ll y^{-1/2} \prod_{p \mid N} (1 - p^{-1/2})^{-1} \ll y^{-1/2} e^{C\sqrt{\log N}}$ .  $\square$

Since  $M = N^{1-\theta'}$  is a fixed power of  $N$ , the second term of Lemma 12 is negligible at  $x \asymp M$ , and  $N/\varphi(N) \ll \log \log N$ ; so  $G(x) \ll e^{-c'\sqrt{\log x}}$  in the relevant range.

### 3.8 Step 7: the main term dies, uniformly in $t$

**Proposition 13.**  $\sup_{1 \leq t < N} |\operatorname{MT}(t)| \ll N e^{-c\sqrt{\log N}}.$

*Proof.* By Lemma 11,  $\operatorname{MT}(t) = \mathfrak{A}(N) \sum_{m < M} \frac{f(m)}{m} \mathcal{T}_m(t) + O(M \log N / D_0)$  with  $\mathcal{T}_m(t)$  as in (17). Put  $m_0 := (N - t)/K$ , so  $\mathcal{T}_m(t) = t$  for  $m \leq m_0$  and  $\mathcal{T}_m(t) = N - mK$  for  $m > m_0$ ; in particular  $x \mapsto T_x(t)$  is non-increasing, is constant on  $[1, m_0]$ , has derivative  $-K$  on  $(m_0, M)$ , and  $T_M(t) = 0$ . Abel summation against  $G$  of Lemma 12 gives

$$\sum_{m < M} \frac{f(m)}{m} \mathcal{T}_m(t) = G(M) T_M(t) + K \int_{m_0}^M G(x) dx = K \int_{m_0}^M G(x) dx.$$

Hence, uniformly in  $t$ ,

$$\left| \sum_{m < M} \frac{f(m)}{m} \mathcal{T}_m(t) \right| \leq K \int_1^M |G(x)| dx \ll K \int_1^M e^{-c\sqrt{\log x}} dx \ll KM e^{-c'\sqrt{\log M}} \ll N e^{-c''\sqrt{\log N}},$$

using  $\log M \asymp \log N$ . The truncation error is  $\ll M \log N / D_0 = o(N(\log N)^{-A})$ .  $\square$

The cancellation is genuine and global: the term  $m = 1$  alone contributes  $\mathcal{T}_1(t) \asymp N$  — the endpoint of (17), not the weighted sum of Theorem 1 — so the smallness of  $\operatorname{MT}$  is entirely due to the cancellation in  $\sum f(m)/m$ .



### 3.9 Conclusion

Combining (11), (12), (15), (20) and Proposition 13:

$$T_1(t) = \underbrace{P(t)}_{N^{o(1)}} - \underbrace{MT(t)}_{\ll Ne^{-c\sqrt{\log N}}} - \underbrace{O(N(\log N)^{-A})}_{BV} - \underbrace{C(t)B(K)}_{\ll Ne^{-c\sqrt{\log N}}},$$

uniformly in  $t < N$ . This proves Theorem 1, and exhibits Bombieri–Vinogradov as the only ingredient that does not give an exponential saving. The bound cannot be improved to  $Ne^{-c\sqrt{\log N}}$  by this argument: the Bombieri–Vinogradov input yields  $N(\log N)^{-A}$  for each fixed  $A$  and no more, because the Siegel–Walfisz range in its proof is  $q \leq (\log N)^B$  with  $B$  fixed.  $\square$

## 4 Proof of Corollary 2

Fix  $k$  and set  $a_n := \Lambda(n)\mu(N-n)(\mathbf{1}_{n \equiv N(k)} - 1/\varphi(k))$ , so that  $E_\mu(t; k) = \sum_{n \leq t} a_n$  and the bracket in (5) is  $\sum_{n < N} a_n \log(N-n)$ . Since  $\log(N-n)$  vanishes at  $n = N-1$  and has derivative  $-1/(N-t)$ , Abel summation gives

$$\sum_{n < N} a_n \log(N-n) = \int_1^{N-1} \frac{E_\mu(t; k)}{N-t} dt.$$

The weight  $\log(N-n)$  does not depend on  $k$ , so multiplying by  $\mu(k)$  and summing over  $k < K$  with  $(k, N) = 1$  (a finite sum) gives the exact identity

$$E_4(\alpha) = \int_1^{N-1} \frac{T_1(t)}{N-t} dt.$$

Hence  $|E_4(\alpha)| \leq (\sup_{t < N} |T_1(t)|) \log N \ll_A N(\log N)^{1-A}$  by Theorem 1. As  $A$  is arbitrary this is Corollary 2. Huang–Li’s Lemma 4 is invoked only to bound  $E_4$ ; with the above it is not needed, and the only remaining appearance of  $EH_\mu$  in their §3 is through  $E_3(\alpha)$ .  $\square$

## 5 Proof of Theorem 3: the permanent closure

Take the weight  $w_k = \log k$ , i.e. the functional  $E_3$  of (4). Steps 0–1 are unchanged, and the complete divisor sum of Lemma 7 is replaced by the following.

**Lemma 14.** *For squarefree  $u$ , write  $u = u_N u'$  where  $u_N$  is the largest divisor of  $u$  composed of primes dividing  $N$ . Then  $\sum_{k|u, (k, N)=1} \mu(k) \log k = -\Lambda(u')$ .*

*Proof.* The sum is over  $k | u'$  and equals  $-\Lambda(u')$  by  $\mu * \log = \Lambda$ .  $\square$

Hence the complete piece of  $E_3$  is

$$- \sum_{1 \leq u < N} \Lambda(N-u) \mu(u) \Lambda(u') = - \sum_{u < N, (u, N)=1} \Lambda(N-u) \mu(u) \Lambda(u) + O(N^{o(1)} \log^2 N).$$

Now  $\mu(u)\Lambda(u)$  is supported on primes  $u = p$ , where it equals  $-\log p$ ; prime powers  $u = p^\ell$ ,  $\ell \geq 2$ , have  $\mu(u) = 0$ . So the complete piece equals

$$\sum_{p < N} \Lambda(N-p) \log p + O(N^{o(1)} \log^2 N) = \sum_{n < N} \Lambda(n) \Lambda(N-n) + O(N^{1/2+\varepsilon}), \quad (21)$$

the binary Goldbach sum itself. The door is free: divisor switching costs nothing, and immediately behind it stands the object one was trying to avoid.

Two further terms have to be evaluated, and neither vanishes.

(i) *The residual main term.* Repeating Steps 3–6 with the extra factor  $\log k = \log((N - n)/m)$  produces the main term  $\mathfrak{A}(N) \sum_{m < M} \mu(m) \lambda(m) m^{-1} \int_{mK}^N \log(v/m) dv$ . Substituting  $v = mt$  evaluates the integral as  $N \log(N/m) - N - mK \log K + mK$ , of which only the  $-N \log m$  piece survives the cancellation: by Lemma 12 the coefficients  $\sum f(m)/m$  and  $\sum f(m)$  both die, and what is left is  $-\mathfrak{A}(N) N \tilde{G}(1)$  with  $\tilde{G}(1) = \lim_x \sum_{m \leq x} \mu(m) \lambda(m) \mathbf{1}_{(m, N)=1} \log m/m$ . The constant is fixed by

$$\mathfrak{A}(N) \tilde{G}(1) = -\mathfrak{S}(N), \quad (22)$$

so that the residual main term is  $+\mathfrak{S}(N) N + O(N(\log N)^{-A})$ .

The sign in (22) is forced, and it is the whole identity: with  $F(s) = \sum_m f(m) m^{-s} = \zeta(s)^{-1} H(s)$  as in Lemma 12 one has  $\tilde{G}(1) = -F'(1) = -H(1) < 0$ , while  $\mathfrak{A}(N) H(1) = \mathfrak{S}(N)$  identically — the local factors pair as

$$\left(1 - \frac{1}{p(p-1)}\right) \left(1 - \frac{1}{(p^2-p-1)(p-1)}\right) = 1 - \frac{1}{(p-1)^2}$$

at  $p \nmid N$ , and as  $p/(p-1)$  at  $p \mid N$ . Numerically,  $\mathfrak{A}(N) \tilde{G}(x) = -1.760250$  at  $x = 4 \cdot 10^6$  against  $\mathfrak{S}(N) = 1.760432$  (`audit_E3_constant.py`).

(ii) *The subtracted mean term.* It carries the factor

$$B_{\log}(K) := \sum_{\substack{k \leq K \\ (k, N)=1}} \frac{\mu(k) \log k}{\varphi(k)} = -\mathfrak{S}(N) + O(e^{-c\sqrt{\log K}} \log K),$$

which is exactly Huang–Li’s Lemma 1: subtracting  $\sum_{k \leq K} \mu(k) \varphi(k)^{-1} \log(K/k) = \mathfrak{S}(N) + O(e^{-c_1 \sqrt{\log K}})$  from  $\log K \cdot \sum_{k \leq K} \mu(k)/\varphi(k) = O(\log K e^{-c_1 \sqrt{\log K}})$  gives the claim. Independently: with  $f(s) = \prod_{p \nmid N} (1 - \frac{p^{-s}}{p-1})$  one has  $f(s) = \zeta(s+1)^{-1} h(s)$  with  $h(0) = \mathfrak{S}(N)$ , so  $B_{\log}(\infty) = -f'(0) = -\mathfrak{S}(N)$ . Since the mean term is  $-C(N) B_{\log}(K)$  with  $C(N) = \sum_{n < N} \Lambda(n) \mu(N-n)$ , it contributes  $+\mathfrak{S}(N) C(N)$ .

Collecting (21), (i) and (ii):

$$E_3(\alpha) = \sum_{n < N} \Lambda(n) \Lambda(N-n) - \mathfrak{S}(N) N + \mathfrak{S}(N) C(N) + O_A\left(\frac{N}{(\log N)^A}\right),$$

which is the assertion of Theorem 3. Since  $\mathfrak{S}(N) \asymp 1$ , the bound  $E_3(\alpha) \ll_A N(\log N)^{-A}$  is therefore equivalent to

$$\tilde{r}(N) = \mathfrak{S}(N)(N - C(N)) + O_A(N(\log N)^{-A}),$$

which is Huang–Li’s equation (22). Given that,  $\tilde{r}(N)$  is positive for large even  $N$  — binary Goldbach — because  $|C(N)| \leq \mathfrak{A}(N) N(1 + o(1))$  with  $\mathfrak{A}(N) < 1$ ; and  $\tilde{r}(N) \sim \mathfrak{S}(N) N$  holds if and only if  $C(N) = o(N)$ , which is not implied.  $\square$

*Note 15* (the identity is not accessible to computation). The finite- $N$  residual  $R(N, \theta') = |E_3(N; \theta') - (\tilde{r}(N) - \mathfrak{S}(N)(N - C(N)))|/N$  does not become small in any accessible range, and it cannot be made small by either free parameter. Over  $N = 2 \cdot 10^5, 4 \cdot 10^5, 8 \cdot 10^5$  at  $\theta' = 0.56$  it reads 0.4558, 0.3729, 0.3108, while the right-hand side it is compared against is  $\asymp 10^{-3} N$ . Swept over  $\theta' \in [0.51, 0.95]$ ,  $R$  is essentially monotone *increasing* — at  $N = 8 \cdot 10^5$  from 0.170167 at  $\theta' = 0.51$  to 4.991178 at  $\theta' = 0.90$  — because the finite- $N$  error is dominated by the main-term cancellation over  $m < M = N^{1-\theta'}$ , which  $\theta' \rightarrow 1$  destroys. The best  $\theta'$  is the smallest admissible one, just above  $1/2$ . Increasing  $N$  points the wrong way as well: at the best  $\theta'$  the ratio of error to signal reads 15.19, 18.38, 26.84 at the three  $N$  above, because  $C(N) = o(N)$  being true kills the right-hand side fast while the finite- $N$  error dies only at a slow power of  $\log$ . So Theorem 3 is a statement no computation confirms, at any  $N$ ; the proof is the only access to it (`lab_theta_sweep.py`).

*Note 16.* The mechanism is structural, not accidental. The identity  $\mu * \log = \Lambda$  is the starting point of the Huang–Li argument (their (10)); it is therefore also what any divisor switch inside their log  $k$ -weighted functional must return. No choice of  $\theta'$ , truncation, or smoothing can circumvent an identity. What the identity does not fix is the *strength* at which  $E_3$  must be bounded, and that is Section 6.

## 6 How strong the surviving demand really is

*Proof of Proposition 5.* By Theorem 3,  $\tilde{r}(N) = \mathfrak{S}(N)(N - C(N)) + E_3(\alpha) + O_A(N(\log N)^{-A})$ . The triangle inequality gives  $|C(N)| \leq U(N) := \sum_{n < N} \Lambda(n) \mu^2(N - n)$ , and  $U(N) = \mathfrak{A}(N)N(1 + o(1))$  — proved in [2, §3.3] itself by Bombieri–Vinogradov, with error  $O(N(\log N)^{-A-1})$ , and classically by Mirsky’s theorem on squarefree shifted primes [5] together with partial summation. Hence  $\mathfrak{S}(N)(N - C(N)) \geq \mathfrak{S}(N)(1 - \mathfrak{A}(N))N(1 + o(1))$ , and  $\mathfrak{A}(N) < 1$  for every  $N$ , so the right-hand side is positive. Substituting and using that the  $O_A$  term may be taken below any fixed power of  $N/\log N$  gives  $\tilde{r}(N) > 0$  under (7).

For the size of the threshold,  $\mathfrak{A}(N) = \prod_{q|N} (1 - \frac{1}{q(q-1)})$  is largest when  $N$  carries the most small primes, so  $1 - \mathfrak{A}(N)$  is smallest at primorials, where  $1 - \mathfrak{A}(N) \asymp \sum_{q > p} \frac{1}{q(q-1)} \asymp 1/(p \log p)$  with  $p \asymp \log N$ ; and  $\mathfrak{S}(N) \gg 1$ .  $\square$

*Proof of Proposition 6.* Immediate from Theorem 3,  $|E_3(\alpha)| \leq B(N)$ , and  $|C(N)| \leq \mathfrak{A}(N)N(1 + o(1))$  as above:  $\tilde{r}(N) \geq \mathfrak{S}(N)(1 - \mathfrak{A}(N))N(1 + o(1)) - B(N) + O_A(N(\log N)^{-A})$ .  $\square$

**Measurement 17** (the threshold, and the margin). Over even  $N \leq 1.6 \cdot 10^7$  the threshold constant  $\mathfrak{S}(N)(1 - \mathfrak{A}(N))$  has median 0.333459 and 0.1-percentile 0.119639; its minimum is 0.060890, attained at  $N = 9\,699\,690 = 2 \cdot 3 \cdots 19$ , the largest primorial in range, and the ten smallest values are all primorial-like. Multiplied by  $\log N \log \log N$  the minimum over  $N \geq 10^5$  is 2.482019, attained at  $N = 510\,510$ ; over  $N \geq 10^3$  it is 1.737799 at  $N = 2310$ ; over  $N \geq 16$  it is 0.810202 at  $N = 30$ . Every argmin is a primorial, which is the mechanism of Proposition 5 made visible; the constant  $c$  there is therefore stated as a constant and not as 1. The step that produces the margin is exact to five decimals:  $U(N)/(\mathfrak{A}(N)N)$  has mean 0.999996 with standard deviation 0.000170 over the top octave. Checked against the truth, no even  $N \in [10^5, 1.6 \cdot 10^7]$  has  $\tilde{r}(N) < \mathfrak{S}(N)(1 - \mathfrak{A}(N))N$ ; the slack between them has minimum 3.7038, median 4.2902 and maximum 95.0889, the maximum at  $N = 9\,699\,690$  — so the margin is thinnest exactly where the Goldbach count is largest (`lab_onesided_margin.py`).

**Measurement 18** (what the weakening relocates). At  $\theta' = 0.56$  and  $N = 2 \cdot 10^5$  through  $3.2 \cdot 10^6$ , the quantity  $B(N)$  of (8) satisfies

$$B(N)/N = 0.8086, 0.7395, 0.7303, 0.6547, 0.5916$$

against a threshold of  $0.3745N$ , so the ratio  $B(N)/(\mathfrak{S}(N)(1 - \mathfrak{A}(N))N)$  falls 2.159, 1.975, 1.950, 1.748, 1.580. The object Proposition 6 constrains is therefore already within a factor 1.6 of the threshold at the top of the accessible range, and falling; but a constant-factor bound valid for *all* large  $N$  is what is needed, and no computation supplies that (`lab_onesided_demand.py`).

Proposition 5 does not reopen the divisor switch. The no-go of [7] costs a factor  $\exp(c_1 \sqrt{\frac{1}{2} \log N})$ , which exceeds  $\log N \log \log N$  as it exceeds every power of  $\log N$ ; so the weaker threshold is still out of reach of that design space.

## 7 The correction $\Delta$ to equation (18)

We record a defect in [2], together with its repair. The scope of the report has to be stated: we have worked from the arXiv version, and have checked the passage below in **both** v1 (May 2020)

and v2 (August 2022), where it appears identically. We have not seen the version published in the Springer volume and make no claim about it.

Huang–Li define

$$S_2(\alpha) = \sum_{n < N} \Lambda(n) \mu^2(N - n) \tilde{\Lambda}_\alpha(N - n), \quad \tilde{\Lambda}_\alpha(u) = \sum_{d|u, d > \alpha} \mu(d) \log(1/d),$$

and substitute  $k = u/d$ , so that the constraint  $d > \alpha$  becomes

$$k < \frac{N - n}{\alpha}, \tag{23}$$

an  $n$ -dependent bound. Their displayed equation (18) reads

$$S_2(\alpha) = \sum_{k < \frac{N-1}{\alpha}} \mu(k) \sum_{\substack{n < N \\ n \equiv N \pmod{k}}} \Lambda(n) \mu(N - n) \log\left(\frac{k}{N - n}\right),$$

in which (23) has been replaced by the  $n$ -free bound  $k < (N - 1)/\alpha$ . The two differ. Writing  $N - n = mk$ , the condition (23) is  $m > \alpha$ , so the right-hand side of (18) contains, in addition to  $S_2(\alpha)$ , exactly the terms with  $m \leq \alpha$ . On those terms  $\mu(k)\mu(N - n) = \mu(m)\mu^2(k)\mathbf{1}_{(k,m)=1}$  and  $\log(k/(N - n)) = -\log m$ , so

$$S_2(\alpha) = \text{RHS of (18)} + \Delta, \quad \Delta = \sum_{2 \leq m \leq \alpha} \mu(m) \log m \sum_{\substack{k < (N-1)/\alpha \\ (k,m)=1}} \mu^2(k) \Lambda(N - mk). \tag{24}$$

(The term  $m = 1$  drops out since  $\log 1 = 0$ .) The trivial bound is  $\Delta \ll N(\log N)^2$ , which exceeds the target  $O(N(\log N)^{-A})$ , so  $\Delta$  is not negligible and needs its own treatment.

It does close, by the machinery already used above and under hypotheses Huang–Li already assume. Indeed  $\Delta$  has exactly the shape of the residual (16): the Möbius factor sits on the short variable  $m \leq \alpha$ , and the long variable  $k$  carries only  $\mu^2 \geq 0$ . Since  $\mu(m) \neq 0$  on the range of summation, the two conditions on  $k$  collapse into one,

$$\mu^2(k) \mathbf{1}_{(k,m)=1} = \mu^2(mk), \tag{25}$$

and expanding  $\mu^2(mk) = \sum_{b^2|mk} \mu(b)$  with the truncation  $b \leq B := (\log N)^{A+4}$  — the same one Huang–Li use in their §3.1 — reduces  $\Delta$  to sums of  $\Lambda$  over progressions to moduli  $[m, b^2] \leq \alpha B^2 = \alpha(\log N)^{2A+8}$ , plus a main term  $\mathfrak{A}(N) \sum_{m \leq \alpha} \mu(m) \lambda(m) (\log m) \mathcal{T}_m/m$  which is  $O(Ne^{-c\sqrt{\log N}})$  by Lemma 12 and partial summation. The level is then exactly the one their hypothesis supplies, with no enlargement of  $A$ . Hence:

- In the Corollary-1 regime,  $\alpha = N^{1-\theta'} < N^{1/2}$  and Bombieri–Vinogradov closes  $\Delta$  *unconditionally*.
- In the general Theorem-1 regime,  $\alpha \asymp N^\theta$  and  $\Delta$  closes under the hypothesis  $EH(N^\theta(\log N)^{2A+8})$ , which is assumed there.

So [2]’s Theorem 1 and their Corollary 1 stand as stated; what is missing is the treatment of  $\Delta$ .

## 8 Relation to the literature

This section is a placement, not a survey, and it is deliberately conservative about what is offered here as new.

**Theorem 1.** The mechanism is standard. After the divisor switch the Möbius on the long variable squares away and the surviving Möbius sits on a variable of length at most  $N^{1/2-\delta}$ , which is the classical configuration in which Bombieri–Vinogradov applies. Every ingredient (Bombieri–Vinogradov [6]; the Goldston–Yıldırım estimate [1] as used in [2]; the elementary density identity of Lemma 11) is off the shelf. The statement is offered as a *lemma about the Huang–Li reduction*, not as a contribution to the theory of Bombieri–Vinogradov; its content is that this particular consumption of  $EH_\mu$  is unnecessary, and what difficulty it has lies in the bookkeeping — in particular that the condition  $(k, N) = 1$  must be discarded by the degeneracy argument (Lemma 8) rather than expanded over  $e \mid N$ , and that main terms may be assigned only to classes with  $(q, N) = 1$ , the second of which shifts the density by exactly  $N/\varphi(N)$  if missed (Remark 9).

**Theorem 3.** We are not aware of the identity in the literature. Its ingredients are again classical; what it does is locate, exactly, where the Goldbach content of the Huang–Li reduction sits — in the weight  $\log k$ , by  $\mu * \log = \Lambda$ .

**Beyond the square-root barrier.** Since [2] appeared, Lichtman [4] has shown that the primes have level of distribution  $66/107 \approx 0.617$  using triply well-factorable weights, and has used it to improve upper bounds for Goldbach representations — the first use of a level beyond the square-root barrier on that problem. That work does not collide with the present one: it concerns the primes rather than  $\sum_{n < N} \Lambda(n)\mu(N-n)$ , and it yields upper bounds rather than the asymptotic that [2] targets. It is nevertheless the state of the art against which any claim of the form “beyond  $1/2$ ” must now be measured.

A level of distribution  $3/5$  for the Möbius function with triply well-factorable weights belongs to the earlier part of that line of work, [3]; we cite it here as the object the question below refers to rather than as a result we have checked against the published version. It is *not* in [4], which does not treat the Möbius function at all — the  $3/5$  appearing there is Maynard’s prior record for the *primes*, which [4] improves to  $66/107$ . Since [2] require  $EH_\mu(N^{\theta'})$  for a single  $\theta' > 1/2$ , and since Theorem 3 identifies  $w_k = \log k$  as the weight carrying the Goldbach content, the following question is well posed and, so far as we know, open:

Does the weight that the Huang–Li consumption actually requires lie in the triply well-factorable class for which level  $3/5$  is known for  $\mu$ ?

Two cautions attach to it. First,  $EH_\mu$  as stated in (1) carries  $\max_a |\cdot|$ , whereas well-factorable results bound signed weighted sums  $\sum_q \lambda_q E(x; q, a)$ ; the comparison must therefore be made against what the argument *consumes*, which is a signed sum, and not against the absolute-value form as written. Second, the no-go of [7] closes extraction *by divisor switching*; the technology of [4] is the Deshouillers–Iwaniec spectral large sieve, a different mechanism, and the no-go says nothing about it.

## 9 Summary

- Theorem 1: the Möbius-weighted, fixed-class correlation sum with weight  $w_k = 1$  is  $\ll_A N(\log N)^{-A}$ , unconditionally, for any level  $N^{\theta'}$  with  $\theta' > 1/2$ .
- Corollary 2: hence Huang–Li’s  $E_4$  (their Lemma 4) needs no hypothesis, and the whole  $EH_\mu$  demand of their reduction collapses to the scalar  $E_3$ .
- Theorem 3: that scalar is unconditionally equivalent to Huang–Li’s equation (22), hence already gives binary Goldbach for large even  $N$ , and gives the asymptotic  $\tilde{r}(N) \sim \mathfrak{S}(N)N$  exactly when  $C(N) = o(N)$ .

- Propositions 5 and 6: Goldbach itself needs only a one-sided bound on  $E_3$  at a threshold  $\asymp N$ , equivalently a constant-factor bound on the object  $EH_\mu$  bounds — weaker than the consumed bound by a factor  $(\log N)^A$  for every  $A$ .
- Section 7: equation (18) of [2] omits an  $n$ -dependent constraint; the missing term  $\Delta$  is exhibited and closes under hypotheses already assumed.
- Section 1.4: the net progress toward Goldbach is zero.

## References

- [1] D. Goldston and C. Yıldırım, *Higher correlations of divisor sums related to primes I*, *Integers* **3** (2003), A5.
- [2] J.-J. Huang and H. Li, *On the connection between the Goldbach conjecture and the Elliott–Halberstam conjecture*, arXiv:2005.03811 [math.NT]; v1 (May 2020), v2 (August 2022). Also published in a Springer volume, doi:10.1007/978-3-030-67996-5\_17. References here are to the arXiv version, which we have read, and not to the published version, which we have not.
- [3] J. D. Lichtman, *Primes in arithmetic progressions to large moduli, II: well-factorable estimates*, arXiv:2006.07088 [math.NT], 2020.
- [4] J. D. Lichtman (with an appendix by S. Drappeau), *Primes in arithmetic progressions to large moduli, and Goldbach beyond the square-root barrier*, arXiv:2309.08522 [math.NT], 2023.
- [5] L. Mirsky, *The number of representations of an integer as the sum of a prime and a  $k$ -th power free integer*, *Amer. Math. Monthly* **56** (1949), 17–19.
- [6] H. Montgomery and R. Vaughan, *Multiplicative number theory I: classical theory*, Cambridge University Press, 2007.
- [7] *No weight extracts  $\sum_{n < N} \Lambda(n) \mu(N - n)$ : a no-go for the divisor-switch route, with the exact identities of its design space*, companion paper.
- [8] C.-D. Pan, *A new attempt on Goldbach conjecture*, *Chinese Ann. Math.* **3** (1982), 555–560.

## A Reproducibility

Every numerical figure printed above comes from one of the following scripts; each is standalone, runs under Python with numpy on a laptop, prints its own pass/fail criteria, and writes one result file.

| Script                                 | Result file                             | Supports       |
|----------------------------------------|-----------------------------------------|----------------|
| <code>audit_switch_identity.py</code>  | <code>audit_switch_identity.txt</code>  | (13)           |
| <code>audit_density_identity.py</code> | <code>audit_density_identity.txt</code> | Lemma 11       |
| <code>audit_E3_constant.py</code>      | <code>audit_E3_constant.txt</code>      | (22)           |
| <code>lab_theta_sweep.py</code>        | <code>lab_theta_sweep.txt</code>        | Remark 15      |
| <code>lab_onesided_margin.py</code>    | <code>lab_onesided_margin.txt</code>    | Measurement 17 |
| <code>lab_onesided_demand.py</code>    | <code>lab_onesided_demand.txt</code>    | Measurement 18 |

No computation is used as a premise of any proof. The computations verify exact identities ((13), Lemma 11), fix the sign of a constant that the analysis determines ((22)), or measure a quantity whose size is stated only over the range measured.