Study of investigational REGN5093-M114 in patients with advanced **non small-cell lung cell cancer**



Primary Objective: evaluate the safety, tolerability, and pharmacokinetics of a METxMET bispecific antibody-drug conjugate in patients with MET-overexpressing NSCLC



PATIENTS WITH ADVANCED NSCLC (N≈83)

OPEN LABEL INTERVENTION

Phase 1

Dose escalation: REGN5093-M114 (IV)

Phase 2

Dose expansion: REGN5093-M114 (IV)

Primary Endpoints

Phase 1: Safety and tolerability

Phase 1: Pharmacokinetics

Phase 2: Objective response rate

Secondary Endpoints

Phase 1: Objective response rate

Duration of response Disease control rate Time to response Progression-free survival

Overall survival

Phase 2: Safety and tolerability
Phase 2: Pharmacokinetics



FIND OUT MORE

Scan here to find out more about this study at https://clinicaltrials.gov/ct2/show/NCT04982224

This information is intended for investigators interested in open clinical trials.

REGN5093-M114 is an investigational agent and has not been evaluated by any regulatory authority.

IV, intravenous; N, number of patients; NSCLC, non-small cell lung cancer; MET, mesenchymal epithelial transition.



REGN5093-M114

An Investigational METxMET Bispecific Antibody-Drug Conjugate

Designed to target two distinct epitopes on the MET receptor to deliver a cytotoxic drug to MET-overexpressing tumor cells¹

SELECTED INCLUSION CRITERIA^a



Histologically confirmed NSCLC that is advanced and for which there are no approved therapies available



Newly obtained biopsy tissue availability



Expansion only:

At least one lesion that is measurable by RECIST v1.1



MET overexpression by central IHC analysis



ECOG PS 0 or 1



Adequate organ and bone marrow function

SELECTED EXCLUSION CRITERIA®



Prior treatment with an approved or investigational systemic therapy or investigational device within 14 days of first study drug administration



Unresolved previous acute toxicity from prior therapy



Received radiation therapy or major surgery within 14 days of first administration of study drug



Secondary malignancy that is progressing or requires active treatment



Untreated or active primary brain tumor, CNS metastases, leptomeningeal disease, or spinal cord compression



Uncontrolled infection with HIV, hepatitis B, or hepatitis C infection



Encephalitis, meningitis, organic brain disease (e.g. Parkinson's disease) or uncontrolled seizures in the year prior to first dose of study therapy



For more information, visit <u>www.clinicaltrials.gov</u> or please call [+353 (0)61 533 400 or 844 REGN-MID]. NCT04982224

https://clinicaltrials.gov/ct2/show/NCT04982224

This information is intended for investigators interested in open clinical trials. REGN5093-M114 is an investigational agent and has not been evaluated by any regulatory authority.

Inclusion/exclusion criteria include a summary of selected criteria. Please review the complete study design on clinicaltrials.gov for complete details.

CNS, central nervous system; ECOG PS, Eastern Cooperative Oncology Group performance status; HIV, human immunodeficiency virus; IHC, immunohistochemistry; MET, mesenchymal epithelial transition; NSCLC, non small-cell lung cancer; RECIST, Response Evaluation Criteria in Solid Tumors.

Current per clinicaltrials.gov as of November 29, 2023.

1. Regeneron Investor Event ESMO 2021. Available at: https://investor.regeneron.com/static-files/95baba22-304d-484f-a02c-a95f3043f58c. Accessed July 13, 2023.

REGN5093-M114-EM-0003 v2.0 January 2024.

©2024 Regeneron Pharmaceuticals, Inc. All rights reserved.

