

Study of investigational odronextamab in patients with B-cell non-Hodgkin lymphoma



Primary Objective: evaluate the safety and efficacy of odronextamab in patients with relapsed or refractory B-NHL



PATIENTS WITH B-NHL (N~512)

OPEN LABEL INTERVENTION

Odronextamab (IV)

FL (Grade 1-3a)
Cohort

DLBCL
Cohort

MCL
Cohort

MZL
Cohort

Other B-NHL
Cohort^a

Primary Endpoint

Objective response rate^{b,c}

Secondary Endpoints

Objective response rate^{b,d}

Complete response rate^{b,c,d}

Progression-free survival^{b,c,d}

Overall survival

Duration of response^{b,c,d}

Disease control rate^{b,c,d}

Patient reported outcomes

Pharmacokinetics

Immunogenicity

Safety and tolerability



FIND OUT MORE

Scan here to find out more about this study at <https://classic.clinicaltrials.gov/ct2/show/NCT03888105>

This information is intended for investigators interested in open clinical trials.

Odronextamab is an investigational agent and has not been evaluated by any regulatory authority.

^aPatients with B-NHL other than FL grade 1-3a, DLBCL, MCL, or MZL that has relapsed after or is refractory to at least 2 prior lines of systemic therapy as defined in the protocol. Patients with a current diagnosis of mixed histology of B-NHL with an aggressive component (such as concurrent FL and DLBCL) will be allowed.

^bAccording to the Lugano Classification of response in malignant lymphoma (Cheson BD, et al. *J Clin Oncol* 2014;32:3059-3067).

^cBased on Independent Central Review (ICR). ^dBased on investigator assessment.

B-NHL, B-cell non-Hodgkin lymphoma; DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; IV, intravenous; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma; N, number of patients.

ODRONEXTAMAB

An Investigational CD20xCD3 Bispecific Antibody

Simultaneously engages CD20 on malignant B cells and CD3 on cytotoxic T cells¹

SELECTED INCLUSION CRITERIA^a



FL grade 1-3a cohort: Central histopathologic confirmation of the FL grade 1 to 3a diagnosis



DLBCL and MZL cohorts: patients that have relapsed after or are refractory to at least 2 prior lines of systemic therapy



MCL after BTK inhibitor therapy cohort: New enrollment is paused until further notice



ECOG PS 0 or 1



Adequate bone marrow, hepatic, and renal function



Measurable disease on cross sectional imaging documented by diagnostic CT or MRI

SELECTED EXCLUSION CRITERIA^a



Primary CNS lymphoma, known involvement by non-primary CNS NHL, or history of CNS movement disorder



Prior treatment with a CD20xCD3 bispecific antibody therapy



Prior treatment with any CAR T-cell therapy



History of allogeneic stem cell transplantation



Treatment with any systemic anti-lymphoma therapy within 5 half-lives or within 28 days prior to administration of study drug



Uncontrolled infection with HIV, hepatitis B or hepatitis C infection, or cytomegalovirus infection



For more information, visit www.clinicaltrials.gov or please call [+353 (0)61 533 400 or 844 REGN-MID].
NCT03888105
<https://classic.clinicaltrials.gov/ct2/show/NCT03888105>

**This information is intended for investigators interested in open clinical trials.
Odronextamab is an investigational agent and has not been evaluated by any regulatory authority.**

^aInclusion/exclusion criteria include a summary of selected criteria. Please review the complete study design on clinicaltrials.gov for complete details.

B-NHL, B-cell non-Hodgkin lymphoma; CAR, chimeric antigen receptor; CD, cluster of differentiation; CNS, central nervous system; CT, computed tomography; ECOG PS, Eastern Cooperative Oncology Group performance status; HIV, human immunodeficiency virus; MRI, magnetic resonance imaging.

1. Smith EJ et al. *Sci Rep*. 2015;5:17943.

Current per clinicaltrials.gov as of April 20, 2023.

Odronextamab-EM-0009 v2.0 February 2024.

©2024 Regeneron Pharmaceuticals, Inc. All rights reserved.