

Study of investigational livoseltamab in patients with relapsed or refractory multiple myeloma



Primary Objective: evaluate the safety and efficacy, and identify the recommended phase 2 dose, of livoseltamab in patients with relapsed or refractory (R/R) multiple myeloma



PATIENTS WITH R/R MM (N≈309)

OPEN LABEL INTERVENTION

Phase 1

Dose escalation: Livoseltamab (IV)



Phase 2

Dose expansion: Livoseltamab (IV)

Primary Endpoints

Phase 1: Safety, tolerability and DLT

Phase 2: Overall response rate^{a,b}

Secondary Endpoints

Phase 1/2: Objective response rate^a

Duration of response^a

Progression-free survival^a

MRD-negative status^a

Overall survival^a

Pharmacokinetics

Immunogenicity

Phase 2: HRQoL, Safety, and tolerability



FIND OUT MORE

Scan here to find out more about this study at <https://clinicaltrials.gov/ct2/show/NCT03761108>

This information is intended for investigators interested in open clinical trials.

Livoseltamab is an investigational agent and has not been evaluated by any regulatory authority.

^aUsing the International Myeloma Working Group (IMWG) response criteria. ^bBased on Blinded Independent Central Review (BICR) in phase 2 portion. ^cBased on investigator assessment. DLT, dose-limiting toxicity; HRQoL, health-related quality of life; IV, intravenous; MM, multiple myeloma; MRD, minimal residual disease; N, number of patients.

LINVOSELTAMAB

An Investigational BCMAxCD3 Bispecific Antibody

Designed to bridge BCMA on MM cells with CD3 on T cells to facilitate local T-cell activation and cytotoxicity¹

SELECTED INCLUSION CRITERIA^a



ECOG PS 0 or 1



Measurable disease according to IMWG consensus criteria



Phase 1: Progression on or after at least 3 lines of therapy, or intolerance to therapy, including a PI, an IMiD, and an anti-CD38 antibody, OR progression on or after an anti-CD38 antibody^b and have disease that is double refractory^c to a PI and an IMiD, or intolerance to therapy



Phase 2: Progression on or after at least 3 prior lines of therapy including a PI, IMiD, and anti-CD38 antibody. Patients must be triple-refractory, defined as being refractory^d to prior treatment with at least one anti-CD38 antibody, PI, and IMiD

SELECTED EXCLUSION CRITERIA^a



Diagnosis of plasma cell leukemia, primary light-chain amyloidosis, Waldenström macroglobulinemia, or POEMS syndrome



Known MM brain lesions or meningeal involvement



Cardiac ejection fraction <40%



Prior treatment with BCMA-directed immunotherapies excluding BCMA antibody-drug conjugates^e



History of allogeneic stem cell transplantation, or autologous stem cell transplantation within 12 weeks of the start of study drug regimen



For more information, visit www.clinicaltrials.gov or please call +353 (0)61 533 400.
NCT03761108
<https://clinicaltrials.gov/ct2/show/NCT03761108>

This information is intended for investigators interested in open clinical trials.

Linvoseltamab is an investigational agent and has not been evaluated by any regulatory authority.

^aInclusion/exclusion criteria include a summary of selected criteria. Please review the complete study design on clinicaltrials.gov for complete details. ^bThe anti-CD38 antibody may have been administered alone or in combination with another agent such as a PI; ^cDefined as lack of response or relapse within 60 days of last treatment; ^dDefined as progression during treatment or within 60 days after completion of therapy, or <25% response to therapy; ^eIncludes BCMA bispecific antibodies, bispecific T Cell engagers, and CAR T Cells.

BCMA, B cell maturation antigen; CD, cluster of differentiation; ECOG PS, Eastern Cooperative Oncology Group performance status; IMiD, immunomodulatory imide drug; IMWG, International Myeloma Working Group; MM, multiple myeloma; PI, proteasome inhibitor; POEMS, polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes.

Current per clinicaltrials.gov as of April 5, 2023.

1. DiLillo DJ et al. *Blood Adv*. 2021;5(5):1291–04.

LNVO-EM-0001 October 2023.

©2023 Regeneron Pharmaceuticals, Inc. All rights reserved.