Back

JGR Space Physics

Method

HARP-FiLM: Global Ionospheric VTEC Forecasting With Solar–Geomagnetic Driver Conditioning and Dual-Baseline Residuals

J. Chen, C. Yang, L. Han

First published: 12 October 2026

https://doi.org/10.1029/2026JA035091

VIEW METRICS

EPUB PDF

Get access to the full version of this article. View access options below.

Institutional Login

Access through your institution

Log in with AGU

Log in with your society credentials

Log in

Log in to Wiley Online Library

Back

Log in with CONNECT

One account for all your research.

Wiley Online Library is part of the CONNECT Network.

Purchase Instant Access

48-Hour online access $16.00

Details

Online-only access $27.00

Details

PDF download and online access $62.00

Details

Check out

Abstract

Accurate 24 hr global ionospheric vertical total electron content (VTEC) forecasting is difficult because diurnal recurrence, solar-cycle variability, and geomagnetic disturbances act on different time scales. We present HARP-FiLM (Horizon-Aware Adaptive Residual Prediction with Feature-wise Linear Modulation), a solar–geomagnetic-driver-conditioned framework that combines multi-scale spatial encoding, Transformer temporal modeling, structured feature modulation, and adaptive dual-baseline residual prediction. Using global ionosphere maps from the Center for Orbit Determination in Europe (CODE) and OMNI2 drivers, HARP-FiLM achieves root-mean-square errors (RMSEs) of 1.356 TECU in 2020 and 4.956 TECU in 2023. Relative to the International Reference Ionosphere 2016 (IRI-2016), RMSE is reduced by 57.7% and 56.3%, while blocked experiments yield reductions of 62.9% and 47.4% in

reported skill is not primarily produced by direct fitting to temporally interpolated targets. External cross-product evaluation against the Institute for Space–Earth Environmental Research (ISEE) ground-based GNSS-derived TEC product also shows lower errors than IRI-2016 in all four evaluated years, although the margin narrows in 2023. Regional, seasonal, geomagnetic-activity-stratified, and rapid-forcing analyses show the strongest gains under quiet to moderately disturbed conditions, whereas intense and super-storm periods remain challenging. These results support structured driver conditioning and adaptive temporal priors for global VTEC forecasting while highlighting limitations associated with severe disturbances and gridded target products.

Plain Language Summary

The ionosphere affects satellite navigation and radio signals because its electron content changes with local time, season, solar activity, and geomagnetic disturbances. These changes are especially difficult to forecast during strong space-weather events. We developed a global one-day forecasting method that combines recent ionospheric maps with information about solar and geomagnetic conditions. The method also uses recent observations and the ionospheric state at the same time on the previous day as adaptive reference patterns. Tests across years with different solar activity show lower forecast errors than a standard empirical ionosphere model, simple persistence methods, and several representative learning models. A separate comparison with a gridded electron-content product derived from ground-based satellite-navigation observations shows that the improvement is not limited to agreement with the maps used for training. The largest benefits occur under quiet to moderately disturbed conditions, while the strongest geomagnetic storms remain difficult to predict. These results show that structured use of space-weather information can improve global ionospheric forecasting while also identifying where additional storm-time observations and modeling are still needed.

Conflict of Interest

The authors declare no conflicts of interest relevant to this study.

The CODE Global Ionosphere Map (GIM) data used as the target VTEC data set in this study are from the CODE final product series for the IGS, provided by the Center for Orbit Determination in Europe (CODE) at the Astronomical Institute, University of Bern (Dach et al., **2026**). The hourly solar-wind and geomagnetic driver data used in this study are available from the NASA OMNIWeb OMNI2 database provided by the NASA Goddard Space Flight Center Space Physics Data Facility via **https://omniweb.gsfc.nasa.gov** (NASA Goddard Space Flight Center, **2026**). Version 1.0.1 of the HARP-FiLM source code used to train, evaluate, and reproduce the forecasting model is preserved in Zenodo via **https://doi.org/10.5281/zenodo.20406934**, licensed under the MIT License, and developed openly at **https://github.com/research1011/HARP-FiLM** (Chen, Yang, & Han, **2026**). The ISEE global GNSS-derived absolute TEC AGRID2 NetCDF products used for the external cross-product evaluation are publicly available from the Institute for Space–Earth Environmental Research (ISEE), Nagoya University, through the ISEE GNSS-TEC database (Shinbori et al., **2020**, **2021**; Zossi et al., **2026**). The AGRID2 data used in this study were downloaded from **https://stdb2.isee.nagoya-u.ac.jp/GPS/shinbori/AGRID2/nc/**.

References

Ahoua, S. M., Habarulema, J. B., Obrou, O. K., Cilliers, P. J., & Zaka, Z. K. (2018). Evaluation of the NeQuick model performance under different geomagnetic conditions over South Africa during the ascending phase of the solar cycle (2009–2012). *Annales Geophysicae*, **36**(5), 1161–1170. **https://doi.org/10.5194/angeo-36-1161-2018**

ADS | Web of Science® | Google Scholar

Aol, S., Buchert, S., & Jurua, E. (2020). Ionospheric irregularities and scintillations: A direct comparison of in situ density observations with ground-based L-band receivers. *Earth Planets and Space*, **72**(1), 164. **https://doi.org/10.1186/s40623-020-01294-z**

ADS | Web of Science® | Google Scholar

Bilitza, D., Pezzopane, S. M., Truhlik, V., Altadill, D., Reinisch, B. W., & Pignalberi, A. (2022). The international reference ionosphere model: A review and description of an ionospheric benchmark. *Reviews of Geophysics*, **60**(4), e2022RG000792. **https://doi.org/10.1029/2022RG000792**

ADS | Web of Science® | Google Scholar

2022SW003227

ADS | Web of Science® | Google Scholar

Chen, J., Yang, C., & Han, L. (2026). HARP-FiLM: Global ionospheric VTEC forecasting with solar–geomagnetic driver conditioning and dual-baseline residuals (Version 1.0.1) [Software]. Zenodo. https://doi.org/10.5281/zenodo.20406934

Google Scholar

Chen, J., Yang, C., Han, L., & Yang, S. (2026). Impact of solar and geomagnetic driver selection on 24 h-ahead global VTEC prediction in a deep learning framework: A ConvLSTM case study. *Geosciences*, **16**(5), 169. https://doi.org/10.3390/geosciences16050169

Web of Science® | Google Scholar

Chen, J., Zhi, N., Liao, H., Lu, M., & Feng, S. (2022). Global forecasting of ionospheric vertical total electron contents via ConvLSTM with spectrum analysis. *GPS Solutions*, **26**(3), 69. https://doi.org/10.1007/s10291-022-01253-z

Web of Science® | Google Scholar

Dach, R., Schaer, S., Arnold, D., Brockmann, E., Kalarus, M. S., Kobel, C., et al. (2026). CODE final product series for the IGS, Astronomical Institute, University of Bern. Published by. https://doi.org/10.48620/97773

Google Scholar

Danilchuk, E., Yasyukevich, Y., Vesnin, A., Klyusilov, A., & Zhang, B. (2025). Impact of the May 2024 extreme geomagnetic storm on the ionosphere and GNSS positioning. *Remote Sensing*, **17**(9), 1492. https://doi.org/10.3390/rs17091492

ADS | Web of Science® | Google Scholar

Denardini, C. M., Pinheiro, G. A. S., Barbosa Neto, P. F., Nogueira, P. A. B., Carmo, C. S., Resende, L. C. A., et al. (2020). Ionospheric scale index map based on TEC data for space-weather studies and

ADS | Web of Science® | Google Scholar

Fang, T., Fuller-Rowell, T., Yudin, V., Matsuo, T., & Viereck, R. (2018). Quantifying the sources of ionosphere day-to-day variability. *Journal of Geophysical Research: Space Physics*, **123**(11), 9682–9696. https://doi.org/10.1029/2018JA025525

ADS | Web of Science® | Google Scholar

Gao, X., Yao, Y., & Wang, Y. (2025). Effects of different feature representations on global ionospheric TEC model based on spatiotemporal convolutions. *GPS Solutions*, **29**(2), 84. https://doi.org/10.1007/s10291-024-01799-0

Web of Science® | Google Scholar

Han, B., Yu, Q., Wang, J., Chernogor, L. F., & Zheng, Y. (2025). An ionospheric VTEC forecasting model over China based on the kernel ridge regression. *Advances in Space Research*, **76**(1), 353–370. https://doi.org/10.1016/j.asr.2025.04.049

ADS | Web of Science® | Google Scholar

Huba, J. D., Maute, A., & Crowley, G. (2017). SAMI3_ICON: Model of the ionosphere/plasmasphere system. *Space Science Reviews*, **212**(1), 731–742. https://doi.org/10.1007/s11214-017-0415-z

CAS | ADS | PubMed | Web of Science® | Google Scholar

Kim, J., & Kwak, Y. (2025). Validating the IRI-2020 model for ionospheric storms over the North-East Asian sector induced by extreme geomagnetic storms. *Advances in Space Research*, **75**(5), 4347–4369. https://doi.org/10.1016/j.asr.2024.07.032

ADS | Web of Science® | Google Scholar

Li, L., Liu, Y., Zhou, H., Yang, K., Yan, H., & Li, J. (2025). Ionformer: A data-driven deep learning baseline for global ionospheric TEC forecasting. *IEEE Transactions on Geoscience and Remote Sensing*, **63**, 1–12. https://doi.org/10.1109/TGRS.2025.3542182

Lin, Y., Fang, H., Duan, D., Huang, H., Xiao, C., & Ren, G. (2024). Enhanced forecasting of global ionospheric vertical total electron content maps using deep learning methods. *Atmosphere*, **15**(11), 1319. https://doi.org/10.3390/atmos15111319

ADS | Web of Science® | Google Scholar

Liu, H., Wang, H., Le, H., Yuan, J., Shen, W., Wu, Y., & Chen, Y. (2025). CGAOA-STRA-BiConvLSTM: An automated deep learning framework for global TEC map prediction. *GPS Solutions*, **29**(1), 55. https://doi.org/10.1007/s10291-025-01814-y

Web of Science® | Google Scholar

Liu, L., Zou, S., Yao, Y., & Wang, Z. (2020). Forecasting global ionospheric TEC using deep learning approach. *Space Weather*, **18**(11), e2020SW002501. https://doi.org/10.1029/2020SW002501

ADS | Web of Science® | Google Scholar

Materassi, M., Alberti, T., Migoya-Orué, Y., Radicella, S. M., & Consolini, G. (2023). Chaos and predictability in ionospheric time series. *Entropy*, **25**(2), 368. https://doi.org/10.3390/e25020368

ADS | PubMed | Web of Science® | Google Scholar

Mehta, P. M., & Linares, R. (2020). Real-time thermospheric density estimation from satellite position measurements. *Journal of Guidance, Control, and Dynamics*, **43**(9), 1656–1670. https://doi.org/10.2514/1.G004793

ADS | Web of Science® | Google Scholar

NASA Goddard Space Flight Center. (2026). OMNIWeb: OMNI2 hourly solar wind and geomagnetic data [Dataset]. NASA Space Physics Data Facility. Retrieved from https://omniweb.gsfc.nasa.gov

Google Scholar

10.1016/j.jastp.2008.01.015

ADS | Web of Science® | Google Scholar

Paziewski, J., Høeg, P., Sieradzki, R., Jin, Y., Jarmołowski, W., Hoque, M. M., et al. (2022). The implications of ionospheric disturbances for precise GNSS positioning in Greenland. *Journal of Space Weather and Space Climate*, **12**, 33. https://doi.org/10.1051/swsc/2022029

ADS | Web of Science® | Google Scholar

Ren, X., Yang, P., Liu, H., Chen, J., & Liu, W. (2022). Deep learning for global ionospheric TEC forecasting: Different approaches and validation. *Space Weather*, **20**(5), e2021SW003011. https://doi.org/10.1029/2021SW003011

ADS | Web of Science® | Google Scholar

Ren, X., Zhao, B., Ren, Z., Wang, Y., & Xiong, B. (2024). Deep learning-based prediction of global ionospheric TEC during storm periods: Mixed CNN-BiLSTM method. *Space Weather*, **22**(7), e2024SW003877. https://doi.org/10.1029/2024SW003877

ADS | Web of Science® | Google Scholar

Schunk, R. W., Scherliess, L., Sojka, J. J., Thompson, D. C., Anderson, D. N., Codrescu, M., et al. (2004). Global assimilation of ionospheric measurements (GAIM). *Radio Science*, **39**(1), RS1S02. https://doi.org/10.1029/2002RS002794

Web of Science® | Google Scholar

Shinbori, A., Otsuka, Y., Sori, T., Tsugawa, T., & Nishioka, M. (2020). Temporal and spatial variations of total electron content enhancements during a geomagnetic storm on 27 and 28 September 2017. *Journal of Geophysical Research: Space Physics*, **125**(7), e2019JA026873. https://doi.org/10.1029/2019JA026873

ADS | Web of Science® | Google Scholar

Relationship between the locations of the ... trough and plasmapause using ... and Arase satellite observation data. *Journal of Geophysical Research: Space Physics*, **126**(5), e2020JA028943. https://doi.org/10.1029/2020JA028943

ADS | Web of Science® | Google Scholar

Sivavaraprasad, G., & Ratnam, D. V. (2017). Short-term forecasting of ionospheric total electron content over a low-latitude global navigation satellite system station. *IET Radar, Sonar & Navigation*, **11**(8), 1309–1320. https://doi.org/10.1049/iet-rsn.2017.0011

Web of Science® | Google Scholar

Tang, J., Ding, M., Yang, D., Fan, C., Khonsari, N., & Mao, W. (2024). Different data-driven prediction of global ionospheric TEC using deep learning methods. *International Journal of Applied Earth Observation and Geoinformation*, **130**, 103889. https://doi.org/10.1016/j.jag.2024.103889

Web of Science® | Google Scholar

Tsurutani, B. T., Mannucci, A. J., Iijima, B. A., Abdu, M. A., Sobral, J. H. A., Gonzalez, W. D., et al. (2004). Global dayside ionospheric uplift and enhancement associated with interplanetary electric fields. *Journal of Geophysical Research*, **109**(A8), A08302. https://doi.org/10.1029/2003JA010342

Web of Science® | Google Scholar

Yu, F., Yuan, H., Chen, S., Luo, R., & Luo, H. (2024). Graph-enabled spatio-temporal transformer for ionospheric prediction. *GPS Solutions*, **28**(4), 203. https://doi.org/10.1007/s10291-024-01734-3

Web of Science® | Google Scholar

Yuan, Y., Xia, G., Zhang, X., & Zhou, C. (2023). Synthesis-style auto-correlation-based transformer: A learner on ionospheric TEC series forecasting. *Space Weather*, **21**(10), e2023SW003472. https://doi.org/10.1029/2023SW003472

ADS | Web of Science® | Google Scholar

Zhang, Q., & Zhao, Q. (2019). Analysis of the data processing strategies of spherical harmonic expansion model on global ionosphere mapping for moderate solar activity. *Advances in Space*

ADS | Web of Science® | Google Scholar

Zhang, R., Li, H., Shen, Y., Yang, J., Li, W., Zhao, D., & Hu, A. (2025). Deep learning applications in ionospheric modeling: Progress, challenges, and opportunities. *Remote Sensing*, **17**(1), 124. https://doi.org/10.3390/rs17010124

ADS | Web of Science® | Google Scholar

Zossi, B. S., Medina, F. D., Duran, T., Zamora, D. J., & Elias, A. G. (2026). Global long-term trends in ionospheric TEC from GNSS observations with local-time dependence. *Geophysical Research Letters*, **53**(6), e2026GL121731. https://doi.org/10.1029/2026GL121731

ADS | Web of Science® | Google Scholar

Download PDF

Back to Top

AGU ADVANCING EARTH AND SPACE SCIENCES

AGU PUBLICATIONS

AGU.ORG

AGU MEMBERSHIP

RESOURCES

PUBLICATION INFO

© 2026 American Geophysical Union

ABOUT WILEY ONLINE LIBRARY

Privacy Policy
Terms of Use
About Cookies
Manage Cookies
Accessibility
Wiley Research DE&I Statement and Publishing Policies

HELP & SUPPORT

Contact Us
Training and Support
DMCA & Reporting Piracy
Sitemap

OPPORTUNITIES

Subscription Agents
Advertisers & Corporate Partners

CONNECT WITH WILEY

The Wiley Network
Wiley Press Room

Copyright © 1999-2026 John Wiley & Sons, Inc or related companies. All rights reserved, including rights for text and data mining and training of artificial intelligence technologies or similar technologies.