Semantic Collapse Geometry and Nested Uplifts Inevitability: A Geometric–Dynamic Path Toward the Riemann Hypothesis

https://chatgpt.com/share/68f02e82-a9d4-8010-9215-0872077567cf

Abstract

This article proposes a geometric–dynamic research path toward the Riemann Hypothesis (RH) by unifying two complementary frameworks: Semantic Collapse Geometry (SCG) and Nested Uplifts Inevitability (INU). SCG recasts prime-gaps arithmetic into a curvature field on a discrete trajectory, from which an intrinsic "collapse Laplacian" emerges whose spectral modes encode the imaginary parts of ζ-zeroes. INU supplies the missing temporal dimension: a sequential-evidence and small-gain mechanism whose whitening threshold aligns with the zeta critical line. Within this union, RH is reinterpreted as an equilibrium law: the critical line is the unique curvature-balance locus where the collapse field attains minimal energy subject to a dynamical whitening constraint. The paper develops the objects and equivalences required for this transposition, outlines verifiable implications, and clarifies how this route differs from and complements spectral and physical approaches in the Hilbert–Pólya, Berry–Keating, and noncommutative geometry lines (Berry and Keating 1999; Connes 1999). The emphasis is on a closed-loop structure linking (i) a discrete curvature extracted from prime gaps, (ii) a self-adjoint collapse generator, and (iii) an INU whitening condition interpreted directly on zeta's error processes. The result is a testable, geometry-driven path that transforms RH from a purely analytic conjecture into a stability claim about a coupled curvature–evidence system.

---

1. Introduction

The Riemann zeta function encodes deep information about the primes through its analytic continuation and nontrivial zeroes. The Riemann Hypothesis asserts that all nontrivial zeroes of $\zeta(s)$ lie on the critical line $\mathrm{Re}(s) = 1/2$. In the conventional analytic setting, $\zeta(s)$ is introduced for $\mathrm{Re}(s) > 1$ by the Dirichlet series

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} \qquad (1.1)$$

and continued meromorphically elsewhere. While modern number theory connects ζ to automorphic forms, L-functions, and random matrix predictions for the zero statistics (Montgomery 1973; Odlyzko 1987; Berry and Keating 1999), the central obstacle remains: we lack a canonical geometric or dynamical object whose intrinsic symmetries force the zeroes onto $\mathrm{Re}(s) = 1/2$.

This paper develops a new route by unifying two ideas:

1. Semantic Collapse Geometry (SCG).
   SCG treats the prime sequence as a discrete trajectory whose "local shape" is measured by a curvature extracted from adjacent prime gaps. Intuitively, irregularities of the prime gaps produce signed curvature fluctuations. SCG posits that the principal oscillatory modes of this curvature field correspond to zeta's nontrivial zeroes. The "critical line" is reinterpreted as a curvature-balance locus: a stationarity condition of a natural energy functional on the discrete trajectory.

2. Nested Uplifts Inevitability (INU).
   INU is a sequential-evidence and small-gain framework for open systems with thresholds. It formalizes when accumulated evidence crosses a critical value and subsequently whitens (i.e., residual fluctuations become decorrelated and scale-stable). When mapped to zeta's error processes, the INU whitening threshold corresponds to Re(s) = 1/2. Deviations from the line manifest as heavy-tailed residue and log–log rescaling effects; whitening at the threshold is the signature of alignment with the critical line.

The synthesis is a closed loop: discrete curvature (SCG) generates a canonical self-adjoint operator (the collapse Laplacian) whose spectrum must be real; simultaneously, the INU whitening threshold selects the same equilibrium as the unique stable operating point. Thus RH becomes equivalent to the coincidence of (i) geometric energy minimization under curvature balance and (ii) dynamical whitening at the INU threshold.

This approach is conceptually close to—but distinct from—Hilbert–Pólya proposals, which seek a self-adjoint operator with eigenvalues matching the imaginary parts of zeta zeroes (Pólya 1926; Hilbert 1914). It also contrasts with Hamiltonian-inspired models (Berry and Keating 1999) and the spectral interpretations arising in noncommutative geometry (Connes 1999). The difference is twofold: SCG provides a constructive geometric origin for the operator (the collapse Laplacian), and INU supplies a statistically verifiable dynamical criterion (whitening) that fixes the relevant spectral locus.

The remainder of the article proceeds as follows. Section 2 defines the SCG curvature from prime gaps and formulates the semantic zeta transform. Section 3 constructs the collapse Laplacian and states the spectral equivalence principle. Section 4 translates INU's evidence thresholds into a whitening condition on zeta residues. Section 5 closes the loop, showing how the geometric and dynamical conditions pick out a unique equilibrium—precisely the critical line. Section 6 sketches empirical pathways. Section 7 discusses relations to existing programs and implications. Section 8 concludes.

---

2. Semantic Collapse Geometry: From Prime Gaps to Curvature and Modes

2.1 Discrete trajectory and curvature

Let p_n denote the n-th prime and define the prime gaps $g_n = p_{n+1} - p_n$. SCG models the prime sequence as a discrete trajectory whose local irregularity is encoded by a curvature observable κ_n computed from consecutive gaps. A symmetric, scale-free choice that responds to relative gap variation is

$$\kappa_n = 2\cdot(g_{n+1} - g_n) / (g_{n+1} + g_n) \qquad (2.1)$$

The factor 2 normalizes κ_n so that $\kappa_n \in (-2, 2)$ whenever $g_{n+1}, g_n > 0$. Other normalizations are possible; the essential feature is to capture signed, relative changes in consecutive gaps in a way that is insensitive to uniform rescaling of the sequence.

For technical developments, we consider κ as a signed signal on the index set $n \in \mathbb{N}$. Define partial sums $K(N)$ and energy density $E(N)$:

$$K(N) = \sum_{n\le N} \kappa_n \qquad (2.2)$$

$$E(N) = (1/N)\cdot\sum_{n\le N} \kappa_n^2 \qquad (2.3)$$

$E(N)$ is the empirical curvature energy over the first N steps. The SCG viewpoint treats $E(N)$ as an observable approaching a limiting energy density E_* in an appropriate averaging sense. The

“curvature-balance” hypothesis is that the RH-critical behavior corresponds to a stationary point of an energy functional built from κ and its correlations.

2.2 Semantic zeta transform and oscillatory modes

To connect κ to spectral data reminiscent of zeta zeroes, we introduce a semantic Dirichlet transform of κ. Define

$$Z_{sem}(s) = \sum_{n=1}^{\infty} \kappa_n \cdot n^{-s} \qquad (2.4)$$

for s in the half-plane $\mathrm{Re}(s) > 1 + \varepsilon$ where the series absolutely converges (ε depends on tail behavior of κ_n). The analytic continuation of $Z_{sem}(s)$ can be studied via summability methods or by smoothing κ with compactly supported windows. The guiding principle is that persistent oscillatory structure in κ should manifest as poles/zeroes or sharp features in $Z_{sem}(s)$, analogous to how oscillations in arithmetic functions shape the analytic structure of corresponding Dirichlet series.

A complementary frequency-domain view uses the discrete Fourier transform of a windowed κ:

$$\hat{K}_N(\theta) = \sum_{n=1}^{N} \kappa_n \cdot e^{-i n \theta} \qquad (2.5)$$

Sharp peaks of $\hat{K}_N(\theta)$ (as $N \to \infty$ along subsequences) indicate dominant oscillatory modes of the curvature field. SCG posits a mode–zero correspondence principle:

Dominant modes of κ ↔ ordinates of nontrivial ζ-zeroes. (2.6)

This principle is not asserted as proved here; rather, it is the organizing hypothesis that connects curvature irregularity to zeta spectral data. Section 3 elevates this to an operator-theoretic statement via the collapse Laplacian.

2.3 Curvature balance and the critical line

SCG reframes the critical line $\mathrm{Re}(s) = 1/2$ as a balance locus for curvature modes. Intuitively, if the κ-field can be decomposed into approximate normal modes with phases tied to a spectral parameter t, then “being on the critical line” is the condition that phases align to minimize a curvature energy functional under a neutrality constraint (no persistent bias in signed curvature). A minimal formal statement is the existence of a functional $\mathscr{E}$ on signed sequences satisfying

$$\mathscr{E}[\kappa] = \limsup_{N\to\infty} (1/N)\cdot\sum_{n\le N} \Phi(\kappa_n, \kappa_{n+1}, \ldots) \qquad (2.7)$$

where Φ is convex in κ entries and enforces a neutrality (zero-mean) constraint. The **curvature-balance hypothesis** asserts that the minimizing configurations correspond to phases associated with $\mathrm{Re}(s) = 1/2$. In Section 5, this geometric stationarity is shown to coincide with the INU whitening threshold, producing a unique equilibrium consistent with RH.

2.4 Relation to classical arithmetic signals

Classical transforms often use the Möbius function μ(n) or the von Mangoldt function Λ(n). SCG does not replace these but complements them: κ_n is a high-pass, local-contrast observable on the prime trajectory, while μ and Λ encode multiplicative structure. One may introduce mixed observables—e.g., a curvature-weighted Mangoldt series—to probe cross-correlations:

$$M_{sem}(s) = \sum_{n=1}^{\infty} \kappa_n \cdot \Lambda(n) \cdot n^{-s} \qquad (2.8)$$

Such hybrids can amplify or filter specific oscillations. The core thesis does not depend on any single hybrid, only on the existence of a stable curvature signal whose dominant modes tie to the

ζ spectrum.

---

3. The Collapse Laplacian and the Spectral Equivalence Principle

3.1 Semantic state space and inner product

Let $\mathcal{S}$ be the space of real, square-summable sequences on $\mathbb{N}$ with zero mean, i.e., sequences $x = (x_n)$ with $\sum_{n=1}^{\infty} x_n^2 < \infty$ *and* $\lim_{N\to\infty} (1/N)\cdot\sum_{n\le N} x_n = 0$. Equip $\mathcal{S}$ with the standard ℓ^2 inner product

$$\langle x, y\rangle = \sum_{n=1}^{\infty} x_n y_n \qquad (3.1)$$

SCG treats $\kappa = (\kappa_n)$ as a stochastic–deterministic element of $\mathcal{S}$ (after suitable truncation or tapering). The zero-mean constraint expresses curvature neutrality, a prerequisite for "balance" on the critical line.

3.2 Discrete collapse Laplacian

Define the collapse Laplacian Δ_c on $\mathcal{S}$ by a weighted graph Laplacian that couples consecutive curvature entries with adaptive weights $w_n > 0$:

$$(\Delta_c x)_n = w_{n-1}(x_n - x_{n-1}) - w_n(x_{n+1} - x_n) \qquad (3.2)$$

with the convention $w_0 = 0$ and appropriate boundary conditions (e.g., $x_0 = 0$). The weights can be chosen as smooth functions of local curvature energy (or of gaps g_n), for example

$$w_n = f(g_n, g_{n+1}) = 2\cdot g_n g_{n+1} / (g_n + g_{n+1}) \qquad (3.3)$$

which is the harmonic mean up to scale; other smooth positive choices are admissible provided $\sum w_n$ diverges and w_n is slowly varying in the sense of bounded relative increments. Under the ℓ^2 inner product, Δ_c with such symmetric edge weights is self-adjoint on its natural domain (Sturm–Liouville type on a line graph).

Self-adjointness (formal): for all compactly supported $x, y \in \mathcal{S}$,

$$\langle \Delta_c x, y\rangle = \langle x, \Delta_c y\rangle \qquad (3.4)$$

The verification is a discrete Green identity. The import is that Δ_c possesses a real spectrum $\mathrm{Spec}(\Delta_c) \subseteq \mathbb{R}$ and a complete set of generalized eigenmodes under mild conditions.

3.3 Semantic wave operator and unitary flow

Introduce the first-order semantic wave generator A_c by

$$A_c = J\,\Delta_c^{1/2} \qquad (3.5)$$

where J is the complex structure on $\mathcal{S} \otimes \mathbb{C}$ ($J^2 = -I$) and $\Delta_c^{1/2}$ is defined via spectral calculus. Then $U_t = \exp(-i\,t\,A_c)$ is a unitary flow on $\mathcal{S} \otimes \mathbb{C}$. The frequencies of the flow are the moduli of the eigenvalues of A_c, i.e., the square-roots of the nonnegative spectrum of Δ_c. Denote the discrete (or resonant) frequency set by $\{t_k\}$.

$$t_k = \sqrt{\lambda_k} \text{ where } \lambda_k \in \mathrm{Spec}(\Delta_c), \quad U_t\,\varphi_k = e^{-i\,t\,t_k}\,\varphi_k \qquad (3.6)$$

The SCG thesis is that the dominant semantic frequencies $\{t_k\}$ match the ordinates of

nontrivial zeta zeroes ρ_k = 1/2 + i t_k, up to a universal scaling fixed by normalization. This is the spectral equivalence principle:

Semantic Spectral Equivalence: There exists a normalization of Δ_c such that the resonant set {t_k} coincides with the multiset of ordinates of nontrivial zeroes of ζ(s). (3.7)

This statement is in the spirit of Hilbert–Pólya (Pólya 1926; Berry and Keating 1999) but here the operator is geometrically constructed from prime-gap curvature rather than postulated.

3.4 Curvature forcing and energy stationarity

Define the curvature energy functional on $\mathcal{S}$:

$\mathscr{E}[x] = \limsup_{N\to\infty} (1/N)\cdot\sum_{n\le N} [\, \alpha\cdot x_n^2 + \beta\cdot(x_{n+1} - x_n)^2 \,]$ (3.8)

with α, β > 0. Critical points of $\mathscr{E}$ under the zero-mean constraint satisfy the Euler–Lagrange equation

$\alpha\cdot x_n - \beta\cdot(x_{n+1} - 2x_n + x_{n-1}) = 0$ (3.9)

which, after inserting the adaptive weights, generalizes to $\Delta_c x = \mu x$ for a Lagrange multiplier μ. Thus $\mathscr{E}$-stationarity selects Δ_c-eigenmodes. The SCG balance claim is:

Curvature-Balance Principle: The physical (statistically stable) curvature configurations lie on Δ_c eigenmodes whose frequencies land on the critical line ordinates t = Im(ρ). (3.10)

In Section 5 we align this geometric selection with the INU whitening threshold, showing both pick the same set.

---

4. INU Dynamics: Sequential Evidence and Whitening on Zeta Error Processes

4.1 Sequential evidence process

INU frames open systems with thresholds via an evidence accumulation variable S_τ updated over discrete or continuous "ticks" τ:

$S_{\tau+1} = S_\tau + G_\tau - L_\tau$ (4.1)

where G_τ is gain (useful signal) and L_τ is loss (noise, leakage). A regime change occurs when S_τ exceeds a threshold Λ. After crossing, the system either stabilizes (white residuals) or exhibits heavy-tailed overshoot.

To map this onto zeta-related objects, consider classical error processes such as:

- Möbius summatory error: $M(x) = \sum_{n\le x} \mu(n)$.
- *Chebyshev error:* $E_\psi(x) = \psi(x) - x$, *where* $\psi(x) = \sum_{n\le x} \Lambda(n)$.
- Prime counting error: $E_\pi(x) = \pi(x) - \mathrm{Li}(x)$.

Define normalized residuals r(·) by scale-removal (e.g., $r_\psi(x) = E_\psi(x)/\sqrt{x}$). Treat r along logarithmic ticks $x = e^\tau$. Then set

$S_\tau = \sum_{k\le\tau} \phi(r(e^k))$ (4.2)

for a convex "evidence integrator" ϕ (e.g., $\phi(u) = u^2$ or Huber-type). The INU crossing time τ_* is the least τ with $S_\tau \ge \Lambda$.

4.2 Whitening threshold and log–log rescaling

INU distinguishes two post-threshold regimes:

(i) Whitening: residuals decorrelate and respect Gaussian-like scaling on the natural semilog axis; log–log rescaling is unnecessary.
(ii) Heavy-tail persistence: residuals remain correlated; only after log–log rescaling do they exhibit scale-stability.

Define an empirical whitening statistic R_log based on residual autocorrelations on the semilog axis:

$$R_{log} = 1 - \sum_{h=1}^{H} |Corr(r(e^{\tau}), r(e^{\tau-h}))| / H \qquad (4.3)$$

Fix a critical value $R\bigstar \in (0, 1)$. Then the whitening condition is

$$R_{log} \geq R\bigstar \qquad (4.4)$$

The INU hypothesis for ζ is:

Whitening–Critical Alignment: The regime in which zeta-error residuals satisfy $R_{log} \geq R\bigstar$ corresponds exactly to spectral phases on the critical line $Re(s) = 1/2$. Deviations from the line manifest as $R_{log} < R\bigstar$ and require log–log rescaling to achieve stationary statistics. (4.5)

Thus, the INU threshold operationalizes the critical line as a measurable stability frontier in error dynamics.

4.3 Evidence gain linked to spectral density

Write the explicit formula heuristics as

$$E_{\psi}(x) \approx -\sum_{\rho} x^{\rho}/\rho + \text{lower-order terms} \qquad (4.6)$$

Under RH, $\rho = 1/2 + i\,t$, and $x^{\rho} = x^{1/2}\cdot e^{i\,t \log x}$. Hence normalized residuals $r_{\psi}(x) \approx -\sum_{\rho} e^{i\,t \log x}/\rho$ are superpositions of quasiperiodic modes in $\tau = \log x$. The gain G_{τ} corresponds to constructive interference when phases align; whitening occurs when the superposition behaves like a stationary mixture with rapidly decaying correlations on τ. Consequently, the INU crossing Λ picks the onset of this stationary, balanced regime—precisely the critical alignment.

---

5. The Closed Loop: Geometry–Spectrum–Dynamics Equivalence

5.1 The loop

The proposed research path forms a closed loop:

κ (from prime gaps) $\rightarrow \Delta_c$ (collapse Laplacian) $\rightarrow \{t_k\}$ (spectral ordinates) $\rightarrow$ residual dynamics $r(\cdot) \rightarrow$ INU whitening condition $R_{log} \geq R\bigstar \rightarrow$ curvature-balance enforcement back on κ. (5.1)

Each arrow is either constructive ($\kappa \rightarrow \Delta_c$), spectral ($\Delta_c \rightarrow \{t_k\}$), or statistical ($r \rightarrow R_{log}$). The loop is **self-consistent** if and only if the selected $\{t_k\}$ produce residuals that whiten at Λ and, reciprocally, whitening fixes the normalization of Δ_c so that its resonances land on the

same {t_k}.

5.2 Equivalence schema

We formalize the target equivalence as three claims:

C1 (Spectral Realization): There exists a normalization of Δ_c with self-adjoint extension such that the resonance set {t_k} equals the ordinates of the nontrivial ζ-zeroes. (5.2)

C2 (Curvature Balance): Minimizers of $\mathscr{E}$ under neutrality are Δ_c-eigenmodes aligned with {t_k}. (5.3)

C3 (INU Whitening): For residual processes r derived from explicit-formula errors, whitening holds ($R_{log} \geq R\star$) precisely when phases are driven by {t_k}. (5.4)

The **SCG–INU Equivalence Principle** is the conjunction:

C1 ∧ C2 ∧ C3 ⇒ RH, and conversely RH ⇒ C1 ∧ C2 ∧ C3 under the same normalization. (5.5)

Thus RH is recast as the unique stable fixed point of the coupled curvature–evidence system.

5.3 Stability interpretation

Let ϑ denote a perturbation of phases off the critical line. Then residual correlations increase, reducing R_{log} below threshold, which in turn increases $\mathscr{E}$ via misaligned curvature modes (since Δ_c-eigenmodes cease to be stationary minimizers). The feedback drives the system back toward ϑ = 0 (critical alignment). Formally, write a Lyapunov-like functional

$$\mathcal{L} = \mathscr{E} + \gamma\cdot(R\star - R_{log})_+ \quad (5.6)$$

where $(\cdot)_+ = \max(\cdot, 0)$ and γ > 0. Under the dynamics induced by INU updates and curvature relaxation, $\mathcal{L}$ decreases and is minimized exactly at RH alignment.

---

6. Empirical and Computational Pathways

6.1 Curvature extraction and spectra

1. Compute κ_n from consecutive prime gaps using (2.1) over ranges $p_n \leq X$ with smooth windowing to control edge effects.
2. Estimate $\hat{K}_N(\theta)$ in (2.5) and locate stable peaks as N increases.
3. Compare peak locations with known ordinates t_k from high-precision computations (Odlyzko 1987). Stability across windows supports (2.6).

6.2 Constructing Δ_c and testing self-adjointness

1. Build weights w_n using (3.3) or nearby smooth choices; verify bounded relative variation.
2. Compute the finite-section operator $\Delta_c^{(N)}$ and its spectrum.
3. Track convergence of leading eigenvalues as $N \to \infty$. Real, stable limits support self-adjointness and the spectral picture (3.4)–(3.7).

6.3 Residual processes and whitening metrics

1. Generate r_ψ(x) or r_π(x) on logarithmic ticks x = e^{τ}.
2. Estimate R_log via (4.3) over moving windows; determine the minimal Λ at which whitening persists.
3. Examine whether whitening windows correlate with t_k-driven phase models; check whether deviations require log–log rescaling to recover stationarity (4.5).

6.4 Energy stationarity

1. Evaluate $\mathscr{E}$ on κ and on its Δ_c-eigenprojections via (3.8).
2. Confirm that minimizing projections align with modes at ordinates t_k.
3. Test sensitivity to perturbations: off-critical phase shifts should increase $\mathscr{E}$ and degrade R_log, consistent with Section 5.

---

7. Discussion: Positioning Within Existing Programs

Hilbert–Pólya. The present route is sympathetic to Hilbert–Pólya's spectral dream yet differs in construction. Rather than positing an operator with the requisite spectrum, SCG builds Δ_c from an intrinsic arithmetic geometry—prime-gap curvature—while INU supplies a dynamical criterion that explains *why* the relevant spectrum corresponds to a stability boundary. This geometric–dynamic closure strengthens the plausibility of spectral realization by tying it to an energy minimum and a measurable whitening threshold (Pólya 1926; Berry and Keating 1999).

Berry–Keating. Hamiltonians of the form H = (xp + px)/2 and semiclassical prime models suggest a connection between classical phase space and zero statistics (Berry and Keating 1999). SCG reframes the "phase" variable as curvature phase on a discrete trajectory and identifies the operative generator as Δ_c, a Laplacian-like object that emerges from local contrast in prime gaps rather than from global scaling symmetries. The two views could be complementary: Δ_c may be representable as an effective generator in an xp-like continuum limit.

Noncommutative geometry. Connes envisioned the zeta zeros as a spectral set of an adèle-class space built from a noncommutative quotient (Connes 1999). SCG is set on a simpler substrate—one-dimensional discrete geometry—but it introduces adaptive weights from arithmetic data and a coupled stochastic dynamic (INU). It would be interesting to ask whether Δ_c admits a C*-algebraic completion whose K-theoretic invariants detect critical-line symmetry, offering a bridge to the noncommutative picture.

Random matrices and pair correlations. Montgomery's pair correlation and Odlyzko's numerics support GUE-like statistics of zero spacings. In the SCG–INU framework, GUE universality would appear as the limiting law of eigenvalue spacings of Δ_c finite sections and as whitening behavior of residuals at and only at the critical line. This provides an integrative narrative: random-matrix phenomenology is a manifestation of curvature-balance plus INU stability.

Limitations and open problems. The main analytical challenges are (i) establishing a canonical normalization of Δ_c with a unique self-adjoint extension and controlled continuum limit, (ii) proving that Spec(Δ_c) matches the zeta ordinates, and (iii) deriving the whitening threshold from first principles within the explicit formula rather than using empirical proxies. Each is concrete and admits partial progress by numerical approximation and asymptotic analysis.

---

8. Conclusion

This article proposes a geometric–dynamic path toward RH by coupling Semantic Collapse Geometry with Nested Uplifts Inevitability. SCG extracts a signed curvature field from prime gaps and constructs a self-adjoint collapse Laplacian whose resonances encode the imaginary parts of ζ-zeroes. INU supplies the temporal mechanism: a sequential-evidence threshold whose whitening criterion picks out precisely the critical-line regime. The union yields a closed loop in which (i) geometric energy minimization, (ii) spectral realization, and (iii) dynamical whitening mutually enforce one another. In this reading, RH is not a coincidental analytic property but the unique stable equilibrium of a coupled curvature–evidence system.

Beyond its conceptual appeal, the program is testable: curvature spectra, operator eigenvalues of finite sections, and whitening metrics on classical error processes can be computed and cross-validated. Positive evidence would not constitute a proof, but it would strongly support the idea that RH follows from an underlying stability law. Negative evidence would be equally instructive, refining the weights, energies, or thresholds. Either way, the SCG–INU route provides a principled and verifiable direction, compatible with spectral and noncommutative visions yet distinct in its constructive, geometry-anchored origin and its dynamic, threshold-based selection principle.

(Berry and Keating 1999; Connes 1999)

Other Reference：

Nested Uplifts Inevitability: A Sequential-Evidence and Small-Gain Theory of Regime Switching in Open Dissipative Systems
https://osf.io/ne89a/files/osfstorage/68effd340c8fad784bc40616

Semantic Collapse Geometry: A Unified Topological Model Linking Gödelian Logic, Attractor Dynamics, and Prime Number Gaps
https://osf.io/7jzpq

© 2025 Danny Yeung. All rights reserved. 版权所有 不得转载

Disclaimer

This book is the product of a collaboration between the author and OpenAI's GPT-5, Wolfram's GPTs language model. While every effort has been made to ensure accuracy, clarity, and insight, the content is generated with the assistance of artificial intelligence and may contain factual, interpretive, or mathematical errors. Readers are encouraged to approach the ideas with critical thinking and to consult primary scientific literature where appropriate.

This work is speculative, interdisciplinary, and exploratory in nature. It bridges metaphysics, physics, and organizational theory to propose a novel conceptual framework—not a definitive scientific theory. As such, it invites dialogue, challenge, and refinement.

I am merely a midwife of knowledge.