Blessed G/Michael Catholic Secondary School

First Semester Agriculture final exam for Grade 11th

January, 2018 E.C Time allowed 50'

Part I: - Choose the best answer from the given alternatives.

C 1. Which one of the following types of animal identification is NOT recommended?
A. Tattooing B. Ear notching C. Branding D. Ear tag

C 2. Which of the following is the highest livestock population species in Ethiopia?
A. Sheep B. Chickens C. cattle D. Goats

E 3. Which type of animal unable to digest much cellulose feed materials?
A. goats B. Sheep C. horse D. Cow E. none

C 4. Which of the following is NOT the component of a mono gastric digestive system?
A. Esophagus B. Small intestine C. Reticulum D. mouth E. none

C 5. among the following, which one is **an energy-rich concentrates feed** for livestock's?
A. Cotton seed cake B. dried blood meal C. Middling D. Fish meal

A 6. which of the following is true about livestock feed?
A. more than 18 % CF feeds are roughage feeds C. Less than 18% CP feeds are concentrate feeds
B. More than 60% TDN feeds are roughage feeds D. Less than 60% TDN feeds are concentrate feeds

E (7.)----------are inorganic elements that required in small amounts but are important components of animal feed.
A. Proteins B. Fats C. Carbohydrates D. Vitamin's E. none

B 8. Which of the following is true about the limitation of cross-breeding?
A. Reduction in production B. Delivery difficulties C. Loss of fertility D. Promote heterosis

C 9. Which of the following reproductive techniques helps the rapid exploitation of superior females in the herd?
A. Animal breeding B. Artificial insemination C. Embryo transfer D. Animal genetics

C 10. All livestock feeds are roughages, EXCEPT?
A. Teff straw B. Hay C. Molasses D. Succulent feed

D 11. Which of the following is correct about the shortcoming of hay?
A. Hay making incurs high cost C. Hay is difficult to transport and store
B. It can be fed with little or no wastage D. Hay making is dependent on the weather conditio

B (12) the most important criterion for selection of a breed for commercial production is--
A. Body color B. Adaptability to local environment C. Horn shape D. Breed name

B 13. Which of the following should be avoided while selecting a breed?

A. High production potential

C. Resistance to diseases

B. Poor adaptability

D. Good reproductive efficiency

B (14) Which type of breeding is commonly used in commercial poultry and pig production?

A. Inbreeding B. Line breeding C. Cross breeding D. Pure breeding

A 15. Which of the following feeds improve to milk fat percentage in dairy animals?

A. High protein B. High roughage diet C. Vitamin supplement. D. Mineral mixture

A 16. Among the following forms of extensive production system, which one involves relatively a better livestock management?

A. Ranching B. Agro-pastoralist C. Nomadism D. Pastoralism

D 17. The primary function of water for livestock production?

A. transport oxygen in the blood stream B. Building muscle mass

C. for metabolic process D. All of the above

C 18. The final holding area for a function as an excretory and reproductive duct of avian is------

A. Gizzard B. Crop C. Cloaca D. Colon

C 19. The type of breeds that helps to produce pure breed and to preserve specific qualities in an animals is -----

A. Cross breeding B. Line breeding C. In- breeding D. Out breeding

D 20. Which of the following is true about the important of record keeping in livestock farms?

A. Helps in making culling decisions C. Eases the selection of new breeding stock

B. Helps producers identify area of weakness D. All E. B&C

Part II:- Match items under column A with items under column B

Column A	Column B
D 21. Heat detection	A. Liquid nitrogen
B 22. AI gun	B. To precisely deliver semen into an animal's reproductive tract
G 23. Inbreeding depression.	C. 34°C water bath
A 24. AI materials	D. Hormonal control of heat
F 25. Estrus synchronization	E. Semen stored at –196°C
C 26. Thawing	F. Standing to be mounted
	G. Reduced fertility & vigor

Part III: - work out (weight mark 4 pts)

27. Framer wants to be formulate a poultry ration having a maize **8%** CP, sorghum **10%**CP, soybear meal **44%** CP and fish meal **65%** CP will provide a mix of **16.5%** CP? Assume **maize** anc **sorghum** are grouped together and used in the proportion **2:1** and **soybean meal** and **fish meal** are grouped together in the proportion of **3:1** respectively.

25 good

Answer Sheet

Name Muhami- Menole Section JJC No 33

1. C	6. A	11. D	16. A	21. D F
2. C	7. E	12. B	17. D	22. B
3. E C	8. B	13. B	18. C	23. G
4. C	9. C	14. B C	19. C	24. A
5. C	10. C	15. A B	20. D	25. E D

26. C

Work out

27. Given

maize = 8%

sorghum = 10%

SBM = 44%

FM = 65%

CP = 16.5


M : S = 2 : 1

SBM : FM = 3 : 1

mass = 100kg

$M+S = \frac{8*2+10*1}{3} = \frac{16+10}{3} = \frac{26}{3} = 8.7\%$

$SBM+FM = \frac{44*3+65*1}{4} = \frac{132+65}{4} = 49.25\%$



$= 40.6$

Next Page

$M = \frac{32800*2}{405*3} = \frac{65600}{1218}$ kg

$S = \frac{32800*1}{405*3} = \frac{32800}{1218}$ kg

$M = \frac{65600}{1218}$ kg

$S = \frac{32800}{1218}$ kg

$M+S = \frac{32.8*100}{40.6} = \frac{32800}{405}$ kg

$SBM+FM = \frac{7.8*100}{40.6} = \frac{7800}{405}$

$SBM = \frac{23400}{1624}$ kg

$FM = \frac{7800}{1624}$ kg

$SBM = \frac{3*7800}{405*4} = \frac{23400}{1624}$ kg


$FM = \frac{1}{4}*\frac{7800}{406} = \frac{7800}{1624}$ kg

C 38. The project was feasible only if the government p...
A. The project was impossible regardless of government help.
... project was still a success.

M = 8%
S = 10%
SBM = 44%
FM = 65%
M = 100 kg

$M+S = \frac{8+2+1000}{3} = \frac{26}{3} = 8.7\%$

$SBM+FM = \frac{44+3+65}{4} = \frac{132+65}{4} = \frac{197}{4} = 49.3\%$



$M+S = \frac{32.8 \times 100}{40.6} = \frac{328 \times 100}{406} = \frac{32800}{406}$ kg

$SBM+FM = \frac{7.8 \times 100}{40.6} = \frac{7800}{406}$ kg

$M = \frac{2}{3} \times \frac{32800}{406} = \frac{65600}{1218}$ kg

$S = \frac{1}{3} \times \frac{32800}{406} = \frac{3280}{1218}$ kg

$SBM = \frac{3}{4} \times \frac{7800}{406} = \frac{23400}{1624}$ kg

$FM = \frac{1}{4} \times \frac{7800}{406} = \frac{7800}{1624}$ kg

Blessed G/Michael Catholic secondary school

Agriculture Group work for grade 11 students (25-36)

Submission date 17/4/2018 E.C

1. What combination of a maize **8%** CP, sorghum **10%**CP, soybean meal **44%** CP and fish meal **65%** CP will provide a mix of **16.5%** CP? Assume **maize** and **sorghum** are grouped together and used in the proportion **2:1** and **soybean meal** and **fish meal** are grouped together in the proportion of **3:1** respectively.

Given

Maize = M = 8% CP

Sorghum = S = 10% CP

Soybean meal = SBM = 44% CP

Fish meal = F = 65% CP

Mix of = 16.5 % CP.


M : S = 2 : 1

SBM : F = 3 : 1

Soln

$= \frac{2*8+1*10}{3} = \frac{16+10}{3} = \frac{26}{3} = 8.7\%$

$= \frac{3*44+65*1}{4} = \frac{197}{4} = 49.3\%$



lets take 100Kg here.

$MS = \frac{32.8*100}{40.6} = 80.8$ Kg

$SF = \frac{7.8*100}{3} = 19.2$ Kg

$S = \frac{80.8}{3} = 26.9$ Kg

$M = \frac{80.8*2}{3} = 53.9$ Kg

$SBM = \frac{19.2*3}{4} = 14.4$ Kg

$F = \frac{19.2}{4} = 4.8$ Kg

SO there will be 26.9 Kg of Sorghum, 53.9 Kg of Maize, 14.4 SBM and 4.8 Kg of Fish meal.