

# Iterative Algorithms in Optimization, Variational Analysis and Fixed Point Theory

**Unit 01: Introduction. Fundamentals of iterative algorithms.**



# About the course

## Description

This course provides the theoretical bases for the numerical approximation of solutions to optimization problems, variational inequalities and fixed point problems by means of iterative algorithms.

## Assessment


Computational exercises 20%, Homework assignments 50%, Exam 30%.

## Dynamics and Schedule

16 sessions: Lectures, exercises. Mo, 9-11. Tu, 9-11. We, 17-19.


# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$




# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$




# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$




# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$




# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$



# Introduction to iterative algorithms


Example: Bisection method to solve  $g(x) = 0$






# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$




# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$




# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$



# Introduction to iterative algorithms

Example: Bisection method to solve  $g(x) = 0$



After  $k$  iterations, the distance to a solution is  $|x_k - \hat{x}| \leq \frac{b - a}{2^k}$ .

# Problems to be solved

We shall approximate points in a set  $S$  of solutions in different contexts:

- $S = \text{Argmin}(f)$ . Minimizers of an extended real-valued function  $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ , especially in the convex case.

# Problems to be solved

We shall approximate points in a set  $S$  of solutions in different contexts:

- $S = \text{Argmin}(f)$ . Minimizers of an extended real-valued function  $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ , especially in the convex case.
- $S = \text{Zer}(A)$ . Zeroes of a set-valued operator  $A : \mathbb{R}^N \rightarrow 2^{\mathbb{R}^N}$ , especially in the monotone case.

# Problems to be solved

We shall approximate points in a set  $S$  of solutions in different contexts:

- $S = \text{Argmin}(f)$ . Minimizers of an extended real-valued function  $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ , especially in the convex case.
- $S = \text{Zer}(A)$ . Zeroes of a set-valued operator  $A : \mathbb{R}^N \rightarrow 2^{\mathbb{R}^N}$ , especially in the monotone case.
- $S = \text{Fix}(T)$ . Fixed points of a function  $T : \mathbb{R}^N \rightarrow \mathbb{R}^N$ , especially in the nonexpansive case.

# Quality of approximate solutions

A **merit function**  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}_+$  for a problem with solution set  $S$  satisfies:

- $\Phi(z) = 0$  if  $z \in S$ , and  $\Phi(z) > 0$  otherwise.
- If  $\Phi(z) < \Phi(w)$ , then  $z$  is a better approximate solution than  $w$ .



# Quality of approximate solutions

A **merit function**  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}_+$  for a problem with solution set  $S$  satisfies:

- $\Phi(z) = 0$  if  $z \in S$ , and  $\Phi(z) > 0$  otherwise.
- If  $\Phi(z) < \Phi(w)$ , then  $z$  is a better approximate solution than  $w$ .

## Examples

- $\Phi(z) = f(z) - \inf(f)$  for  $S = \text{Argmin}(f)$ .

# Quality of approximate solutions

A **merit function**  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}_+$  for a problem with solution set  $S$  satisfies:

- $\Phi(z) = 0$  if  $z \in S$ , and  $\Phi(z) > 0$  otherwise.
- If  $\Phi(z) < \Phi(w)$ , then  $z$  is a better approximate solution than  $w$ .

## Examples

- $\Phi(z) = f(z) - \inf(f)$  for  $S = \text{Argmin}(f)$ .

Also  $\Phi(z) = \|\nabla f(z)\|$ , if  $f$  is convex and differentiable.

# Quality of approximate solutions

A **merit function**  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}_+$  for a problem with solution set  $S$  satisfies:

- $\Phi(z) = 0$  if  $z \in S$ , and  $\Phi(z) > 0$  otherwise.
- If  $\Phi(z) < \Phi(w)$ , then  $z$  is a better approximate solution than  $w$ .

## Examples

- $\Phi(z) = f(z) - \inf(f)$  for  $S = \text{Argmin}(f)$ .  
Also  $\Phi(z) = \|\nabla f(z)\|$ , if  $f$  is convex and differentiable.
- $\Phi(z) = \inf\{\|v\| : v \in Az\}$  for  $S = \text{Zer}(A)$ .

# Quality of approximate solutions

A **merit function**  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}_+$  for a problem with solution set  $S$  satisfies:

- $\Phi(z) = 0$  if  $z \in S$ , and  $\Phi(z) > 0$  otherwise.
- If  $\Phi(z) < \Phi(w)$ , then  $z$  is a better approximate solution than  $w$ .

## Examples

- $\Phi(z) = f(z) - \inf(f)$  for  $S = \text{Argmin}(f)$ .  
Also  $\Phi(z) = \|\nabla f(z)\|$ , if  $f$  is convex and differentiable.
- $\Phi(z) = \inf\{\|v\| : v \in Az\}$  for  $S = \text{Zer}(A)$ .
- $\Phi(z) = \|Tz - z\|$  or  $\Phi(z) = \|Tz - z\|^2$  for  $S = \text{Fix}(T)$ .

# Quality of approximate solutions

A **merit function**  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}_+$  for a problem with solution set  $S$  satisfies:

- $\Phi(z) = 0$  if  $z \in S$ , and  $\Phi(z) > 0$  otherwise.
- If  $\Phi(z) < \Phi(w)$ , then  $z$  is a better approximate solution than  $w$ .

## Examples

- $\Phi(z) = f(z) - \inf(f)$  for  $S = \text{Argmin}(f)$ .  
Also  $\Phi(z) = \|\nabla f(z)\|$ , if  $f$  is convex and differentiable.
- $\Phi(z) = \inf\{\|v\| : v \in Az\}$  for  $S = \text{Zer}(A)$ .
- $\Phi(z) = \|Tz - z\|$  or  $\Phi(z) = \|Tz - z\|^2$  for  $S = \text{Fix}(T)$ .
- $\Phi(z) = \text{dist}(z, S)$  in all cases ( $\Phi(z) = \|z - \hat{x}\|$  if  $S = \{\hat{x}\}$ ).

# Introduction to iterative algorithms

An **iterative algorithm** is a procedure that computes a sequence  $(x_k)$  of points in  $\mathbb{R}^N$  that approximate a solution to a problem.

# Introduction to iterative algorithms

An **iterative algorithm** is a procedure that computes a sequence  $(x_k)$  of points in  $\mathbb{R}^N$  that approximate a solution to a problem. It requires:

- An **initial guess**  $x_0$ .

# Introduction to iterative algorithms

An **iterative algorithm** is a procedure that computes a sequence  $(x_k)$  of points in  $\mathbb{R}^N$  that approximate a solution to a problem. It requires:

- An **initial guess**  $x_0$ .
- A sequence  $(u_k)$  in  $\mathbb{R}^M$  of **parameters**.



# Introduction to iterative algorithms

An **iterative algorithm** is a procedure that computes a sequence  $(x_k)$  of points in  $\mathbb{R}^N$  that approximate a solution to a problem. It requires:

- An **initial guess**  $x_0$ .
- A sequence  $(u_k)$  in  $\mathbb{R}^M$  of **parameters**.
- An operator  $F : \mathbb{R}^N \times \mathbb{R}^M \rightarrow \mathbb{R}^N$  used to compute  $x_{k+1}$ , given  $x_k$ :

$$x_{k+1} = F(u_k, x_k).$$

# Introduction to iterative algorithms

An **iterative algorithm** is a procedure that computes a sequence  $(x_k)$  of points in  $\mathbb{R}^N$  that approximate a solution to a problem. It requires:

- An **initial guess**  $x_0$ .
- A sequence  $(u_k)$  in  $\mathbb{R}^M$  of **parameters**.
- An operator  $F : \mathbb{R}^N \times \mathbb{R}^M \rightarrow \mathbb{R}^N$  used to compute  $x_{k+1}$ , given  $x_k$ :

$$x_{k+1} = F(u_k, x_k).$$

- A stopping rule, related to a (normalized) merit function, that is activated when the approximation is **sufficiently good**.

# Introduction to iterative algorithms

An **iterative algorithm** is a procedure that computes a sequence  $(x_k)$  of points in  $\mathbb{R}^N$  that approximate a solution to a problem. It requires:

- An **initial guess**  $x_0$ .
- A sequence  $(u_k)$  in  $\mathbb{R}^M$  of **parameters**.
- An operator  $F : \mathbb{R}^N \times \mathbb{R}^M \rightarrow \mathbb{R}^N$  used to compute  $x_{k+1}$ , given  $x_k$ :

$$x_{k+1} = F(u_k, x_k).$$

- A stopping rule, related to a (normalized) merit function, that is activated when the approximation is **sufficiently good**.

Important questions: **convergence** and **complexity**.

# Convergence

A sequence  $(x_k)$  in  $\mathbb{R}^N$  **converges** to a **limit**  $\hat{x} \in \mathbb{R}^N$  as  $k \rightarrow \infty$  if, for every  $\varepsilon > 0$ , there is  $K \in \mathbb{N}$  such that

$$\|x_k - \hat{x}\| < \varepsilon$$

for every  $k \geq K$ .

# Convergence

A sequence  $(x_k)$  in  $\mathbb{R}^N$  **converges** to a **limit**  $\hat{x} \in \mathbb{R}^N$  as  $k \rightarrow \infty$  if, for every  $\varepsilon > 0$ , there is  $K \in \mathbb{N}$  such that

$$\|x_k - \hat{x}\| < \varepsilon$$

for every  $k \geq K$ .

## Proposition

*The following statements about a sequence  $(x_k)$  in  $\mathbb{R}^N$  are equivalent:*

- i) it is convergent;*
- i) it has Cauchy's property; and*
- i) it is bounded and has at most one limit point.*

# Keynote examples

## Theorem (Banach-Picard Iterations and the Fixed Point Theorem)

Let  $q \in (0, 1)$ , and let  $T : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be  $q$ -Lipschitz. Then,  $T$  has a unique fixed point  $\hat{x}$ . Moreover, for each  $x_0 \in \mathbb{R}^N$ , the sequence  $(x_k)$ , defined by iterating  $x_{k+1} = Tx_k$ , converges to  $\hat{x}$ .

# Keynote examples

## Theorem (Banach-Picard Iterations and the Fixed Point Theorem)

Let  $q \in (0, 1)$ , and let  $T : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be  $q$ -Lipschitz. Then,  $T$  has a unique fixed point  $\hat{x}$ . Moreover, for each  $x_0 \in \mathbb{R}^N$ , the sequence  $(x_k)$ , defined by iterating  $x_{k+1} = Tx_k$ , converges to  $\hat{x}$ .

## Theorem (Krasnoselskii-Mann Iterations)

Let  $\gamma \in (0, 1)$  and let  $T : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be nonexpansive (1-Lipschitz), with  $\text{Fix}(T) \neq \emptyset$ . For each  $x_0 \in \mathbb{R}^N$ , the sequence  $(x_k)$ , defined by iterating  $x_{k+1} = \gamma Tx_k + (1 - \gamma)x_k$ , converges to a fixed point of  $T$ .

# Useful shortcuts

## Lemma (Opial)

Let  $\Omega$  be a nonempty subset of  $\mathbb{R}^N$ , and let  $(x_k)$  be a sequence in  $\mathbb{R}^N$ . Suppose that:

- i) For every  $y \in \Omega$ ,  $\lim_{k \rightarrow \infty} \|x_k - y\|$  exists; and
- ii) Every limit point of  $(x_k)$  belongs to  $\Omega$ .

Then,  $(x_k)$  converges, as  $k \rightarrow \infty$ , to a point in  $\Omega$ .



# Useful shortcuts

## Lemma (Opial)

Let  $\Omega$  be a nonempty subset of  $\mathbb{R}^N$ , and let  $(x_k)$  be a sequence in  $\mathbb{R}^N$ . Suppose that:

- i) For every  $y \in \Omega$ ,  $\lim_{k \rightarrow \infty} \|x_k - y\|$  exists; and
- ii) Every limit point of  $(x_k)$  belongs to  $\Omega$ .

Then,  $(x_k)$  converges, as  $k \rightarrow \infty$ , to a point in  $\Omega$ .

## Remark (Fejér)

If  $\|x_{k+1} - y\| \leq \|x_k - y\|$  for all  $k \geq 0$  and  $y \in \Omega$ , then  $\lim_{k \rightarrow \infty} \|x_k - y\|$  exists for every  $y \in \Omega$ .

# Convergence rates

Let  $(\phi_k)$  be nonnegative, and let  $(\rho_k)$  be positive, with  $\lim_{k \rightarrow \infty} \rho_k = 0$

We shall use the following notation:

- $\phi_k = \mathcal{O}(\rho_k)$  means that  $\phi_k \rightarrow 0$  at least as fast as  $\rho_k$ :

$$C := \sup_{k \geq 0} \left[ \frac{\phi_k}{\rho_k} \right] < +\infty,$$

# Convergence rates

Let  $(\phi_k)$  be nonnegative, and let  $(\rho_k)$  be positive, with  $\lim_{k \rightarrow \infty} \rho_k = 0$

We shall use the following notation:

- $\phi_k = \mathcal{O}(\rho_k)$  means that  $\phi_k \rightarrow 0$  at least as fast as  $\rho_k$ :

$$C := \sup_{k \geq 0} \left[ \frac{\phi_k}{\rho_k} \right] < +\infty,$$

- $\phi_k = o(\rho_k)$  means that  $\phi_k \rightarrow 0$  strictly faster than  $\rho_k$ :

$$\lim_{k \rightarrow \infty} \left[ \frac{\phi_k}{\rho_k} \right] = 0.$$

# Convergence rates

Let  $(\phi_k)$  be nonnegative, and let  $(\rho_k)$  be positive, with  $\lim_{k \rightarrow \infty} \rho_k = 0$

We shall use the following notation:

- $\phi_k = \mathcal{O}(\rho_k)$  means that  $\phi_k \rightarrow 0$  at least as fast as  $\rho_k$ :

$$C := \sup_{k \geq 0} \left[ \frac{\phi_k}{\rho_k} \right] < +\infty,$$

- $\phi_k = o(\rho_k)$  means that  $\phi_k \rightarrow 0$  strictly faster than  $\rho_k$ :

$$\lim_{k \rightarrow \infty} \left[ \frac{\phi_k}{\rho_k} \right] = 0.$$

Clearly,  $\phi_k = o(\rho_k)$  implies  $\phi_k = \mathcal{O}(\rho_k)$ , but the constant  $C$  is important!

# Convergence rates

## Some terminology

- If  $\phi_k = \mathcal{O}(q^k)$  for some  $q \in (0, 1)$ , we say  $\phi_k$  converges **linearly** to zero.

# Convergence rates

## Some terminology

- If  $\phi_k = \mathcal{O}(q^k)$  for some  $q \in (0, 1)$ , we say  $\phi_k$  converges **linearly** to zero.
  - It is the case for Banach-Picard iterations.

# Convergence rates

## Some terminology

- If  $\phi_k = \mathcal{O}(q^k)$  for some  $q \in (0, 1)$ , we say  $\phi_k$  converges **linearly** to zero.
  - It is the case for Banach-Picard iterations.
- If  $\phi_k \neq \mathcal{O}(q^k)$  for every  $q \in (0, 1)$ ,  $\phi_k$  converges **sublinearly**.
  - En many frequent cases, we still have  $\phi_k = \mathcal{O}(k^{-p})$  for some  $p > 0$ .

# Convergence rates

## Some terminology

- If  $\phi_k = \mathcal{O}(q^k)$  for some  $q \in (0, 1)$ , we say  $\phi_k$  converges **linearly** to zero.
  - It is the case for Banach-Picard iterations.
- If  $\phi_k \neq \mathcal{O}(q^k)$  for every  $q \in (0, 1)$ ,  $\phi_k$  converges **sublinearly**.
  - En many frequent cases, we still have  $\phi_k = \mathcal{O}(k^{-p})$  for some  $p > 0$ .
- Finally, if  $\phi_k = o(q^k)$  for all  $q \in (0, 1)$ , then  $\phi_k$  converges **superlinearly**.



# Convergence rates

## Some terminology

- If  $\phi_k = \mathcal{O}(q^k)$  for some  $q \in (0, 1)$ , we say  $\phi_k$  converges **linearly** to zero.
  - It is the case for Banach-Picard iterations.
- If  $\phi_k \neq \mathcal{O}(q^k)$  for every  $q \in (0, 1)$ ,  $\phi_k$  converges **sublinearly**.
  - En many frequent cases, we still have  $\phi_k = \mathcal{O}(k^{-p})$  for some  $p > 0$ .
- Finally, if  $\phi_k = o(q^k)$  for all  $q \in (0, 1)$ , then  $\phi_k$  converges **superlinearly**.
  - It is the case for Newton's method, under favorable conditions.

# Practical aspects

## Linear regression

Consider a set  $\{(u_k, v_k)\}_{k=1}^K$  of samples corresponding to  $v \sim mu + b$ , where  $m$  and  $b$  are unknown.

# Practical aspects

## Linear regression

Consider a set  $\{(u_k, v_k)\}_{k=1}^K$  of samples corresponding to  $v \sim mu + b$ , where  $m$  and  $b$  are unknown. Define

$$\Psi(m, b) = \frac{1}{2} \sum_{k=1}^K (v_k - mu_k - b)^2.$$

We look for the values of  $m$  and  $b$  that minimize  $\Psi$ .

# Practical aspects

## Linear regression

Consider a set  $\{(u_k, v_k)\}_{k=1}^K$  of samples corresponding to  $v \sim mu + b$ , where  $m$  and  $b$  are unknown. Define

$$\Psi(m, b) = \frac{1}{2} \sum_{k=1}^K (v_k - mu_k - b)^2.$$

We look for the values of  $m$  and  $b$  that minimize  $\Psi$ . The relation  $\nabla \Psi(\hat{m}, \hat{b}) = 0$  gives a system of two equations with two unknowns:

$$\begin{cases} \hat{m} \sum u_k^2 + \hat{b} \sum u_k &= \sum u_k v_k \\ \hat{m} \sum u_k + \hat{b} K &= \sum v_k \end{cases}$$

# Practical aspects

## Linear regression

Consider a set  $\{(u_k, v_k)\}_{k=1}^K$  of samples corresponding to  $v \sim mu + b$ , where  $m$  and  $b$  are unknown. Define

$$\Psi(m, b) = \frac{1}{2} \sum_{k=1}^K (v_k - mu_k - b)^2.$$

We look for the values of  $m$  and  $b$  that minimize  $\Psi$ . The relation  $\nabla \Psi(\hat{m}, \hat{b}) = 0$  gives a system of two equations with two unknowns:

$$\begin{cases} \hat{m} \sum u_k^2 + \hat{b} \sum u_k &= \sum u_k v_k \\ \hat{m} \sum u_k + \hat{b} K &= \sum v_k \end{cases}$$

If we suspect that  $\Phi(x_k)$  converges linearly to zero, we can use this with  $u_k = k$ , and  $v_k = \ln(\Phi(x_k))$ .

# Complexity

Consider a problem with merit function  $\Phi$  and a sequence  $(x_k)$  produced by an algorithm

Suppose that for every  $\varepsilon > 0$ , which represents a **tolerance** or **precision level**, there is a number  $k_\varepsilon$  such that

$$\Phi(x_{k_\varepsilon}) < \varepsilon.$$

A **complexity bound** is either a map  $\varepsilon \mapsto k_\varepsilon$ , or an inequality that maps  $\varepsilon$  to a set of valid  $k_\varepsilon$ 's.

# Complexity

Consider a problem with merit function  $\Phi$  and a sequence  $(x_k)$  produced by an algorithm

Suppose that for every  $\varepsilon > 0$ , which represents a **tolerance** or **precision level**, there is a number  $k_\varepsilon$  such that

$$\Phi(x_{k_\varepsilon}) < \varepsilon.$$

A **complexity bound** is either a map  $\varepsilon \mapsto k_\varepsilon$ , or an inequality that maps  $\varepsilon$  to a set of valid  $k_\varepsilon$ 's.

## Complexity bounds from convergence rates

- From  $\Phi(x_k) \leq C/k^p$ , we get  $k > \sqrt[p]{C/\varepsilon}$ .

# Complexity

Consider a problem with merit function  $\Phi$  and a sequence  $(x_k)$  produced by an algorithm

Suppose that for every  $\varepsilon > 0$ , which represents a **tolerance** or **precision level**, there is a number  $k_\varepsilon$  such that

$$\Phi(x_{k_\varepsilon}) < \varepsilon.$$

A **complexity bound** is either a map  $\varepsilon \mapsto k_\varepsilon$ , or an inequality that maps  $\varepsilon$  to a set of valid  $k_\varepsilon$ 's.

## Complexity bounds from convergence rates

- From  $\Phi(x_k) \leq C/k^p$ , we get  $k > \sqrt[p]{C/\varepsilon}$ .
- From  $\Phi(x_k) \leq Ae^{-Bk}$ , we get  $k > \ln \left( \sqrt[B]{A/\varepsilon} \right)$ .



# Examples

## Banach-Picard Iterations

We have

$$\|Tx_k - x_k\| \leq \|Tx_0 - x_0\| q^k.$$

# Examples

## Banach-Picard Iterations

We have

$$\|Tx_k - x_k\| \leq \|Tx_0 - x_0\| q^k.$$

## Krasnoselskii-Mann Iterations

We can prove that

$$\|Tx_k - x_k\|^2 \leq \frac{\text{dist}(x_0, S)^2}{\gamma(1-\gamma)k} = \mathcal{O}\left(\frac{1}{k}\right) \quad \text{and} \quad \|Tx_k - x_k\|^2 = o\left(\frac{1}{k}\right).$$

# Exercises

- ❶ Let  $(a_k)$ ,  $(b_k)$  and  $(c_k)$  be nonnegative sequences, and suppose

$$a_{k+1} + b_k \leq a_k + c_k$$

for all  $k \geq 0$ . Assume  $C := \sum c_k < +\infty$ . Show that

$$\lim_{k \rightarrow \infty} a_k \text{ exists,} \quad \text{and} \quad \sum b_k \leq a_0 + C < +\infty.$$

- ❷ Let  $(d_k)$  be nonnegative, with  $D := \sum d_k < +\infty$ , and write  $\delta_k = \min\{d_0, \dots, d_k\}$ . Prove that

$$(k+1)\delta_k \leq D \quad \text{and} \quad \lim_{k \rightarrow \infty} (k+1)\delta_k = 0.$$

## Exercises, continued

- ③ Let  $T$  be nonexpansive, with  $\text{Fix}(T) \neq \emptyset$ . Apply Krasnoselskii-Mann iterations

$$x_{k+1} = \gamma T x_k + (1 - \gamma)x_k.$$

- ① Show that  $\|x_{k+1} - \hat{x}\|^2 + \gamma(1 - \gamma)\|T x_k - x_k\|^2 \leq \|x_k - \hat{x}\|^2$  for every  $\hat{x} \in \text{Fix}(T)$ . Conclude that  $(x_k)$  converges to a fixed point of  $T$ .
- ② Verify that  $k \mapsto \|T x_k - x_k\|^2$  is nonincreasing.
- ③ Prove that the following convergence rates hold:

$$\|T x_k - x_k\|^2 \leq \frac{\text{dist}(x_0, S)^2}{\gamma(1 - \gamma)k} = \mathcal{O}\left(\frac{1}{k}\right) \quad \text{and} \quad \|T x_k - x_k\|^2 = o\left(\frac{1}{k}\right).$$