

a female (FlyWire): the turning DNs
paper's model (known signs, eLN ×0.25)
+ 0.2 Hz substrate
published lobe (Shiu signs, eLN ×1)
+ 0.2 Hz substrate
known signs, eLN ×1
+ 0.2 Hz substrate
Shiu signs, eLN ×0.25
+ 0.2 Hz substrate
known signs, eLN ×0.5
+ 0.2 Hz substrate
known signs, eLN ×0.1
+ 0.2 Hz substrate
Zenodo table, ≥ 5 synapses (first version)
+ 0.2 Hz substrate
Zenodo table, all connections
+ 0.2 Hz substrate
0 5 10 15 20 25 30
turning DNs DNa01/02, mean of both sides (Hz)
b female: suppression index
cVA, Or47b: DNa 0 → 12 Hz with touch
−2.0 −1.5 −1.0 −0.5 0.0 0.5 1.0
1 − DNa(+ touch) / DNa(alone)
c male (MaleCNS): MDN (backward walking)
paper's model (eLN ×0.25)
+ 0.2 Hz substrate
published lobe (eLN ×1)
+ 0.2 Hz substrate
eLN ×0.25, unclear LNs inhibitory
+ 0.2 Hz substrate
eLN ×0.25, unclear LNs silent
+ 0.2 Hz substrate
0 5 10 15 20 25 30 35
MDN, backward walking (Hz)
walking drive (100 Hz, 600 ms, 5 seeds)
food odor
cVA
Or47b (fly odor)
ppk23/25 alone → ppk25 + bristles
substrate alone
a, c: the walking drive alone
a, c: drive + head bristles (100 Hz)
b: hollow = DNa alone < 2 Hz (not driven)
b: index > 0.5 (< 0 = the bristles excite)
the ≥ 5-synapse connection threshold