1. The probability of a leap year selected at random contain 53 Sunday is:							
(a) 53/ 366 (b) 1/7 (c) 2/7 (d) 53/365							
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at							
random. The probability of drawing a black ball is :							
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5							
3. The probability that it will rain tomorrow is 0.85. What is the							
probability that it will not rain tomorrow							
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these							
4. What is the probability that a number selected from the numbers							
(1, 2, 3,,15) is a multiple of 4?							
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3							
5. What are the total outcomes when we throw three coins?							
(a) 4 (b) 5 (c) 8 (d) 7							
6. The probability that a prime number selected at random from the							
numbers (1,2,3,35) is :							
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these							
7. The sum of the probability of an event and non event is :							
(a) 2 (b) 1 (c) 0 (d) none of these.							
8. The following probabilities are given; choose the correct answer							
for that which is not possible.							
(a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.							
9. If three coins are tossed simultaneously, than the probability of							
getting at least two heads, is:							
(a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/8$							
10. A letter is chosen at random from the letters of the word							
♦ ASSASSINATION ♦ . The probability that the letter chosen has:							
(a) 6/13 (b) 7/13 (c) 1 (d) none of these.							
(3) 1, 10 (3) 1, 10							
11. A dice is thrown. Find the probability of getting an even number.							
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2							
12. Two coins are thrown at the same time. Find the probability of getting both heads.							
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0							
13. Two dice are thrown simultaneously. The probability of getting a							

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/9)			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4		(C) 1/4	(D) 2	29/100			
•	a blue ball is d	ouble that of a		the probability the number of			
16. A box of	f 600 bulbs co random from ve bulb is:	ntains 12 defections this box. Then	ctive bulbs. C				
mixed thoro	oughly. One ca lity that the nu	mbers 2 to 101 rd is drawn fro mber on card i (C) 3/10	m this box ra s a perfect so	ndomly, then quare.			
18. What is (A) 1/7	the probabilit (B) 53/366	y of getting 53 (C) 2/7	Mondays in (D) 7/3				
 19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13 							
equally likel 1,2,312	ly to come to r then the proba	nsists of spinn est pointing to ability that it w (C) 7/12	one of the ni ill point to an	umber odd number is:			
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4							

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225				
2. Then the pr	x is chosen at ra cobability that x ² < 2/5 (C) 3/5	: 2 is?	numbers -2, -1, 0 , 1,				
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) 7/50 (B) 4/25 (C) 1/25 (D) 2/25							
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
	-		hes are 50, 70, 82,				
	ne standard devia b) 25.49		d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$ 30. X is a varia	b) $^1\!/_3$ ate between 0 and	d 3. The value of	d) $\frac{1}{6}$ E(X ²) is				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of the probability?	•	alues, whic	h one is not po	ssible in			
a) $P(x) = 1$	b) ∑ x d) P(x	P(x) = 3 () = -0.5					
33.If E(x) = a) 2	2 and E(z) = 4 b) 6	c) 0		nsufficient data			
34.The cova	ariance of two	independe	nt random varia	able is			
a) 1	b) 0	c) - 1	d) (Jndefined			
35.If Σ P(x) a) 0	= k² – 8 then, b) 1	the value o		Insufficient data			
, ,	0.5 and x = 4, b) 0.5	, ,	? d) 2	2			
37.In a discrete probability distribution, the sum of all probabilities is always?							
a) 0	b) Infinite	c) 1	d) U	ndefined			
38.If the pr	obability of hi	tting the tai	get is 0.4, find	mean and			
	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16			
-	-	mbs are dr		ace will strike the an and variance? d) 4, 1.6			
a) 2	e mean of toss b) 4 s the mean and	c) 8	d) 1	rmal distribution?			

c) 5

d) 7

a) 3

b) 4

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0								
42. Variance of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2)$ – $(E(X))2$ d) $(E(X))2$								
	43.Mean of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2) - (E(X))2$ d) $(E(X))2$							
44.Mean of a constant 'a' is a) 0								
45.Variance of a constant 'a' is . a) 0								
46.Find the mean and variance of X?								
	Х	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		
a) 2,	, 4/3	b) 3	, 4/3	(c) 2, 2/3		d) 3, 2/3	

47. Find the expectation of a random variable X?

	х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) (0.5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2
- d) \sqrt{npq}