

# CS685: DATA MINING MODEL-BASED CLUSTERING METHODS

Arnab Bhattacharya  
arnabb@cse.iitk.ac.in

Computer Science and Engineering,  
Indian Institute of Technology, Kanpur  
<http://web.cse.iitk.ac.in/~cs685/>

1<sup>st</sup> semester, 2020-21  
Mon 1030-1200 (online)

# Probabilistic Model-Based Clustering

- Assume a model for each cluster
- Observed data points are just samples from these models

# Probabilistic Model-Based Clustering

- Assume a model for each cluster
- Observed data points are just samples from these models
- Dataset  $D$  consists of  $k$  clusters,  $C_1, \dots, C_k$
- Each cluster has a *prior* probability  $\omega_j$  that captures its background probability
- Assuming that there are only  $k$  clusters,  $\sum_{j=1}^k \omega_j = 1$
- This constitutes a **mixture model**

# Probabilistic Theory of Clustering

- Probability density function in each  $C_j$  is given by  $f_j$

# Probabilistic Theory of Clustering

- Probability density function in each  $C_j$  is given by  $f_j$
- A data point  $O_i$  is generated from a cluster  $C_j$  with probability

$$P(O_i|C_j) = f_j(O_i)$$

# Probabilistic Theory of Clustering

- Probability density function in each  $C_j$  is given by  $f_j$
- A data point  $O_i$  is generated from a cluster  $C_j$  with probability

$$P(O_i|C_j) = f_j(O_i)$$

- $O_i$  is generated from the mixture model  $C = (C_1, \dots, C_k)$  with probability

$$P(O_i|C) = \sum_{j=1}^k (\omega_j \cdot f_j(O_i))$$

# Probabilistic Theory of Clustering

- Probability density function in each  $C_j$  is given by  $f_j$
- A data point  $O_i$  is generated from a cluster  $C_j$  with probability

$$P(O_i|C_j) = f_j(O_i)$$

- $O_i$  is generated from the mixture model  $C = (C_1, \dots, C_k)$  with probability

$$P(O_i|C) = \sum_{j=1}^k (\omega_j \cdot f_j(O_i))$$

- The entire dataset  $D$  is generated from  $C$  with probability (assuming all  $O_i$ 's to be independent)

$$P(D|C) = \prod_{i=1}^n P(O_i|C) = \prod_{i=1}^n \left[ \sum_{j=1}^k (\omega_j \cdot f_j(O_i)) \right]$$

# Probabilistic Theory of Clustering

- Probability density function in each  $C_j$  is given by  $f_j$
- A data point  $O_i$  is generated from a cluster  $C_j$  with probability

$$P(O_i|C_j) = f_j(O_i)$$

- $O_i$  is generated from the mixture model  $C = (C_1, \dots, C_k)$  with probability

$$P(O_i|C) = \sum_{j=1}^k (\omega_j \cdot f_j(O_i))$$

- The entire dataset  $D$  is generated from  $C$  with probability (assuming all  $O_i$ 's to be independent)

$$P(D|C) = \prod_{i=1}^n P(O_i|C) = \prod_{i=1}^n \left[ \sum_{j=1}^k (\omega_j \cdot f_j(O_i)) \right]$$

- Clustering: find  $C$  (along with its parameters) that *maximize*  $P(D|C)$



# Expectation-Maximization (EM)

- How to find the parameters of the clusters?
- May not be analytically solvable or may be computationally intractable

# Expectation-Maximization (EM)

- How to find the parameters of the clusters?
- May not be analytically solvable or may be computationally intractable
- **Expectation-maximization (EM)** framework
- General framework to solve many parameter estimation problems
- Two important steps

# Expectation-Maximization (EM)

- How to find the parameters of the clusters?
- May not be analytically solvable or may be computationally intractable
- **Expectation-maximization (EM)** framework
- General framework to solve many parameter estimation problems
- Two important steps
- **Expectation step (E-step)**: Given the current estimate of the model parameters, distribute (may be probabilistically) the data to models
  - Distribute according to expectation

# Expectation-Maximization (EM)

- How to find the parameters of the clusters?
- May not be analytically solvable or may be computationally intractable
- **Expectation-maximization (EM)** framework
- General framework to solve many parameter estimation problems
- Two important steps
- **Expectation step (E-step)**: Given the current estimate of the model parameters, distribute (may be probabilistically) the data to models
  - Distribute according to expectation
- **Maximization step (M-step)**: Given the current distribution of data to models, update the parameters of the models to maximize the generation of the data
  - Maximize according to data

# Expectation-Maximization (EM)

- How to find the parameters of the clusters?
- May not be analytically solvable or may be computationally intractable
- **Expectation-maximization (EM)** framework
- General framework to solve many parameter estimation problems
- Two important steps
- **Expectation step (E-step)**: Given the current estimate of the model parameters, distribute (may be probabilistically) the data to models
  - Distribute according to expectation
- **Maximization step (M-step)**: Given the current distribution of data to models, update the parameters of the models to maximize the generation of the data
  - Maximize according to data
- Keep iterating

# Expectation-Maximization (EM)

- How to find the parameters of the clusters?
- May not be analytically solvable or may be computationally intractable
- **Expectation-maximization (EM)** framework
- General framework to solve many parameter estimation problems
- Two important steps
- **Expectation step (E-step)**: Given the current estimate of the model parameters, distribute (may be probabilistically) the data to models
  - Distribute according to expectation
- **Maximization step (M-step)**: Given the current distribution of data to models, update the parameters of the models to maximize the generation of the data
  - Maximize according to data
- Keep iterating
- Mostly gets stuck in local maxima

# EM for Clustering

- *Expectation step (E-step)*: Given the current cluster parameters, assign each data point to the cluster that is most likely to generate it

# EM for Clustering

- *Expectation step (E-step)*: Given the current cluster parameters, assign each data point to the cluster that is most likely to generate it
- *Maximization step (M-step)*: Given the current assignment of data points to a cluster, update cluster parameters such that likelihood of points generated from it is maximum



# EM for Clustering

- *Expectation step (E-step)*: Given the current cluster parameters, assign each data point to the cluster that is most likely to generate it
- *Maximization step (M-step)*: Given the current assignment of data points to a cluster, update cluster parameters such that likelihood of points generated from it is maximum
- K-means can be thought of as a type of EM algorithm
  - E-step: Assign a data point to the nearest cluster centre
  - M-step: Update cluster centre to minimize SSE of its points

# EM for Mixture Models

- Assume univariate Gaussian mixture models
- Set of parameters  $\theta = \{\theta_1, \dots, \theta_k\}$ , each  $\theta_j = (\mu_j, \sigma_j)$  for Gaussian
- Dataset  $D = \{O_1, \dots, O_n\}$

# EM for Mixture Models

- Assume univariate Gaussian mixture models
- Set of parameters  $\theta = \{\theta_1, \dots, \theta_k\}$ , each  $\theta_j = (\mu_j, \sigma_j)$  for Gaussian
- Dataset  $D = \{O_1, \dots, O_n\}$
- Start with random  $\theta_j$ 's

# EM for Mixture Models

- Assume univariate Gaussian mixture models
- Set of parameters  $\theta = \{\theta_1, \dots, \theta_k\}$ , each  $\theta_j = (\mu_j, \sigma_j)$  for Gaussian
- Dataset  $D = \{O_1, \dots, O_n\}$
- Start with random  $\theta_j$ 's
- *E-step*: For each point  $O_i$ , probability that  $O_i$  is assigned to cluster  $C_j$  is

$$P(\theta_j|O_i, \theta) = \frac{P(O_i|\theta_j)}{\sum_{l=1}^k P(O_i|\theta_l)}$$

# EM for Mixture Models

- Assume univariate Gaussian mixture models
- Set of parameters  $\theta = \{\theta_1, \dots, \theta_k\}$ , each  $\theta_j = (\mu_j, \sigma_j)$  for Gaussian
- Dataset  $D = \{O_1, \dots, O_n\}$
- Start with random  $\theta_j$ 's
- *E-step*: For each point  $O_i$ , probability that  $O_i$  is assigned to cluster  $C_j$  is

$$P(\theta_j|O_i, \theta) = \frac{P(O_i|\theta_j)}{\sum_{l=1}^k P(O_i|\theta_l)}$$

- *M-step*: For each cluster  $C_j$ , adjust  $\theta_j$  such that expected likelihood of points, i.e.,  $P(O|\theta)$  is maximized

$$\mu_j = \frac{1}{k} \sum_{i=1}^n O_i \frac{P(\theta_j|O_i, \theta)}{\sum_{l=1}^n P(\theta_j|O_l, \theta)} = \frac{1}{k} \frac{\sum_{i=1}^n O_i \cdot P(\theta_j|O_i, \theta)}{\sum_{i=1}^n P(\theta_j|O_i, \theta)}$$
$$\sigma_j = \sqrt{\frac{\sum_{i=1}^n (O_i - \mu_j)^2 \cdot P(\theta_j|O_i, \theta)}{\sum_{i=1}^n P(\theta_j|O_i, \theta)}}$$