

The universal property of infinite direct sums in C*-categories and W*-categories

Tobias Fritz and Bas Westerbaan

ABSTRACT. When formulating universal properties for objects in a dagger category, one usually expects a universal property to characterize the universal object up to unique *unitary* isomorphism. We observe that this is automatically the case in the important special case of C*-categories if one uses enrichment in Banach spaces. We then formulate such a universal property for infinite direct sums in C*-categories, and prove the equivalence with the existing definition due to Ghez, Lima and Roberts in the case of W*-categories. This notion specializes to the standard infinite direct sums in the case of a W*-category of normal representations of a W*-algebra.

Finding a universal property for the more general case of direct integrals remains an open problem.

1. Introduction

Among the most central and useful concepts of category theory is the notion of universal property. Its utility relates to the fact that a universal property characterizes an object up to unique isomorphism, and to the vast number of mathematical construction which can be defined or characterized in terms of universal properties. Among these is the concept of *direct sum* of two objects A_1 and A_2 in an additive category \mathcal{C} : the direct sum $A_1 \oplus A_2$, which can be defined in several equivalent ways via the following standard result:

1.1. THEOREM (e.g. [1, Proposition 1.2.4]). *Let \mathcal{C} be an additive category and $A_1, A_2 \in \mathcal{C}$ two objects. Then the following pieces of structure are equivalent for an object $A \in \mathcal{A}$:*

(a) *A represents the functor*

$$\mathcal{C} \longrightarrow \mathbf{Ab}, \quad B \longmapsto \mathcal{C}(A_1, B) \times \mathcal{C}(A_2, B)$$

(b) *A represents the functor*

$$\mathcal{C}^{\text{op}} \longrightarrow \mathbf{Ab}, \quad B \longmapsto \mathcal{C}(A_1, B) \times \mathcal{C}(A_2, B)$$

2010 *Mathematics Subject Classification*. Primary: ; Secondary:

Acknowledgements. We thank Robert Furber for copious help and discussion. Part of this work was conducted while the first author was at the Max Planck Institute for Mathematics in the Sciences in Leipzig, Germany.

- (c) *There are morphisms $p_i : A \rightarrow A_i$ and $\kappa_i : A_i \rightarrow A$ satisfying the biproduct equations*

$$p_i \kappa_j = \delta_{ij}, \quad \kappa_1 p_1 + \kappa_2 p_2 = 1.$$

Here, (a) and (b) state that A is a coproduct and product of A_1 and A_2 , respectively. (c) is a purely equational characterization which also proves that direct sums are preserved by every additive functor.

Traditionally, universal properties have been formulated mostly for purely algebraic structures only. Treating analytical structures categorically is straightforward in some cases, but very challenging in many other cases. It usually involves equipping the categories under consideration themselves with certain analytical structures. An interesting case of where this happens is the theory of W^* -categories, which model categories of Hilbert space representations of C^* -algebras or W^* -algebras, with the most paradigmatic example of a W^* -category being Hilb , the category of Hilbert spaces with bounded linear maps.

In this paper, we study *infinite* direct sums of objects in W^* -categories and in the more general case of C^* -categories. (Our definitions and results also apply to the case of finite direct sums, but reduce to Theorem 1.1 in this case.) As the most basic example, the Hilbert space $\ell^2(I)$ for a (finite or) infinite set I is the direct sum of I many copies of \mathbb{C} . Part of the *raison d'être* for analytic structure on an algebraic object is that it often makes the objects under consideration more well-behaved. This is indeed what happens in our case as well: while Theorem 1.1 does not generalize to infinite direct sums since infinite coproducts and infinite products do no longer coincide in a general additive category (even when they exist), our main result (Theorem 5.1) provides such a generalization for infinite direct sums in W^* -categories.

Summary. After giving some background on C^* -categories and W^* -categories in Section 2, we start by investigating universal properties for objects in C^* -categories in Section 3. We find that defining universal properties in terms of representability of a functor enriched in Banach spaces provides a good notion of universal property in a C^* -category, since it guarantees uniqueness up to unique *unitary* isomorphism of the representing object. We then introduce our definition of the particular universal property of infinite direct sums in C^* -categories in Section 4. In Section 5, we specialize to the case of W^* -categories, and prove an analog of Theorem 1.1 in this setting, showing that our infinite direct sums are equivalent to the direct sums in W^* -categories as introduced originally by Ghez, Lima and Roberts [3].

Direct integrals. We have also attempted to formulate a similar universal property for *direct integrals* in W^* -categories, but we have not been able to make this idea work. As has been argued by Simon Henry¹, finding such a universal property requires the consideration of additional structure on the W^* -category. Our current best bet is that doing so will involve a notion of *W^* -category with rigged objects*, generalizing rigged Hilbert spaces [2].

¹See mathoverflow.net/questions/313158/what-does-it-mean-for-a-category-to-admit-direct-integrals.

2. C*-categories and W*-categories

State the basic definitions

2.1. EXAMPLE. If A is a C*-algebra, then the category of non-degenerate representations $\pi : A \rightarrow \mathcal{B}(\mathcal{H})$ on Hilbert spaces \mathcal{H} is a W*-category. The morphisms $(\mathcal{H}_1, \pi_1) \rightarrow (\mathcal{H}_2, \pi_2)$ are exactly the *intertwiners*, i.e. the bounded linear maps $f : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ such that the diagram

$$\begin{array}{ccc} \mathcal{H}_1 & \xrightarrow{f} & \mathcal{H}_2 \\ \pi_1(a) \downarrow & & \downarrow \pi_2(a) \\ \mathcal{H}_1 & \xrightarrow{f} & \mathcal{H}_2 \end{array}$$

commutes for all $a \in A$. It is well-known that this is a W*-category [3, Example 2.4], where the Banach space structure on each hom-set is the usual one on a space of operators using the operator norm, and the involution is given by the usual Hermitian adjoint. We denote this W*-category by $\text{Rep}(A)$. In the most basic case $A = \mathbb{C}$, we get $\text{Rep}(\mathbb{C}) \cong \text{Hilb}$, the category of Hilbert spaces.

2.2. EXAMPLE. If N is a W*-algebra, then the category of normal unital representations on Hilbert spaces is a W*-category, with intertwiners as morphisms as above. We denote this W*-category by $\text{nRep}(N)$. For a C*-algebra A and A^{**} its universal enveloping W*-algebra, we have $\text{nRep}(A^{**}) \cong \text{Rep}(A)$, since non-degenerate representations of A extend uniquely to normal unital representations of A^{**} .

3. Universal properties in C*-categories

In this section, we write \mathcal{C} for a C*-category [3]. We start with a generalization of [7, Lemma 7] to C*-categories before discussing universal properties.

3.1. LEMMA. *Let $A, B \in \mathcal{C}$. Then for any $a : A \rightarrow B$ with $\|a\| \leq 1$ and projections $s : A \rightarrow A$, $t : B \rightarrow B$, the following are equivalent:*

- (a) $a^*ta \leq 1 - s$,
- (b) $asa^* \leq 1 - t$,
- (c) $tas = 0$,
- (d) $sa^*t = 0$.

PROOF. We will first prove $a^*ta \leq 1 - s \Leftrightarrow tas = 0$. Assume $a^*ta \leq 1 - s$. Then $0 \leq sa^*tas \leq s(1 - s)s = 0$. Thus $sa^*tas = 0$, which is to say $tas = 0$. Now, assume $tas = 0$. Then $ta = ta(1 - s)$ and $a^*t = (1 - s)a^*t$. Hence $a^*ta = (1 - s)a^*ta(1 - s) \leq 1 - s$, as desired. A similar argument proves $asa^* \leq 1 - t \Leftrightarrow sa^*t = 0$. To finish the proof, it is sufficient to show $tas = 0 \Leftrightarrow sa^*t = 0$, which follows directly by applying $(\)^*$. \square

3.2. LEMMA. *Let $a : A \hookrightarrow B : b$ be any two morphisms such that $\|a\| \leq 1$, $\|b\| \leq 1$ and $ab = 1$. Then b is an isometry with $a = b^*$.*

PROOF. To start, note $\|a^*a\| = \|a\|^2 \leq 1$, so $a^*a \leq 1$. Similarly $b^*b \leq 1$. Combined $1 = b^*a^*ab \leq b^*b \leq 1$, hence $b^*b = 1$. Similarly $aa^* = 1$. Clearly $bb^*bb^* = b1b^* = bb^*$, so b is an isometry. Now $a(1 - bb^*)a^* = aa^* - abb^*a^* = 1 - 1 = 0$. So $a^*(1 - 0)a \leq bb^*$ by Lemma 3.1. Similarly $bb^* \leq a^*a$. Hence $bb^* = a^*a$. We are done: $a = aa^*a = abb^* = b^*$. \square

We will need the full power of this observation only in the next section. However, the following special case is generally useful when working with universal properties in C^* -categories:

3.3. COROLLARY. *Let $u : A \rightarrow B$ be an invertible morphism with $\|u\| \leq 1$ and $\|u^{-1}\| \leq 1$. Then u is unitary.*

Now let \mathbf{Ban} be the category of Banach spaces with bounded linear maps of norm ≤ 1 . Every C^* -category is \mathbf{Ban} -enriched. Our notion of universal property in a C^* -category is the standard one for enriched categories [5, Section 1.10], namely with enrichment in \mathbf{Ban} :

3.4. DEFINITION. *A functor $F : C \rightarrow \mathbf{Ban}$ is representable if there is an object $A \in C$ together with a natural isomorphism $C(A, -) \cong F$ that is a componentwise isometric isomorphism of Banach spaces.*

Corollary 3.3 and the (weak) enriched Yoneda lemma [5, Section 1.9] imply:

3.5. COROLLARY. *For any representable functor $C \rightarrow \mathbf{Ban}$, the representing object is unique up to unique unitary isomorphism*

PROOF. If both A and B represent F via $C(A, -) \cong F \cong C(B, -)$, then the enriched Yoneda lemma gives us an isomorphism $u : A \rightarrow B$. Since the natural isomorphism between the hom-functors is required to be a componentwise isometry by \mathbf{Ban} -enrichment, we necessarily have $\|u\| = \|u^{-1}\| = 1$, making u into a unitary by Corollary 3.3. \square

This uniqueness up to unique unitary is a desideratum of a good notion of universal property for C^* -categories, and more generally for all dagger categories [4]. Corollary 3.3 tells us that the compatibility with the dagger structure is automatic thanks to the enrichment in \mathbf{Ban} , so that we do not need to formulate our universal properties in the form of dagger limits as in [4].

4. The universal property of infinite direct sums

T: I've generalized the defn of direct sum from C^* -categories, where it works just the same way as for W^* -categories, except for Theorem 5.1. Does this seem reasonable?

We now turn to the study of infinite direct sums by defining them in terms of a universal property as in Definition 3.4, and then proving that the existing notions of direct sum in the category of Hilbert spaces and more generally all W^* -categories indeed satisfy this universal property. While our considerations also apply to finite direct sums as a special case, in this case our discussion provides nothing new, since then our universal property reduces to the standard one of biproducts in an additive category.

We again let \mathcal{C} be a C^* -category. Let I be a finite or infinite index set. For $(A_i)_{i \in I}$ a family of objects in \mathcal{C} , we consider the functor

$$\bigoplus_i \mathcal{C}(A_i, -) : \mathcal{C} \rightarrow \mathbf{Ban}$$

given by associating to every object B the set

$$\bigoplus_i \mathcal{C}(A_i, B) := \left\{ (f_i : A_i \rightarrow B)_{i \in I} \mid \sum_i f_i f_i^* < \infty \right\}. \quad (4.1)$$

Here, $\sum_i f_i f_i^*$ can be interpreted as converging in the ultraweak topology² in the case where $\mathcal{C}(B, B)$ is a W^* -algebra. If this is not the case, what we mean by the condition $\sum_i f_i f_i^* < \infty$ is that there is a fixed element which upper bounds all partial sums $\sum_{i \in S} f_i f_i^*$ for finite $S \subseteq I$, or that the sum converges ultraweakly in the double dual W^* -algebra $\mathcal{C}(B, B)^{**}$.

The set (4.1) is a vector space under componentwise addition and scalar multiplication, where closure under addition follows from $f_x g_x^* + g_x f_x^* \leq f_x f_x^* + g_x g_x^*$. We furthermore consider this vector space as a normed space under the norm

$$\|(f_i)_{i \in I}\| := \left\| \sum_i f_i f_i^* \right\|^{1/2},$$

where the triangle inequality again follows from $f_x g_x^* + g_x f_x^* \leq f_x f_x^* + g_x g_x^*$, and we interpret the sum as before either in terms of a supremum over the norm of finite partial sums, or as converging in the double dual $\mathcal{C}(B, B)^{**}$.

4.1. LEMMA. *Let \mathcal{C} be a C^* -category, $(A_i)_I \in \mathcal{C}$ a family of objects and $B \in \mathcal{C}$. Then $\bigoplus_i \mathcal{C}(A_i, B)$ is complete, and therefore a Banach space.*

The proof generalizes the standard proof showing that $\ell^2(I)$ is a Hilbert space.

PROOF. Suppose that we are given a Cauchy sequence of families $(f^{(n)})_{n \in \mathbb{N}} = ((f_i^{(n)})_{i \in I})_{n \in \mathbb{N}}$. For given j , the projection map

$$\bigoplus_i \mathcal{C}(A_i, B) \longrightarrow \mathcal{C}(A_j, B)$$

is bounded, and therefore $(f_j^{(n)})_{n \in \mathbb{N}}$ is Cauchy as well; let us denote the limit by $f_j : A_j \rightarrow B$. We will now show that $f \in \bigoplus_i \mathcal{C}(A_i, B)$ for $f := (f_j)_{j \in I}$, which amounts to verifying the relevant square summability. The Cauchy assumption implies that the sequence $(f^{(n)})_{n \in \mathbb{N}}$ is uniformly bounded by some constant C ,

$$\left\| \sum_i f_i^{(n)} (f_i^{(n)})^* \right\|^{1/2} \leq C.$$

²For positive elements $(x_i)_{i \in I}$ of a W^* -algebra, convergence and value of the sum $\sum_{i \in I} x_i$ coincides in the ultraweak and ultrastrong topology and, in fact, equals the supremum of the partial sums.

Restricting to a finite partial sum over $S \subseteq I$ and taking $n \rightarrow \infty$ shows that

$$\left\| \sum_{i \in S} f_i f_i^* \right\|^{1/2} \leq C$$

as well. Since S was arbitrary, we conclude that $\|f\| \leq C$.

The convergence $f_i \rightarrow f$ works similarly. By assumption, for every $\varepsilon > 0$ we have $\|f^{(n)} - f^{(m)}\| \leq \varepsilon$ for almost all $n, m \in \mathbb{N}$. In particular, we know that for any finite $S \subseteq I$,

$$\left\| \sum_{i \in S} (f_i^{(n)} - f_i^{(m)})^* (f_i^{(n)} - f_i^{(m)}) \right\|^{1/2} \leq \varepsilon$$

for almost all $n, m \in \mathbb{N}$. Taking $m \rightarrow \infty$ gives

$$\left\| \sum_{i \in S} (f_i^{(n)} - f_i)^* (f_i^{(n)} - f_i) \right\|^{1/2} \leq \varepsilon.$$

Since S was arbitrary, we conclude $\|f^{(n)} - f\| \leq \varepsilon$ for almost all n . \square

The map $B \mapsto \bigoplus_i \mathbb{C}(A_i, B)$ is a functorial in B in the obvious way, giving a functor

$$\bigoplus_i \mathbb{C}(A_i, -) : \mathbb{C} \rightarrow \mathbf{Ban}.$$

We now get to our main definition:

4.2. DEFINITION. *Let \mathbb{C} be a C^* -category. An I -indexed direct sum of the family of objects $(A_i)_{i \in I}$ is an object $\bigoplus_i A_i$ which represents the \mathbf{Ban} -enriched functor $\bigoplus_i \mathbb{C}(A_i, -)$.*

By Corollary 3.5, this characterizes infinite direct sums up to unique unitaries. Since \mathbb{C} is a dagger category, the same holds dually, meaning that $\bigoplus_i A_i$ also represents the functor

$$\bigoplus_i \mathbb{C}(-, A_i) : \mathbb{C}^{\text{op}} \rightarrow \mathbf{Ban}$$

given by associating to every object B the Banach space

$$\bigoplus_i \mathbb{C}(B, A_i) := \left\{ (f_i : B \rightarrow A_i)_{i \in I} \mid \sum_i f_i^* f_i < \infty \right\},$$

where the sum is, as before, taken in the ultraweak topology of $\mathbb{C}(B, B)^{**}$ or to be understood in terms of uniform bounds on finite partial sums, and again considered a Banach space with respect to the norm

$$\|(f_i)_{i \in I}\| := \left\| \sum_i f_i^* f_i \right\|^{1/2}.$$

This Banach space is contravariantly functorial in B in the obvious way.

4.3. REMARK. For finite I , our direct sums are also dagger limits in the sense of [4]. For infinite I , this is no longer the case, since infinite direct sums are not even limits; compare the degeneracy result [4, Theorem 5.2], which indicates that infinite dagger products cannot be expected to exist. In this sense, our notion of universal property via Definition 4.2 seems better adapted to the category of Hilbert spaces or other W^* -categories than that of [4].

4.4. REMARK. We could also consider *direct sums with kernel* in the following sense. Let $K = (K_{ij}) \in \mathbb{C}^{I \times I}$ be a positive semidefinite matrix. Then given a family of objects $(A_i)_{i \in I}$ as before, we can consider the functor

$$\mathcal{C} \longrightarrow \text{Ban}, \quad B \longmapsto \bigoplus_i \mathcal{C}(A_i, B) := \left\{ (f_i : A_i \rightarrow B)_{i \in I} \mid \sum_{i,j} K_{ij} f_i f_j^* < \infty \right\},$$

where the completeness proof with respect to the norm $f \mapsto \|\sum K_{ij} f_i f_j^*\|^{1/2}$ works in the same way as the proof of Lemma 4.1.

We can thus define the *direct sum with kernel* K to be an object representing this functor in the same way as above. For example in Hilb with $A_i = \mathbb{C}$ for all $i \in I$, the resulting direct sum would be the space of families $(\xi_i)_{i \in I}$ with $\xi_i \in \mathbb{C}$ that are K -square summable in that $\sum_{i,j} K_{ij} \xi_i \bar{\xi}_j < \infty$, turned into a Hilbert space with respect to the inner product with kernel K .

T: Is it worth elaborating on this and prove a generalization of Theorem 5.1?

5. Direct sums in W^* -categories

In a W^* -category, we characterize the existence of direct sums as follows:

5.1. THEOREM. *Let \mathcal{C} be a W^* -category and $(A_i)_{i \in I}$ a family of objects in \mathcal{C} . Then the following pieces of structure are equivalent for an object A :*

- (a) *A is an I -indexed direct sum $\bigoplus_i A_i$.*
- (b) *There is a family of morphisms $(\kappa_j : A_j \rightarrow A)_{j \in I}$ with $\sum_{j \in I} \kappa_j \kappa_j^* < \infty$ such that for any object B and family of morphisms $(f_j : A_j \rightarrow B)_{j \in I}$ with $\sum_{j \in I} f_j f_j^* < \infty$, there is a unique $f : A \rightarrow B$ such that $f \kappa_j = f_j$ for all $j \in I$, and $\|f\|^2 = \|\sum_{j \in I} f_j f_j^*\|$.*
- (c) *There is a family of morphisms $(\kappa_j : A_j \rightarrow A)_{j \in I}$ such that $\kappa_i^* \kappa_j = \delta_{ij}$, and*

$$\sum_{j \in I} \kappa_j \kappa_j^* = 1_A. \tag{5.1}$$

As far as we know, the first definition of infinite direct sums in W^* -categories has been given in [3, p. 100] in the form (c). From our perspective, it is preferable to replace this definition by a universal property as in Definition 4.2, and to regard the original definition of Ghez, Lima and Roberts as an alternative characterization instead.

PROOF. (b) is a simple reformulation of the universal property (a) via the Yoneda lemma, as follows. Given a natural isometric isomorphism $\mathcal{C}(A, B) \cong \bigoplus_i \mathcal{C}(A_i, B)$, we

define the family (κ_i) to be the image of 1_A . Then for any $f : A \rightarrow B$, the naturality diagram

$$\begin{array}{ccc} \mathcal{C}(A, A) & \xrightarrow{\cong} & \bigoplus_i \mathcal{C}(A_i, A) \\ \downarrow f \circ - & & \downarrow (f \circ -) \\ \mathcal{C}(A, B) & \xrightarrow{\cong} & \bigoplus_i \mathcal{C}(A_i, B) \end{array}$$

shows that the universal property takes f to the family $(f_j) := (f\kappa_j)$. By the definition of the norm on $\bigoplus_i \mathcal{C}(A_i, B)$ and the assumption that the universal property is an isometric isomorphism, we indeed have $\|f\|^2 = \left\| \sum_j f_j f_j^* \right\|$, which also gives $\sum_j \kappa_j \kappa_j^* < \infty$ in the case $f = 1_A$. Conversely, it is clear that any family (κ_j) as in (b) gives an isometric isomorphism $\mathcal{C}(A, B) \cong \bigoplus_i \mathcal{C}(A_i, B)$ that is natural in B .

We now derive (c) from (b). Applying the assumption on norms with $(f_j) := (\kappa_j)$ gives $\left\| \sum_i \kappa_i \kappa_i^* \right\| = \|1_A\|$, and therefore $\|\kappa_j\| \leq 1$ for all $j \in I$. For fixed j , the family $(\delta_{ij} : A_i \rightarrow A_j)$ corresponds to some morphism $\pi_j : A \rightarrow A_j$ such that $\pi_j \circ \kappa_i = \delta_{ij}$. Again by the isometry property, we have $\|\pi_j\| = \|1_A\| \leq 1$. By Lemma 3.2, this implies $\pi_j = \kappa_j^*$. We therefore have $\kappa_j^* \kappa_i = \delta_{ij}$, as desired. This also implies that the $\kappa_j \kappa_j^*$ are mutually orthogonal projections. In order to also prove the completeness relation (5.1), let $p : A \rightarrow A$ be any other projection that is orthogonal to each of the $\kappa_j \kappa_j^*$; we need to show $p = 0$. Again by naturality, under the isomorphism $\mathcal{C}(A, A) \cong \bigoplus_i \mathcal{C}(A_i, A)$, the new p corresponds to the family $(p\kappa_i) = (0)$, where $p\kappa_i = 0$ follows from the assumed orthogonality. This implies indeed $p = 0$ by the uniqueness part of the universal property.

We finally show that (c) implies (b). So let $(f_j : A_j \rightarrow B)_{j \in I}$ be any family of morphisms with $\sum_{j \in I} f_j f_j^* < \infty$. The homset $\mathcal{C}(B, A)$ is a self-dual Hilbert $\mathcal{C}(B, B)$ -module with $\mathcal{C}(B, B)$ -valued inner product $\langle g, f \rangle \equiv g^* f$ [3, Proposition 2.15]. The self-duality is equivalent to the fact that $\mathcal{C}(B, A)$ is complete in the *ultranorm topology*; this is the topology generated by the seminorms $\|f\|_\omega \equiv \omega(\langle f, f \rangle)^{\frac{1}{2}}$ indexed by normal states ω on $\mathcal{C}(B, B)$ [8, §149V], also known as the *s-topology* [3, p. 88]. We now turn to the construction of $f : A \rightarrow B$. Pick any normal state ω on $\mathcal{C}(B, B)$ and finite subset $S \subseteq I$. We have

$$\begin{aligned} \left\| \sum_{j \in S} \kappa_j f_j^* \right\|_\omega^2 &\equiv \omega \left(\left(\sum_{i \in S} \kappa_i f_i^* \right)^* \left(\sum_{j \in S} \kappa_j f_j^* \right) \right) \\ &= \omega \left(\sum_{j \in S} f_j f_j^* \right) \quad \text{as } \kappa_j^* \kappa_i = \delta_{ij}. \end{aligned}$$

From this and $\sum_{j \in I} f_j f_j^* < \infty$, it follows that the net $\left(\sum_{j \in S} \kappa_j f_j^* \right)_S$ is ultranorm Cauchy as S ranges over all finite subsets of I . Define $f := \left(\sum_{j \in I} \kappa_j f_j^* \right)^*$, where the sum is understood to converge ultranorm. The assignment $g \mapsto \kappa_j^* g$ yields a bounded $\mathcal{C}(B, B)$ -linear map between the Hilbert $\mathcal{C}(B, B)$ -modules $\mathcal{C}(B, A)$ and $\mathcal{C}(B, A_j)$ and is therefore ultranorm continuous [8, §148]. Hence $\kappa_j^* f^* = \kappa_j^* \sum_i \kappa_i f_i^* = \sum_i \kappa_j^* \kappa_i f_i^* = f_j^*$ and so $f \kappa_j =$

f_j as desired. Next we will show that $\|f\|^2 = \|\sum_j f_j f_j^*\|$. By definition of f^* and [8, §148V] we have

$$\langle f^*, f^* \rangle = \text{uwlim}_{S \subseteq I \text{ finite}} \left\langle \sum_{j \in S} \kappa_j f_j^*, \sum_{j \in S} \kappa_j f_j^* \right\rangle = \text{uwlim}_{S \subseteq I \text{ finite}} \sum_{j \in S} f_j f_j^* = \sum_j f_j f_j^*$$

and so $\|f\|^2 = \|f^*\|^2 = \|\langle f, f \rangle\| = \|\sum_j f_j f_j^*\|$ as desired. Finally, to show uniqueness, assume $f': A \rightarrow B$ is any (other) morphism with $f' \kappa_i = f_i$. Note that $\sum_j \kappa_j \kappa_j^*$ converges ultrastrongly and so ultranorm as well. With similar reasoning as before, the composition map $g \mapsto f'g$ is ultranorm continuous, hence

$$f' = f' \sum_{j \in I} \kappa_j \kappa_j^* = \sum_{j \in I} f' \kappa_j \kappa_j^* = \sum_{j \in I} f_j \kappa_j^* = \sum_{j \in I} f \kappa_j \kappa_j^* = f,$$

where the sums converge ultranorm. \square

It is now straightforward to note:

5.2. COROLLARY. *Every normal *-functor between W*-categories preserves direct sums.*

Rieffel has observed this for the W*-categories $\text{Rep}(A)$ from Example 2.1 as [6, Proposition 4.9]. The following was undoubtedly already known to the authors of [3], but we have not found it stated anywhere explicitly. We are now in a position to give a very simple proof.

5.3. PROPOSITION. *Let N be a W*-algebra. In the W*-category $\text{nRep}(N)$, any ℓ^2 -direct sum*

$$\bigoplus_{i \in I} A_i := \left\{ (\xi_i)_{i \in I} \mid \xi_i \in A_i, \sum_i \langle \xi_i, \xi_i \rangle < \infty \right\}, \quad (5.2)$$

equipped with the componentwise representation of N , is an I -indexed direct sum in the sense of Definition 4.2. Conversely, every I -indexed direct sum in $\text{nRep}(N)$ is of this form.

PROOF. Given a concrete direct sum as in (5.2), it is straightforward to see that the inclusion maps $\kappa_i : A_i \rightarrow A$ satisfy the conditions of Theorem 5.1(c). This shows that all direct sums in $\text{nRep}(N)$ exist; the uniqueness up to unique unitaries therefore implies that every direct sum is of this form. \square

References

- [1] Francis Borceux. *Handbook of categorical algebra. 2*, volume 51 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1994. Categories and structures. \uparrow 1
- [2] I. M. Gel'fand and N. Ya. Vilenkin. *Generalized functions. Vol. 4: Applications of harmonic analysis*. Translated by Amiel Feinstein. Academic Press, New York - London, 1964, 1964. \uparrow 2
- [3] P. Ghez, R. Lima, and J.E. Roberts. W*-categories. *Pacific Journal of Mathematics*, 120:79–109, 1985. \uparrow 2, 3, 7, 8, 9
- [4] Chris Heunen and Martti Karvonen. Limits in dagger categories, 2018. [arXiv:1803.06651](https://arxiv.org/abs/1803.06651). \uparrow 4, 7
- [5] G. M. Kelly. Basic concepts of enriched category theory. *Repr. Theory Appl. Categ.*, (10), 2005. Reprint of the 1982 original [Cambridge Univ. Press, Cambridge]. \uparrow 4

- [6] Marc A. Rieffel. Morita equivalence for C^* -algebras and W^* -algebras. *Journal of Pure and Applied Algebra*, 5(1):51–96, 1974. [↑9](#)
- [7] Abraham Westerbaan and Bas Westerbaan. A universal property for sequential measurement. *Journal of Mathematical Physics*, 57(9):092203, 2016. [↑3](#)
- [8] Bas Westerbaan. *Dagger and dilations in the category of von Neumann algebras*. PhD thesis, Radboud University, 2018. [arXiv:1803.01911](#). [↑8](#), [9](#)

PERIMETER INSTITUTE FOR THEORETICAL PHYSICS
Email address: tfritz@pitp.ca

ADD YOUR AFFILIATION HERE