

Calculus Formulae Sheet

Muktadir Somio

Contents

1	Precalculus	4
1.1	Functions and Notation	4
1.2	Basic Identities	4
1.3	Inverse functions and monotonicity	4
2	Limits	4
2.1	Intuitive notions	4
2.2	Formal epsilon-delta definition	4
2.3	One-sided limits and infinite limits	4
2.4	Standard limit techniques	4
2.5	Common limits	5
2.6	Continuity	5
3	Sequences and Series (Single variable)	5
3.1	Sequences	5
3.2	Infinite series	5
3.3	Tests for convergence	5
3.4	Power series	5
3.5	Taylor and Maclaurin series	5
4	Differential Calculus	6
4.1	Derivatives	6
4.2	Rules of Differentiation	6
4.3	Higher derivatives and Taylor remainder	6
4.4	Mean Value Theorems	6
4.5	Asymptotic notation	6
4.6	Convexity and second derivative test	6
5	Riemann Integration and Measure-theoretic remarks	7
5.1	Riemann sum definition	7
5.2	Riemann integrability criterion	7
5.3	Fundamental Theorems of Calculus	7
5.4	Definite and Indefinite Integrals	7
5.5	Improper integrals	7

6	Integration Techniques	7
6.1	Basic Integration Rules	7
6.2	Techniques of Integration	8
6.3	Trigonometric integrals	8
6.4	Reduction formulas	8
6.5	Special definite integrals	8
6.6	Applications of Integration: Area, Volume, and Density	8
6.7	Integrals in Polar Coordinates	10
7	Special Functions	11
7.1	Gamma function	11
7.2	Beta function	11
7.3	Stirling's approximation	11
7.4	Error function and Gaussian integrals	11
7.5	Bessel, Legendre, orthogonal polynomials (formulas)	11
8	Multivariable Calculus	12
8.1	Functions of several variables	12
8.2	Limits and continuity in \mathbb{R}^n	12
8.3	Vector-Valued Functions	12
8.4	Partial derivatives and differentiability	13
8.5	Gradient, directional derivative, Hessian	13
8.6	Chain rule (multivariable)	13
8.7	Taylor series in several variables	14
8.8	Multiple integrals	14
8.9	Change of variables and Jacobian	14
8.10	Common coordinate systems	14
8.11	Line integrals	14
8.12	Conservative fields and potentials	14
8.13	Surface integrals	14
8.14	Gradient, Divergence, Curl	15
8.15	Integral Theorems	15
8.16	Optimization with constraints	15
9	Ordinary Differential Equations	15
9.1	Classification and existence	15
9.2	Linear first-order ODEs	15
9.3	Separable equations	15
9.4	Second-order linear ODEs with constant coefficients	16
9.5	Method of undetermined coefficients	16
9.6	Variation of parameters (2nd order)	16
9.7	Linear systems and matrix exponentials	16
9.8	Stability of equilibria	16
9.9	Phase plane methods	16
9.10	Boundary value problems and Sturm-Liouville	16
10	Applications	16
10.1	Optimization	16

11 Partial Differential Equations: Classical Examples and Methods	17
11.1 Canonical PDEs	17
11.2 Separation of variables	17
11.3 Fourier series and transforms	17
11.4 Green's functions	17
12 Transforms and Operational Methods	17
12.1 Laplace transform	17
12.2 Z-transform (discrete)	17
12.3 Fourier transform properties	17
13 Numerical Methods	18
13.1 Root finding	18
13.2 Numerical integration	18
13.3 Numerical ODEs	18
14 Miscellaneous Topics	18
14.1 Uniform convergence	18
14.2 Dominated convergence theorem (measure-theory remark)	18
14.3 Asymptotic expansions and stationary phase	18
14.4 Calculus of variations (principles)	18
15 Cheat Sheet: Common Identities and Integrals	19
15.1 Derivatives	19
15.2 Integrals	19
15.3 Useful limits and series	19
16 Reference Theorems (Statements)	19

1 Precalculus

1.1 Functions and Notation

$$f : X \rightarrow Y, \quad \text{dom}(f), \text{ range}(f), \quad f^{-1}(B) = \{x : f(x) \in B\}.$$

Types: polynomial, rational, exponential, logarithmic, trigonometric, inverse trig, hyperbolic.

1.2 Basic Identities

$$\begin{aligned}\sin^2 x + \cos^2 x &= 1, \\ \sin(2x) &= 2 \sin x \cos x, \\ \cos(2x) &= \cos^2 x - \sin^2 x = 2 \cos^2 x - 1 = 1 - 2 \sin^2 x, \\ \tan(A \pm B) &= \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}, \\ \sinh^2 x + 1 &= \cosh^2 x.\end{aligned}$$

1.3 Inverse functions and monotonicity

If f is strictly monotone on interval I then f^{-1} exists on $f(I)$ and $(f^{-1})'(y) = \frac{1}{f'(x)}$ with $y = f(x)$.

2 Limits

2.1 Intuitive notions

Limit describes approaching behavior. Useful heuristics: squeeze theorem, dominant terms for $x \rightarrow \infty$ or $x \rightarrow 0$.

2.2 Formal epsilon-delta definition

Definition 2.1. $\lim_{x \rightarrow a} f(x) = L$ iff $\forall \varepsilon > 0 \exists \delta > 0$ such that $0 < |x - a| < \delta \Rightarrow |f(x) - L| < \varepsilon$.

2.3 One-sided limits and infinite limits

$$\begin{aligned}\lim_{x \rightarrow a^+} f(x), \quad \lim_{x \rightarrow a^-} f(x), \quad \lim_{x \rightarrow a} f(x) = L &\iff \text{both one-sided limits} = L. \\ \lim_{x \rightarrow a} f(x) = \infty &\text{ means } \forall M > 0, \exists \delta > 0 : 0 < |x - a| < \delta \Rightarrow f(x) > M.\end{aligned}$$

2.4 Standard limit techniques

- Algebraic simplification.
- Factor and cancel.
- Multiply by conjugate.
- L'Hôpital's rule: if $\lim f = \lim g = 0$ or $\pm\infty$, then $\lim \frac{f}{g} = \lim \frac{f'}{g'}$ when RHS exists.

- Series expansion.

2.5 Common limits

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1, \quad \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1, \quad \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e.$$

2.6 Continuity

$$f \text{ continuous at } x = a \iff \lim_{x \rightarrow a} f(x) = f(a),$$

Intermediate Value Theorem: f continuous on $[a, b] \Rightarrow \forall k \in [f(a), f(b)], \exists c \in [a, b] : f(c) = k$.

3 Sequences and Series (Single variable)

3.1 Sequences

$$a_n \rightarrow L \iff \forall \varepsilon > 0, \exists N : n > N \Rightarrow |a_n - L| < \varepsilon.$$

Monotone convergence theorem: monotone bounded sequence converges.

3.2 Infinite series

$$\sum_{n=0}^{\infty} a_n \text{ converges} \iff \text{partial sums } s_N = \sum_{n=0}^N a_n \text{ converge.}$$

3.3 Tests for convergence

- **Comparison test:** $0 \leq a_n \leq b_n$ and $\sum b_n$ converges $\Rightarrow \sum a_n$ converges.
- **Limit comparison:** $\lim \frac{a_n}{b_n} = L \in (0, \infty) \Rightarrow$ same behaviour.
- **Ratio test:** $L = \lim |a_{n+1}/a_n|$; $L < 1$ conv, $L > 1$ div, $L = 1$ inconclusive.
- **Root test:** $L = \limsup \sqrt[n]{|a_n|}$ similar to ratio.
- **Alternating series (Leibniz):** if $a_n \downarrow 0$ then $\sum (-1)^n a_n$ converges.
- **Absolute convergence:** $\sum |a_n|$ converges $\Rightarrow \sum a_n$ converges.

3.4 Power series

$$\sum_{n=0}^{\infty} c_n (x - a)^n, \quad R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|c_n|}}.$$

Within radius of convergence series can be differentiated and integrated termwise.

3.5 Taylor and Maclaurin series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n, \quad R_n(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} (x - a)^{n+1}.$$

4 Differential Calculus

4.1 Derivatives

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h},$$
$$(Df)(x) = \frac{df}{dx}, \quad \frac{d}{dx}(c) = 0, \quad \frac{d}{dx}(x^n) = nx^{n-1}.$$

4.2 Rules of Differentiation

$$(uv)' = u'v + uv',$$
$$\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2},$$
$$(f(g(x)))' = f'(g(x))g'(x),$$
$$\frac{d}{dx}[a^x] = a^x \ln a, \quad \frac{d}{dx}[\ln x] = \frac{1}{x},$$
$$\frac{d}{dx}[\sin x] = \cos x, \quad \frac{d}{dx}[\cos x] = -\sin x,$$
$$\frac{d}{dx}[\tan x] = \sec^2 x.$$

4.3 Higher derivatives and Taylor remainder

$$f^{(n)}(x) = \frac{d^n f}{dx^n}.$$

Taylor remainder Lagrange form given above.

4.4 Mean Value Theorems

Rolle's Theorem: $f(a) = f(b)$, $f'(\xi) = 0$ for some $\xi \in (a, b)$,

Lagrange MVT: $f'(\xi) = \frac{f(b) - f(a)}{b - a}$,

Cauchy MVT: $\frac{f'(\xi)}{g'(\xi)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$

4.5 Asymptotic notation

$$f(x) = o(g(x)) \iff \lim_{x \rightarrow a} \frac{f}{g} = 0, \quad f(x) \sim g(x) \iff \lim_{x \rightarrow a} \frac{f}{g} = 1.$$

4.6 Convexity and second derivative test

If $f''(x) > 0$ on interval then f is convex. Local minima if $f'(x_0) = 0$ and $f''(x_0) > 0$.

5 Riemann Integration and Measure-theoretic remarks

5.1 Riemann sum definition

Partition $P = \{x_0 < \cdots < x_n\}$, sample points $\xi_i \in [x_{i-1}, x_i]$:

$$S(P, f) = \sum_{i=1}^n f(\xi_i) \Delta x_i, \quad \int_a^b f(x) dx = \lim_{\|P\| \rightarrow 0} S(P, f).$$

5.2 Riemann integrability criterion

Bounded f on $[a, b]$ is Riemann integrable iff set of discontinuities has measure zero.

5.3 Fundamental Theorems of Calculus

- FTC1: If $F(x) = \int_a^x f(t) dt$ and f is integrable then F is continuous and if f is continuous at x then $F'(x) = f(x)$.
- FTC2: If F is any antiderivative of f on $[a, b]$ then $\int_a^b f(x) dx = F(b) - F(a)$.

5.4 Definite and Indefinite Integrals

$$\begin{aligned} \int f(x) dx &= F(x) + C, \quad F'(x) = f(x), \\ \int_a^b f(x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x, \\ \text{Fundamental Theorem: } \frac{d}{dx} \int_a^x f(t) dt &= f(x). \end{aligned}$$

5.5 Improper integrals

$$\int_a^\infty f(x) dx = \lim_{R \rightarrow \infty} \int_a^R f(x) dx,$$

convergent if limit exists. Comparison and p-test: $\int_1^\infty \frac{1}{x^p} dx$ converges iff $p > 1$.

6 Integration Techniques

6.1 Basic Integration Rules

$$\begin{aligned} \int x^n dx &= \frac{x^{n+1}}{n+1} + C, \quad n \neq -1, \\ \int e^x dx &= e^x + C, \quad \int \frac{1}{x} dx = \ln |x| + C, \\ \int \sin x dx &= -\cos x + C, \quad \int \cos x dx = \sin x + C. \end{aligned}$$

6.2 Techniques of Integration

$$\text{Substitution: } \int f(g(x))g'(x) dx = \int f(u) du,$$

$$\text{Integration by Parts: } \int u dv = uv - \int v du,$$

$$\text{Partial Fractions: } \frac{P(x)}{Q(x)} = \sum_i \frac{A_i}{(x - r_i)^k} + \cdots.$$

6.3 Trigonometric integrals

Use power reduction, substitution $t = \tan(x/2)$ for rational functions of \sin, \cos .

6.4 Reduction formulas

Examples:

$$I_n = \int \sin^n x dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} I_{n-2}.$$

6.5 Special definite integrals

$$\int_0^\infty e^{-ax} x^{\nu-1} dx = \frac{\Gamma(\nu)}{a^\nu}, \quad \operatorname{Re}(a) > 0, \operatorname{Re}(\nu) > 0.$$

6.6 Applications of Integration: Area, Volume, and Density

Area under a Curve (Single Variable)

$$A = \int_a^b f(x) dx, \quad \text{where } f(x) \geq 0,$$
$$A = \int_a^b |f(x)| dx \quad \text{if } f(x) \text{ changes sign on } [a, b].$$

Area between Two Curves

$$A = \int_a^b [f(x) - g(x)] dx, \quad \text{where } f(x) \geq g(x) \text{ on } [a, b].$$

Area using Vertical and Horizontal Slices

$$A_y = \int_{y_1}^{y_2} [x_{\text{right}}(y) - x_{\text{left}}(y)] dy.$$

Volume by Slicing

$$V = \int_a^b A(x) dx, \quad A(x) = \text{cross-sectional area at } x.$$

Volume of Revolution: Disk Method

$$V = \pi \int_a^b [f(x)]^2 dx, \quad (\text{rotation about x-axis}),$$

$$V = \pi \int_c^d [g(y)]^2 dy, \quad (\text{rotation about y-axis}).$$

Volume of Revolution: Washer Method

$$V = \pi \int_a^b ([R(x)]^2 - [r(x)]^2) dx,$$

where $R(x)$ = outer radius, $r(x)$ = inner radius.

Volume by Cylindrical Shells

$$V = 2\pi \int_a^b x f(x) dx \quad (\text{about y-axis}),$$

$$V = 2\pi \int_c^d y g(y) dy \quad (\text{about x-axis}).$$

Density and Mass Distributions

$$\text{Mass: } M = \int_a^b \rho(x) dx,$$

$$\text{Center of Mass: } \bar{x} = \frac{1}{M} \int_a^b x \rho(x) dx,$$

$$\text{Moment of Inertia: } I = \int_a^b r^2 \rho(x) dx.$$

Double Integrals over Rectangular Regions

$$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx, \quad R = [a, b] \times [c, d].$$

Double Integrals over Non-Rectangular Regions

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx,$$

or

$$\iint_R f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy.$$

Physical Applications of Double Integrals

$$\text{Mass: } M = \iint_R \rho(x, y) dA,$$

$$\text{Center of Mass: } \bar{x} = \frac{1}{M} \iint_R x \rho(x, y) dA, \quad \bar{y} = \frac{1}{M} \iint_R y \rho(x, y) dA,$$

$$\text{Moment of Inertia: } I_x = \iint_R y^2 \rho(x, y) dA, \quad I_y = \iint_R x^2 \rho(x, y) dA.$$

6.7 Integrals in Polar Coordinates

Polar Coordinate Relations

$$\begin{aligned} x &= r \cos \theta, & y &= r \sin \theta, \\ r^2 &= x^2 + y^2, & \tan \theta &= \frac{y}{x}, \\ dA &= r dr d\theta. \end{aligned}$$

Conversion of Double Integral to Polar Form

$$\iint_R f(x, y) dA = \iint_R f(r \cos \theta, r \sin \theta) r dr d\theta.$$

Area in Polar Coordinates

$$A = \int_{\theta_1}^{\theta_2} \frac{1}{2} [r(\theta)]^2 d\theta,$$

$$\text{If bounded by } r = f(\theta) : \quad A = \frac{1}{2} \int_{\theta_1}^{\theta_2} [f(\theta)]^2 d\theta.$$

Area Between Two Polar Curves

$$A = \frac{1}{2} \int_{\theta_1}^{\theta_2} ([r_2(\theta)]^2 - [r_1(\theta)]^2) d\theta.$$

Arc Length in Polar Coordinates

$$L = \int_{\theta_1}^{\theta_2} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta.$$

Surface Area of a Solid of Revolution (Polar Form)

$$S = 2\pi \int_{\theta_1}^{\theta_2} r \sin \theta \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta \quad (\text{rotation about x-axis}),$$

$$S = 2\pi \int_{\theta_1}^{\theta_2} r \cos \theta \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta \quad (\text{rotation about y-axis}).$$

Mass and Center of Mass in Polar Coordinates

$$\begin{aligned} M &= \iint_R \rho(r, \theta) r \, dr \, d\theta, \\ \bar{r} &= \frac{1}{M} \iint_R r^2 \rho(r, \theta) \, dr \, d\theta, \\ \bar{\theta} &= \frac{1}{M} \iint_R \theta r \rho(r, \theta) \, dr \, d\theta. \end{aligned}$$

Moments and Inertia in Polar Coordinates

$$\begin{aligned} I_x &= \iint_R (r \sin \theta)^2 \rho(r, \theta) r \, dr \, d\theta, \\ I_y &= \iint_R (r \cos \theta)^2 \rho(r, \theta) r \, dr \, d\theta, \\ I_z &= \iint_R r^2 \rho(r, \theta) r \, dr \, d\theta. \end{aligned}$$

7 Special Functions

7.1 Gamma function

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} \, dt, \quad \Gamma(z+1) = z\Gamma(z), \quad \Gamma(n) = (n-1)!.$$

Reflection and duplication:

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}, \quad \Gamma(z)\Gamma\left(z + \frac{1}{2}\right) = 2^{1-2z} \sqrt{\pi} \, \Gamma(2z).$$

7.2 Beta function

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} \, dt = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

7.3 Stirling's approximation

$$\Gamma(z+1) \sim \sqrt{2\pi z} \left(\frac{z}{e}\right)^z \quad (z \rightarrow \infty).$$

7.4 Error function and Gaussian integrals

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} \, dt, \quad \int_{-\infty}^\infty e^{-x^2} \, dx = \sqrt{\pi}.$$

7.5 Bessel, Legendre, orthogonal polynomials (formulas)

Bessel ODE:

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0, \quad J_\nu(x) = \sum_{m=0}^\infty \frac{(-1)^m}{m! \Gamma(m + \nu + 1)} \left(\frac{x}{2}\right)^{2m+\nu}.$$

8 Multivariable Calculus

8.1 Functions of several variables

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, \quad \text{dom}(f) \subset \mathbb{R}^n.$$

8.2 Limits and continuity in \mathbb{R}^n

Definition uses $\|x - a\|$ and epsilon-delta. Path dependence matters for nonexistence.

8.3 Vector-Valued Functions

Definition and Representation

$$\begin{aligned}\vec{r}(t) &= \langle x(t), y(t), z(t) \rangle, \quad t \in I \subset \mathbb{R}, \\ \text{Domain: } &\{t \in \mathbb{R} : x(t), y(t), z(t) \text{ are defined}\}, \\ \text{Range: } &\{\vec{r}(t) : t \in I\} \subset \mathbb{R}^3.\end{aligned}$$

Limits and Continuity

$$\begin{aligned}\lim_{t \rightarrow a} \vec{r}(t) &= \langle \lim_{t \rightarrow a} x(t), \lim_{t \rightarrow a} y(t), \lim_{t \rightarrow a} z(t) \rangle, \\ \vec{r}(t) \text{ continuous at } t = a &\iff x, y, z \text{ are continuous at } t = a.\end{aligned}$$

Differentiation

$$\begin{aligned}\vec{r}'(t) &= \frac{d\vec{r}}{dt} = \langle x'(t), y'(t), z'(t) \rangle, \\ \text{Tangent vector: } \vec{T}(t) &= \frac{\vec{r}'(t)}{\|\vec{r}'(t)\|}, \\ \text{Derivative rules: } (c\vec{r})' &= c\vec{r}', \quad (\vec{r}_1 + \vec{r}_2)' = \vec{r}_1' + \vec{r}_2', \\ \frac{d}{dt}(\vec{r}_1 \cdot \vec{r}_2) &= \vec{r}_1' \cdot \vec{r}_2 + \vec{r}_1 \cdot \vec{r}_2', \\ \frac{d}{dt}(\vec{r}_1 \times \vec{r}_2) &= \vec{r}_1' \times \vec{r}_2 + \vec{r}_1 \times \vec{r}_2'.\end{aligned}$$

Velocity, Speed, and Acceleration

$$\begin{aligned}\vec{v}(t) &= \vec{r}'(t), \quad \text{velocity vector,} \\ v &= \|\vec{v}(t)\|, \quad \text{speed,} \\ \vec{a}(t) &= \vec{v}'(t) = \vec{r}''(t), \quad \text{acceleration vector.}\end{aligned}$$

Arc Length and Reparameterization

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\vec{r}'(u)\| du, \\ \frac{ds}{dt} &= \|\vec{r}'(t)\|, \quad \frac{d\vec{r}}{ds} = \vec{T}(s).\end{aligned}$$

Curvature and Torsion

$$\kappa = \frac{\|\vec{r}'(t) \times \vec{r}''(t)\|}{\|\vec{r}'(t)\|^3},$$

Principal normal: $\vec{N} = \frac{\vec{T}'}{\|\vec{T}'\|},$

Binormal: $\vec{B} = \vec{T} \times \vec{N},$

$$\tau = -\frac{d\vec{B}/ds \cdot \vec{N}}{1} = \frac{(\vec{r}'(t), \vec{r}''(t), \vec{r}'''(t))}{\|\vec{r}'(t) \times \vec{r}''(t)\|^2}.$$

Tangential and Normal Components of Acceleration

$$a_T = \frac{d}{dt}\|\vec{v}\| = \frac{\vec{v} \cdot \vec{a}}{\|\vec{v}\|},$$

$$a_N = \sqrt{\|\vec{a}\|^2 - a_T^2} = \frac{\|\vec{v} \times \vec{a}\|}{\|\vec{v}\|}.$$

Projectile and Circular Motion

Projectile: $\vec{r}(t) = (v_0 \cos \alpha)t \hat{i} + (v_0 \sin \alpha)t - \frac{1}{2}gt^2 \hat{j},$

Circular: $\vec{r}(t) = \langle r \cos(\omega t), r \sin(\omega t) \rangle,$

$$\vec{v}(t) = r\omega \langle -\sin(\omega t), \cos(\omega t) \rangle, \quad \|\vec{v}\| = r\omega,$$

$$\vec{a}(t) = -r\omega^2 \langle \cos(\omega t), \sin(\omega t) \rangle.$$

8.4 Partial derivatives and differentiability

$$\frac{\partial f}{\partial x_i} = \lim_{h \rightarrow 0} \frac{f(x_1, \dots, x_i + h, \dots) - f(x)}{h}.$$

Differentiable at a iff

$$f(a+h) = f(a) + Df(a)h + o(\|h\|)$$

where $Df(a)$ is the gradient (row) or Jacobian.

8.5 Gradient, directional derivative, Hessian

$$\nabla f = \begin{pmatrix} \partial_{x_1} f \\ \vdots \\ \partial_{x_n} f \end{pmatrix}, \quad D_{\mathbf{u}}f(a) = \nabla f(a) \cdot \mathbf{u}.$$

Hessian $H_{ij} = \partial_{x_i x_j}^2 f$. Second derivative test uses eigenvalues of Hessian.

8.6 Chain rule (multivariable)

If $z = f(x, y)$ with $x = x(t), y = y(t)$ then

$$\frac{dz}{dt} = f_x \frac{dx}{dt} + f_y \frac{dy}{dt}.$$

General matrix form available via Jacobians.

8.7 Taylor series in several variables

Second order expansion:

$$f(\mathbf{x} + \mathbf{h}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top \mathbf{h} + \frac{1}{2} \mathbf{h}^\top H f(\mathbf{x}) \mathbf{h} + o(\|\mathbf{h}\|^2).$$

8.8 Multiple integrals

$$\iint_D f(x, y) dA = \int_{y_0}^{y_1} \int_{x_0(y)}^{x_1(y)} f(x, y) dx dy.$$

Fubini's theorem for integrable functions allows iterated integrals.

8.9 Change of variables and Jacobian

If $\mathbf{x} = \mathbf{x}(\mathbf{u})$ is a C^1 bijection,

$$\iint_{D_x} f(\mathbf{x}) d\mathbf{x} = \iint_{D_u} f(\mathbf{x}(\mathbf{u})) |\det J_{\mathbf{x}}(\mathbf{u})| d\mathbf{u}.$$

8.10 Common coordinate systems

Polar: (r, θ) , $dA = r dr d\theta$.

Cylindrical: (r, θ, z) , $dV = r dr d\theta dz$.

Spherical: (ρ, ϕ, θ) , $dV = \rho^2 \sin \phi d\rho d\phi d\theta$.

8.11 Line integrals

For curve $\mathbf{r}(t)$, $a \leq t \leq b$:

$$\int_C f ds = \int_a^b f(\mathbf{r}(t)) \|\mathbf{r}'(t)\| dt.$$

For vector field \mathbf{F} :

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt.$$

8.12 Conservative fields and potentials

$\mathbf{F} = \nabla \phi$ implies $\int_C \mathbf{F} \cdot d\mathbf{r} = \phi(b) - \phi(a)$. Condition in simply connected domain: $\nabla \times \mathbf{F} = 0$.

8.13 Surface integrals

Parameterize S by $\mathbf{r}(u, v)$:

$$\iint_S f dS = \iint_D f(\mathbf{r}(u, v)) \|\mathbf{r}_u \times \mathbf{r}_v\| du dv.$$

Flux:

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \mathbf{F}(\mathbf{r}) \cdot (\mathbf{r}_u \times \mathbf{r}_v) du dv.$$

8.14 Gradient, Divergence, Curl

$$\begin{aligned}\nabla f &= \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right), \\ \nabla \cdot \vec{F} &= \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}, \\ \nabla \times \vec{F} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix}.\end{aligned}$$

8.15 Integral Theorems

$$\begin{aligned}\text{Green's Theorem: } \oint_C (Pdx + Qdy) &= \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA, \\ \text{Stokes' Theorem: } \oint_C \vec{F} \cdot d\vec{r} &= \iint_S (\nabla \times \vec{F}) \cdot d\vec{S}, \\ \text{Divergence Theorem: } \iiint_V (\nabla \cdot \vec{F}) dV &= \iint_S \vec{F} \cdot d\vec{S}.\end{aligned}$$

8.16 Optimization with constraints

Lagrange multipliers: optimize $f(\mathbf{x})$ subject to $g(\mathbf{x}) = 0$ via

$$\nabla f = \lambda \nabla g.$$

Multiple constraints use vector of multipliers.

9 Ordinary Differential Equations

9.1 Classification and existence

General form: $F(x, y, y', \dots) = 0$. First order explicit: $y' = f(x, y)$. Picard-Lindelöf theorem: if f is Lipschitz in y then local uniqueness and existence.

9.2 Linear first-order ODEs

$$y' + p(x)y = q(x).$$

Integrating factor $\mu(x) = e^{\int p(x) dx}$ gives

$$y = \frac{1}{\mu} \left(\int \mu q dx + C \right).$$

9.3 Separable equations

$$\frac{dy}{dx} = g(x)h(y) \Rightarrow \int \frac{dy}{h(y)} = \int g(x) dx.$$

9.4 Second-order linear ODEs with constant coefficients

$$y'' + ay' + by = f(x).$$

Homogeneous solution from characteristic equation $r^2 + ar + b = 0$. Particular solution via undetermined coefficients or variation of parameters.

9.5 Method of undetermined coefficients

Assume particular solution form matching RHS type (polynomial, exponential, sine/cosine) multiplied by x^k if resonance occurs.

9.6 Variation of parameters (2nd order)

If y_1, y_2 fundamental solutions of homogeneous then particular solution:

$$y_p = -y_1 \int \frac{y_2 f}{W} dx + y_2 \int \frac{y_1 f}{W} dx, \quad W = y_1 y_2' - y_1' y_2.$$

9.7 Linear systems and matrix exponentials

System $\mathbf{x}' = A\mathbf{x}$. Solution:

$$\mathbf{x}(t) = e^{At} \mathbf{x}(0), \quad e^{At} = \sum_{k=0}^{\infty} \frac{(At)^k}{k!}.$$

Compute via diagonalization $A = S\Lambda S^{-1}$ or Jordan form.

9.8 Stability of equilibria

For autonomous $\mathbf{x}' = f(\mathbf{x})$, linearize at equilibrium \mathbf{x}_0 using Jacobian $J = f'(\mathbf{x}_0)$. Eigenvalues of J determine local stability.

9.9 Phase plane methods

Nullclines, critical points classification (node, saddle, spiral, center) by eigenvalues.

9.10 Boundary value problems and Sturm-Liouville

Form $-(p(x)y')' + q(x)y = \lambda w(x)y$ with BCs. Eigenfunctions form orthogonal basis for weighted L^2 .

10 Applications

10.1 Optimization

Critical points: $f'(x) = 0$, $f''(x) > 0 \Rightarrow$ local min, $f''(x) < 0 \Rightarrow$ local max.

11 Partial Differential Equations: Classical Examples and Methods

11.1 Canonical PDEs

- Heat equation: $u_t = \kappa \Delta u$.
- Wave equation: $u_{tt} = c^2 \Delta u$.
- Laplace's equation: $\Delta u = 0$.

11.2 Separation of variables

Assume $u(x, t) = X(x)T(t)$ leading to ODEs and eigenvalue problems.

11.3 Fourier series and transforms

Fourier series on $[-L, L]$:

$$f(x) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right],$$

coefficients given by orthogonality. Fourier transform:

$$\hat{f}(\omega) = \int_{-\infty}^{\infty} f(x) e^{-i\omega x} dx, \quad f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\omega) e^{i\omega x} d\omega.$$

11.4 Green's functions

For linear operator \mathcal{L} , solution to $\mathcal{L}u = f$ can be represented via $u = \mathcal{G}f$ where \mathcal{G} involves Green's function kernel solving $\mathcal{L}G(x, s) = \delta(x - s)$ with BCs.

12 Transforms and Operational Methods

12.1 Laplace transform

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt.$$

Properties: linearity, shift, differentiation in time $\mathcal{L}\{f'\} = sF(s) - f(0)$, convolution theorem $\mathcal{L}\{f * g\} = F(s)G(s)$.

12.2 Z-transform (discrete)

$$\mathcal{Z}\{x[n]\} = X(z) = \sum_{n=0}^{\infty} x[n] z^{-n}.$$

12.3 Fourier transform properties

Linearity, modulation, differentiation, convolution.

13 Numerical Methods

13.1 Root finding

- Newton's method: $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$.
- Secant method and bisection (guaranteed convergence for continuous sign change).

13.2 Numerical integration

$$\text{Trapezoidal rule: } \int_a^b f(x) dx \approx \frac{b-a}{2}(f(a) + f(b))$$

Simpson's rule and composite versions. Error estimates involve higher derivatives.

13.3 Numerical ODEs

- Euler forward: $y_{n+1} = y_n + hf(t_n, y_n)$.
- Improved Euler/Heun and classical RK4:

$$k_1 = f(t_n, y_n), \quad k_2 = f(t_n + \frac{h}{2}, y_n + \frac{h}{2}k_1), \quad k_3 = \dots, \quad k_4 = f(t_n + h, y_n + hk_3),$$

$$y_{n+1} = y_n + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4).$$

14 Miscellaneous Topics

14.1 Uniform convergence

Sequence of functions f_n converges uniformly to f if $\sup_x |f_n(x) - f(x)| \rightarrow 0$. Uniform convergence permits termwise integration and differentiation under conditions.

14.2 Dominated convergence theorem (measure-theory remark)

If $f_n \rightarrow f$ a.e. and $|f_n| \leq g$ with $g \in L^1$ then $\lim \int f_n = \int f$.

14.3 Asymptotic expansions and stationary phase

Common tools for approximating integrals and series for large parameters.

14.4 Calculus of variations (principles)

Euler-Lagrange equation: stationary action for functional $J[y] = \int_a^b L(x, y, y') dx$ yields

$$\frac{\partial L}{\partial y} - \frac{d}{dx} \left(\frac{\partial L}{\partial y'} \right) = 0.$$

15 Cheat Sheet: Common Identities and Integrals

15.1 Derivatives

$$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} \arctan x = \frac{1}{1+x^2}.$$

15.2 Integrals

$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \arctan \frac{x}{a} + C, \quad \int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + C.$$

15.3 Useful limits and series

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}, \quad \sum_{n=0}^{\infty} \frac{x^n}{n!} = e^x.$$

16 Reference Theorems (Statements)

Theorem 16.1 (Taylor's theorem). *If $f^{(n+1)}$ exists on interval containing a and x then*

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k + \frac{f^{(n+1)}(\xi)}{(n+1)!} (x-a)^{n+1}.$$

Theorem 16.2 (Uniform convergence interchange). *If $f_n \rightarrow f$ uniformly on $[a, b]$ and each f_n integrable then $\int_a^b f_n \rightarrow \int_a^b f$.*

Theorem 16.3 (Green-Stokes-Gauss collection). *The respective integral theorems relating derivatives inside a domain to boundary integrals hold under stated regularity.*
