

Exponential Lower-bounds via Exponential Sums

Somnath Bhattacharjee
(Chennai Mathematical Institute)

Joint work with Markus Bläser (Saarland University), Pranjal Dutta
(NUS), Saswata Mukherjee (NUS)

(ICALP 2024)

Outline

Motivation

High level idea

Towards Explicitness

Conclusion

NP Problems: Is Brute Force Optimal?

NP Problems: Is Brute Force Optimal?

Given a word x , check for

$$\bigvee_{e \in \{0,1\}^n} M(x, e) = 1$$

NP Problems: Is Brute Force Optimal?

Given a word x , check for

$$\bigvee_{e \in \{0,1\}^n} M(x, e) = 1$$

M runs in m time

NP Problems: Is Brute Force Optimal?

Given a word x , check for

$$\bigvee_{e \in \{0,1\}^n} M(x, e) = 1$$

M runs in m time

- Running time upper bound : $2^n m$ (Brute force!)

NP Problems: Is Brute Force Optimal?

Given a word x , check for

$$\bigvee_{e \in \{0,1\}^n} M(x, e) = 1$$

M runs in m time

- Running time upper bound : $2^n m$ (Brute force!)
- Improve to $2^{o(n)} \text{poly}(m)$ possible?

NP Problems: Is Brute Force Optimal?

Given a word x , check for

$$\bigvee_{e \in \{0,1\}^n} M(x, e) = 1$$

M runs in m time

- Running time upper bound : $2^n m$ (Brute force!)
- Improve to $2^{o(n)} \text{poly}(m)$ possible?
- **ETH** says **NO!** (informally)

Is Brute Force Optimal?


- Can we ask the same question in *algebraic setting*?

Is Brute Force Optimal?

- Can we ask the same question in *algebraic setting*?
- What is even the *Computational Model* in that setting?

Is Brute Force Optimal?

- Can we ask the same question in *algebraic setting*?
- What is even the *Computational Model* in that setting?
- Hence *Algebraic Circuit* enters the picture



Algebraic Circuits (Constant-free)

- Arithmetic Circuits are **directed acyclic graphs**.

Algebraic Circuits (Constant-free)

- Arithmetic Circuits are **directed acyclic graphs**.
- Each internal node: $+$ or \times gate.

Algebraic Circuits (Constant-free)

- Arithmetic Circuits are **directed acyclic graphs**.
- Each internal node: $+$ or \times gate.
- Each leaf: $\{1, 0, -1\}$ or variables **X**

Algebraic Circuits (Constant-free)

- Arithmetic Circuits are **directed acyclic graphs**.
- Each internal node: $+$ or \times gate.
- Each leaf: $\{1, 0, -1\}$ or variables X
- Computes a polynomial in $\mathbb{Z}[X]$

Algebraic Circuits (Constant-free)

- Arithmetic Circuits are **directed acyclic graphs**.
- Each internal node: $+$ or \times gate.
- Each leaf: $\{1, 0, -1\}$ or variables X
- Computes a polynomial in $\mathbb{Z}[X]$
- Complexity Measure: **# Edges** in the circuit

Tau-Complexity

Given a polynomial $P(\mathbf{x}) \in \mathbb{Z}[\mathbf{x}]$

Tau-Complexity

Given a polynomial $P(\mathbf{X}) \in \mathbb{Z}[\mathbf{X}]$

$\tau(P(\mathbf{X}))$: Size of **smallest circuit** that computes $P(\mathbf{X})$

Tau-Complexity

Given a polynomial $P(\mathbf{X}) \in \mathbb{Z}[\mathbf{X}]$

$\tau(P(\mathbf{X}))$: Size of **smallest circuit** that computes $P(\mathbf{X})$

Examples

1. $\tau(2^{2^k}) = \Theta(k)$, $\tau(x^n) = \Theta(\log n)$

Tau-Complexity

Given a polynomial $P(\mathbf{X}) \in \mathbb{Z}[\mathbf{X}]$

$\tau(P(\mathbf{X}))$: Size of **smallest circuit** that computes $P(\mathbf{X})$

Examples

1. $\tau(2^{2^k}) = \Theta(k)$, $\tau(x^n) = \Theta(\log n)$

2. $\tau(n!) = ?$, $\tau(\prod_{i=1}^n (x + i)) = ?$

Tau-Complexity

Given a polynomial $P(\mathbf{X}) \in \mathbb{Z}[\mathbf{X}]$

$\tau(P(\mathbf{X}))$: Size of **smallest circuit** that computes $P(\mathbf{X})$

Examples

1. $\tau(2^{2^k}) = \Theta(k)$, $\tau(x^n) = \Theta(\log n)$

2. $\tau(n!) = ?$, $\tau(\prod_{i=1}^n (x + i)) = ?$

We believe these are $\geq \Omega(n)$

Tau-Complexity

Blum-Shub-Smale Tau-Conjecture:

Tau-Complexity

Blum-Shub-Smale Tau-Conjecture: For $P(x) \in \mathbb{Z}[x]$

Tau-Complexity


Blum-Shub-Smale Tau-Conjecture: For $P(x) \in \mathbb{Z}[x]$

$$\tau(P(x)) \geq [\# \text{ integer roots of } P(x)]^c$$

Tau-Complexity

Blum-Shub-Smale Tau-Conjecture: For $P(x) \in \mathbb{Z}[x]$

$$\tau(P(x)) \geq [\# \text{ integer roots of } P(x)]^c$$



Exponential sum (VNP_0)

$$P_{n,m}(\mathbf{x}) = \sum_{\mathbf{y} \in \{0,1\}^n} g(\mathbf{x}, \mathbf{y})$$

Exponential sum (VNP_0)

$$P_{n,m}(\mathbf{x}) = \sum_{\mathbf{y} \in \{0,1\}^n} g(\mathbf{x}, \mathbf{y})$$

Circuit size of g is m , $g[\mathbf{X}, \mathbf{Y}] \in \mathbb{Z}[\mathbf{X}, \mathbf{Y}]$.

Exponential sum (VNP_0)

$$P_{n,m}(\mathbf{X}) = \sum_{\mathbf{y} \in \{0,1\}^n} g(\mathbf{X}, \mathbf{y})$$

Circuit size of g is m , $g[\mathbf{X}, \mathbf{Y}] \in \mathbb{Z}[\mathbf{X}, \mathbf{Y}]$.

- Note $\tau(P_{n,m}) \leq 2^n m$

Exponential sum (VNP_0)

$$P_{n,m}(\mathbf{X}) = \sum_{\mathbf{y} \in \{0,1\}^n} g(\mathbf{X}, \mathbf{y})$$

Circuit size of g is m , $g[\mathbf{X}, \mathbf{Y}] \in \mathbb{Z}[\mathbf{X}, \mathbf{Y}]$.

- Note $\tau(P_{n,m}) \leq 2^n m$
- $\tau(P_{m,n}) = 2^{o(n)} \text{poly}(m)$ possible?

Exponential sum (VNP_0)

$$P_{n,m}(\mathbf{X}) = \sum_{\mathbf{y} \in \{0,1\}^n} g(\mathbf{X}, \mathbf{y})$$

Circuit size of g is m , $g[\mathbf{X}, \mathbf{Y}] \in \mathbb{Z}[\mathbf{X}, \mathbf{Y}]$.

- Note $\tau(P_{n,m}) \leq 2^n m$
- $\tau(P_{m,n}) = 2^{o(n)} \text{poly}(m)$ possible?
- Does τ -conjecture imply some lower bound?

Exponential sum (VNP_0)

$$P_{n,m}(\mathbf{X}) = \sum_{\mathbf{y} \in \{0,1\}^n} g(\mathbf{X}, \mathbf{y})$$

Circuit size of g is m , $g[\mathbf{X}, \mathbf{Y}] \in \mathbb{Z}[\mathbf{X}, \mathbf{Y}]$.

- Note $\tau(P_{n,m}) \leq 2^n m$
- $\tau(P_{m,n}) = 2^{o(n)} \text{poly}(m)$ possible?
- Does τ -conjecture imply some lower bound?
- [Bürgisser'07] showed super-polynomial lowerbound on $P_{m,n}$ assuming τ -conjecture

Main result

Conditional Optimal Lower Bound

Assuming τ -conjecture \exists a polynomial family $P_{n,m}(\mathbf{X}) \in \mathbb{Z}[\mathbf{X}]$ of exponential sum which requires $2^{\Omega(n)} \text{poly}(m)$ size circuit.

Bürgisser's Proof analysis

Assume every $P_{n,m}(X)$ has $\text{poly}(m)$ circuit

Bürgisser's Proof analysis

Assume every $P_{n,m}(\mathbf{X})$ has $\text{poly}(m)$ circuit



Lots of Bad things happen

Bürgisser's Proof analysis

Assume every $P_{n,m}(\mathbf{X})$ has $\text{poly}(m)$ circuit



Lots of Bad things happen



$\prod_{i=1}^n (x + i)$ has easy coefficients



It has $\text{poly}(\log n)$ size circuit

Bürgisser's Proof analysis: Observations



Bürgisser's Proof analysis: Observations



Bürgisser's Proof analysis: Observations

Started with Poly circuit for
Exp Sum



We get **Poly (log)** circuit



But we have n many roots



Bürgisser's Proof analysis: Observations

Started with Poly circuit for
Exp Sum



We get **Poly (log)** circuit



But we have n many roots




May be starting with SubExp
gives Sub linear circuit




Bürgisser's Proof analysis: Observations



Bürgisser's Proof analysis: Observations



But we can improve it!



What's the Magic

$$\prod_{i=1}^n (x + i) = \sum_{k=0}^n S_{n,k}(1, \dots, n) x^k$$

What's the Magic

$$\prod_{i=1}^n (x + i) = \sum_{k=0}^n S_{n,k}(1, \dots, n) x^k$$

where

$$S_{n,k}(X_1, \dots, X_n) = \sum_{S \subseteq [n], |S|=k} \prod_{i \in S} X_i$$

What's the Magic

$$1, 2, \dots, n, \quad x^k$$

What's the Magic

$$1 \times \dots \times k, \quad 2 \times \dots \times (k+1), \quad \dots$$

$$1, 2, \dots, n, \quad x^k$$

What's the Magic

$$S_{n,k} := 1 \times \dots \times k + 2 \times \dots \times (k+1) + \dots$$

$$1 \times \dots \times k, \quad 2 \times \dots \times (k+1), \quad \dots$$

$$1, 2, \dots, n, \quad x^k$$

What's the Magic

$$S_{n,n}x^n + S_{n,n-1}x^{n-1} + \cdots + S_{n,0}$$

$$S_{n,k} := 1 \times \cdots \times k + 2 \times \cdots \times (k+1) + \cdots$$

$$1 \times \cdots \times k, \quad 2 \times \cdots \times (k+1), \quad \dots$$

$$1, 2, \dots, n, \quad x^k$$

What's the Magic

$$S_{n,n}x^n + S_{n,n-1}x^{n-1} + \cdots + S_{n,0}$$

3 level

$$S_{n,k} := 1 \times \cdots \times k + 2 \times \cdots \times (k+1) + \cdots$$

2 level

$$1 \times \cdots \times k, \quad 2 \times \cdots \times (k+1), \quad \dots$$

1 level

$$1, 2, \dots, n, \quad x^k$$

0 level

Linear Counting Hierarchy

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,
we define $\mathbf{C}_{lin} \cdot K$ by

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,
we define $\mathbf{C}_{lin}.K$ by

$A \in \mathbf{C}_{lin}.K$ if there is some $B \in K$

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,
we define $\mathbf{C}_{lin}.K$ by

$A \in \mathbf{C}_{lin}.K$ if there is some $B \in K$
and a linear function $\ell : \mathbb{N} \rightarrow \mathbb{N}$, $\ell(n) = O(n)$

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,
we define $\mathbf{C}_{lin}.K$ by

$A \in \mathbf{C}_{lin}.K$ if there is some $B \in K$

and a linear function $\ell : \mathbb{N} \rightarrow \mathbb{N}$, $\ell(n) = O(n)$

and some polynomial time computable function $f : \{0, 1\}^* \rightarrow \mathbb{N}$

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,

we define $\mathbf{C}_{lin}.K$ by

$A \in \mathbf{C}_{lin}.K$ if there is some $B \in K$

and a linear function $\ell : \mathbb{N} \rightarrow \mathbb{N}$, $\ell(n) = O(n)$

and some polynomial time computable function $f : \{0, 1\}^* \rightarrow \mathbb{N}$
such that,

$$x \in A \iff |\{y \in \{0, 1\}^{\ell(|x|)} : \langle x, y \rangle \in B\}| > f(x).$$

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,
We define $\text{C-lin}_0 K := K$ and for all $k \in \mathbb{N}$,
 $\text{C-lin}_{k+1} K := \mathbf{C}_{\text{lin}}.\text{C-lin}_k K$.

Linear Counting Hierarchy

Linear Counting Hierarchy

Given a complexity class K ,

We define $\text{C-lin}_0 K := K$ and for all $k \in \mathbb{N}$,

$\text{C-lin}_{k+1} K := \mathbf{C}_{\text{lin}}.\text{C-lin}_k K$.

The *linear counting hierarchy* is $\text{CH}_{\text{lin}} K := \bigcup_{k \geq 0} \text{C-lin}_k K$

Linear Counting Hierarchy (CH_{lin})

Characterization of CH_{lin}

$(k + 1)^{\text{th}}$ level of CH_{lin} is Exponential sum of k^{th} level

Linear Counting Hierarchy (CH_{lin})

Characterization of CH_{lin}

$(k+1)^{\text{th}}$ level of CH_{lin} is Exponential sum of k^{th} level

Hence Exponential sum is EASY $\implies \text{CH}_{\text{lin}}$ collapses

$$\implies \prod_{i=1}^n (x + i) \text{ is EASY}$$

Permanent

Given a variable matrix

$$\mathbf{X} := \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & & \ddots & \vdots \\ X_{n1} & X_{n2} & \dots & X_{nn} \end{bmatrix}_{n \times n}$$

Permanent

Given a variable matrix

$$\mathbf{X} := \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & & \ddots & \vdots \\ X_{n1} & X_{n2} & \dots & X_{nn} \end{bmatrix}_{n \times n}$$

$$\text{Per}_n(\mathbf{X}) := \sum_{\sigma \in S_n} X_{1,\sigma_1} X_{2,\sigma_2} \dots X_{n,\sigma_n}$$

Completeness of Permanent

Ryser formula: Per_n can be written as Exponential sum (n, n^2)

Completeness of Permanent

Ryser formula: Per_n can be written as Exponential sum (n, n^2)

[Valiant 79]: Any Exponential sum (n, m) can be written as Per of $m^4 \times m^4$ matrix with entries $1, 0, -1, X$

Completeness of Permanent

Ryser formula: Per_n can be written as Exponential sum (n, n^2)

[Valiant 79]: Any Exponential sum (n, m) can be written as Per of $m^4 \times m^4$ matrix with entries $1, 0, -1, X$

- Super Polynomial lower bound on Exponential sum \iff
Super Polynomial lower bound on Per

Completeness of Permanent

Ryser formula: Per_n can be written as Exponential sum (n, n^2)

[Valiant 79]: Any Exponential sum (n, m) can be written as Per of $m^4 \times m^4$ matrix with entries $1, 0, -1, X$

- Super Polynomial lower bound on Exponential sum \iff Super Polynomial lower bound on Per
- NOT true for Exponential lower bounds

Completeness of Permanent

Ryser formula: Per_n can be written as Exponential sum (n, n^2)

[Valiant 79]: Any Exponential sum (n, m) can be written as Per of $m^4 \times m^4$ matrix with entries $1, 0, -1, X$

- Super Polynomial lower bound on Exponential sum \iff Super Polynomial lower bound on Per
- NOT true for Exponential lower bounds
- We gave $2^{n^{1/8}}$ lower bound for Per_n

Completeness of Permanent

Ryser formula: Per_n can be written as Exponential sum (n, n^2)

[Valiant 79]: Any Exponential sum (n, m) can be written as Per of $m^4 \times m^4$ matrix with entries $1, 0, -1, X$

- Super Polynomial lower bound on Exponential sum \iff Super Polynomial lower bound on Per
- NOT true for Exponential lower bounds
- We gave $2^{n^{1/8}}$ lower bound for Per_n (Conditionally)

Other Results

1. We achieved optimal lower bound from tau conjecture for **Parameterized Algebraic classes** defined in [Bläser and Engles 18] (which are analogous to $\#W[t]$ classes)

Other Results

1. We achieved optimal lower bound from tau conjecture for **Parameterized Algebraic classes** defined in [Bläser and Engles 18] (which are analogous to $\#W[t]$ classes)
2. We achieved **completeness** result for parameterized valiant classes.

Open Problems

1. Can we established conditional truly exponential (ie, $2^{\Omega(n)}$ $\text{poly}(n)$) lower bound for Per_n ? (Unconditional will be better :))
2. Can we get Lower Bounds for NP from tau-conjecture? (We don't know even super-polynomial bound)