
192	146M	3M	91M
Count of countries	Total confirmed cases	Total death cases.	Total recovered cases




Here each bubble represent one of the 192 countries.
Bubble size is determined by cases confirmed.
Hover on any bubble for further details.


Steps taken by Indian Government,Hover to know the steps

Steps

Top 10 countries with highest cases confirmed



Bottom10 countries with lowest cases confirmed



Year

- [ ] 2020
- [ ] 2021

61.92%
Recovery rate

Afghanistan
country name


2.12%
fatality rate

countries	Month	confirmed_cases	DEATH_CASES	Recovered_cases	active_cases
Afghanistan	January	1668119	71838	1403684	192597
Afghanistan	February	1553108	67864	1358287	126957
Afghanistan	March	1738232	76254	1542006	119972
Afghanistan	April	1404560	61564	1252270	90726
Afghanistan	May	225674	5007	24129	196538
Afghanistan	June	750944	15016	184768	551160
Afghanistan	July	1077247	33518	688117	355612
Afghanistan	August	1163297	42101	843438	277758
Afghanistan	September	1163868	42987	949104	171777
Afghanistan	October	1245948	46287	1039280	160381
Afghanistan	November	1308111	49133	1058724	200254
Afghanistan	December	1522209	62357	1207801	252051
Albania	January	2088163	39480	1250570	798113
Total		**20661747798**	**507583169**	**12074960658**	**8079203971**


New cases by month

Month	countries	new cases
January	Afghanistan	3497
February	Afghanistan	692
March	Afghanistan	914
April	Afghanistan	4228
May	Afghanistan	13081
June	Afghanistan	16299
July	Afghanistan	5158
August	Afghanistan	1494
September	Afghanistan	1109
October	Afghanistan	2157
Total		**146461460**

Cases By Year and Month



Cases Hierarchy Funnel



First wave from 1st April 2020 to 31st January 2021

first

Second wave from 1st March 2021 to 30th June 2021

second

On 11 January 2020, the WHO confirmed that a novel coronavirus was the cause of a respiratory illness in a cluster of people in India City, Jaipur, India, which was reported to the WHO on 31 December 2019

Protective measures were first applied in January. India began thermal screening of passengers arriving from China

On 17 March, the Government of India issued an advisory, urging all Indian states to take social distancing measures as a preventive strategy

Despite the early lockdown, India's COVID-19 cases surged in September 2020, with over 90,000 cases reported daily
India's Health Ministry published its Guidelines on COVID-19 management, which included approval of hydroxychloroquine.

The Indian government stopped issuing of new visas. Previously issued visas for the nationals of Iran, South Korea, Japan, and Italy were suspended

Over the month of March, multiple states across the country began shutting down schools, colleges, public facilities such as malls, gyms, cinema halls and other public places to contain the spread.

The Government Divided the entire nation into three zones – Green Zone, Red Zone, Orange Zone. Relaxations would be allowed accordingly

Timing of relaxation in restriction policies by the central and state governments, resumption of public transport, opening up of market places and businesses, large gatherings at social, religious and political events, noncompliance of COVID appropriate behaviour by the general public and emergence of new mutant strains presumed to have high infectivity potential, are thought to be factors resulting in the second COVID wave in India.

The focus of the policy that was adopted when the first wave started declining was on test, trace and treat, continued emphasis on COVID-specific prevention behaviour, conservative stepwise approach to reverse lockdown restrictions and anti-COVID-19 vaccination drive

A theoretical simplistic model for control of COVID-19 pandemic can be built around the two-pronged strategy of preventing the spread of infection and reducing mortality among infected people. Designing the policies and strategies for the surge of COVID-19 cases must be based on the lessons learnt and evidence generated during the first wave.

According to my opinion government should have banned entry for foreigners in the Begining and should have let indian residents coming from abroad leave after 15 days of observation.This would help to reduce transmission