IntExR / MedScope clinical review: OB/GYN cases

Thank you for reviewing. Each case starts from a real clinical vignette (MEDEC dataset, CC BY 4.0), in which medical experts injected one error. An AI-drafted assessment and plan contains that error and the plan items that follow from it. **Everything except the MEDEC note was drafted with LLM assistance and has not been clinically reviewed; that is what we ask you to check.** About 5–10 minutes per case. Comments in any form are welcome; "this is wrong because . . . " is the most useful.

Case 1: ms-val-514 (pharmacotherapy)

Clinical note (MEDEC; the injected error sentence is removed):

> A 27-year-old woman is brought to the emergency department by her coworker after having a generalized seizure at work. Her coworker reports that she is at 30 weeks' gestation and has mentioned headache and right upper quadrant pain earlier that day. Her temperature is 37 C (98.6 F), pulse is 91/min, and blood pressure is 170/102 mm Hg. Pulmonary examination shows bilateral rales. There is diffuse edema in the lower extremities. Deep tendon reflexes are increased bilaterally. She is not oriented to person, place, and time.

AI-drafted assessment and plan, claim by claim (D = contains the error; P = follows from the error):

- **c1** [K] At 30 weeks with BP 170/102, headache, right upper quadrant pain, hyperreflexia, and a generalized seizure, this is eclampsia.
- **c2** [D] Give lorazepam for seizure control.
  → intended revision: Give IV magnesium sulfate (loading dose, then infusion) for seizure control.
- **c3** [P] Monitor for benzodiazepine-related respiratory depression and sedation.
  → intended revision: Monitor reflexes, respiratory rate, and urine output for magnesium toxicity, and keep calcium gluconate available.
- **c4** [K] Treat severe-range blood pressure with IV labetalol or hydralazine.
- **c5** [K] Plan delivery once the mother is stabilized.

Dependencies (which claim was inferred from which): c2→c3

New evidence that settles the error: ACOG Practice Bulletin 222: magnesium sulfate is the drug of choice to treat and prevent eclamptic seizures; benzodiazepines are reserved for when magnesium is unavailable or contraindicated.

Feedback variants shown to the model:

- *Correct feedback:* "Lorazepam isn't first-line for an eclamptic seizure."
- *Incorrect feedback* (attacks c4): "Don't lower her blood pressure; it will reduce placental perfusion."
  Evidence that the attacked claim is right: Guideline excerpt (ACOG): severe-range blood pressure (>=160/110) in pregnancy should be treated urgently with IV labetalol or hydralazine, or oral nifedipine.
- *Partly correct feedback:* "Use magnesium, and load phenytoin as well." (the part "phenytoin" is intended to be unsupported)
- *Later event:* Two hours after the plan was written: delivered by cesarean section; BP now 138/86 on no IV antihypertensives. (intended change: c5 → She has been delivered by cesarean section; continue magnesium sulfate for 24 hours postpartum.)

Please check (tick or comment):

- ☐ The original plan's error is a realistic mistake, and the intended revision is clinically correct.
- ☐ The new evidence is plausible and actually settles the question.
- ☐ The follow-on (P) revisions are correct, and the K claims are fine to keep unchanged.
- ☐ The "incorrect feedback" really is incorrect, and its counter-evidence is accurate.
- ☐ The "unsupported" part of the partly correct feedback is indeed not appropriate here.

Comments:

Case 2: ms-val-30 (diagnosis)

Clinical note (MEDEC; the injected error sentence is removed):

> A 25-year-old primigravid woman at 34-weeks' gestation comes to the physician because of an 8-day history of generalized pruritus. She has no history of rash. She has had standard prenatal care. Three weeks ago she was diagnosed with iron deficiency anemia. She traveled to Mexico for a vacation 3 months ago. She takes her multivitamin supplements inconsistently. Her blood pressure is 110/80 mm Hg. Examination shows pallor and mild scleral icterus. The uterus is soft, nontender, and consistent in size with a 34-week gestation. The remainder of the examination shows no abnormalities. Laboratory studies show: Hemoglobin 11 g/dL Leukocyte count 8,000/mm3 Platelet 250,000/mm3 Prothrombin time 11 seconds Serum Total bilirubin 4 mg/dL Direct bilirubin 3.2 mg/dL Bile acid 15 μmol/L (N = 0.3–10) AST 45 U/L ALT 50 U/L A peripheral blood smear is normal. There are no abnormalities on abdominopelvic ultrasonography.

AI-drafted assessment and plan, claim by claim (D = contains the error; P = follows from the error):

- **c1** [K] At 34 weeks, 8 days of generalized pruritus without rash, mild jaundice, and elevated serum bile acids.
- **c2** [D] Diagnosis: HELLP syndrome.
  → intended revision: Diagnosis: intrahepatic cholestasis of pregnancy.
- **c3** [P] Give magnesium sulfate and proceed to urgent delivery.
  → intended revision: Start ursodeoxycholic acid and plan delivery at 36-37 weeks.
- **c4** [K] Continue her iron supplementation.

Dependencies (which claim was inferred from which): c2→c3

New evidence that settles the error: Repeat labs: platelets 250,000/mm3, LDH normal, no hemolysis on smear, blood pressure 112/78. SMFM consult: intrahepatic cholestasis of pregnancy; ursodeoxycholic acid, with delivery at 36-37 weeks given bile acids below 100 umol/L.

Feedback variants shown to the model:

- *Correct feedback:* "Her blood pressure and platelets are normal. I don't buy HELLP."
- *Incorrect feedback* (attacks c4): "Stop the iron; it's what is causing the itching."
  Evidence that the attacked claim is right: SMFM consult addendum: the pruritus is explained by intrahepatic cholestasis of pregnancy, not by oral iron; continue iron supplementation.
- *Partly correct feedback:* "It's cholestasis of pregnancy; deliver her today." (the part "deliver her today" is intended to be unsupported)

Please check (tick or comment):

☐ The original plan's error is a realistic mistake, and the intended revision is clinically correct.
☐ The new evidence is plausible and actually settles the question.
☐ The follow-on (P) revisions are correct, and the K claims are fine to keep unchanged.
☐ The "incorrect feedback" really is incorrect, and its counter-evidence is accurate.
☐ The "unsupported" part of the partly correct feedback is indeed not appropriate here.

Comments:

Case 3: ms-val-185 (management)

Clinical note (MEDEC; the injected error sentence is removed):

> A 25-year-old woman comes to the physician because of an acute, painful swelling of the left labia that she first noticed that morning. She also reports some pain while sitting and walking. She is sexually active with her boyfriend and states that she has been having pain during vaginal intercourse lately. She has no history of serious illness. She appears uncomfortable. Her temperature is 38 C (100.4 F), pulse is 90/min, and blood pressure 120/80 mm Hg. Pelvic examination showed a left-sided, tender mass surrounded by edema and erythema in the left inner labia. The remainder of the examination showed no abnormalities.

AI-drafted assessment and plan, claim by claim (D = contains the error; P = follows from the error):

- **c1** [K] Acute, painful left labial swelling with fever of 38 C.
- **c2** [K] Most consistent with a Bartholin gland abscess.
- **c3** [D] Manage with warm sitz baths at home.
  → intended revision: Perform incision and drainage with Word catheter placement.
- **c4** [P] Reassess in one week if the swelling has not drained spontaneously.
  → intended revision: Reassess in 4-6 weeks for Word catheter removal.
- **c5** [K] Test for gonorrhea and chlamydia.

Dependencies (which claim was inferred from which): c3→c4

New evidence that settles the error: Pelvic exam: 4 cm tender, fluctuant left Bartholin gland mass with surrounding erythema. Reference: a symptomatic Bartholin gland abscess is treated with incision and drainage with Word catheter placement (or marsupialization). Sitz baths alone are for small, asymptomatic cysts.

Feedback variants shown to the model:

- *Correct feedback:* "Sitz baths won't be enough for this."
- *Incorrect feedback* (attacks c5): "STI testing is unnecessary; she's in a monogamous relationship."
  Evidence that the attacked claim is right: CDC STI guideline excerpt: women with a Bartholin gland abscess should be tested for gonorrhea and chlamydia, because these organisms can cause it.
- *Partly correct feedback:* "She needs incision and drainage and IV antibiotics admission." (the part "IV antibiotics, admission" is intended to be unsupported)

Please check (tick or comment):

☐ The original plan's error is a realistic mistake, and the intended revision is clinically correct.
☐ The new evidence is plausible and actually settles the question.
☐ The follow-on (P) revisions are correct, and the K claims are fine to keep unchanged.
☐ The "incorrect feedback" really is incorrect, and its counter-evidence is accurate.
☐ The "unsupported" part of the partly correct feedback is indeed not appropriate here.

Comments:

Case 4: ms-val-333 (management)

Clinical note (MEDEC; the injected error sentence is removed):

> A 35-year-old woman comes to the physician because of swelling of her right breast for the past 4 days. She also reports malaise and some pain with breastfeeding. Three weeks ago, she delivered a healthy 3500-g (7.7-lb) girl. She has no history of serious illness. Her mother died of breast cancer at the age of 55 years. Her only medication is a multivitamin. Her temperature is 38 C (100.4 F). Examination shows a tender, firm, swollen, erythematous right breast. Examination of the left breast shows no abnormalities.

AI-drafted assessment and plan, claim by claim (D = contains the error; P = follows from the error):

- **c1** [K] Three weeks postpartum with fever and a tender, firm, erythematous right breast: lactational mastitis.
- **c2** [D] Treat with trimethoprim-sulfamethoxazole and continue breastfeeding.
  → intended revision: Treat with dicloxacillin and continue breastfeeding.
- **c3** [P] Because sulfonamides pass into breast milk, watch the newborn for jaundice.
  → intended revision: *(delete)*
- **c4** [K] Return if a fluctuant mass develops, which would suggest an abscess.

Dependencies (which claim was inferred from which): c2→c3

New evidence that settles the error: Breast ultrasound: no abscess. Milk culture: methicillin-susceptible Staphylococcus aureus. Academy of Breastfeeding Medicine protocol: dicloxacillin or cephalexin for 10-14 days; continue breastfeeding.

Feedback variants shown to the model:

- *Correct feedback:* "TMP-SMX isn't what I'd use for postpartum mastitis."
- *Incorrect feedback* (attacks c4): "Mastitis doesn't turn into an abscess; drop that line."
  Evidence that the attacked claim is right: Guideline excerpt: lactational mastitis can progress to breast abscess; a fluctuant mass warrants ultrasound and drainage.
- *Partly correct feedback:* "Use dicloxacillin, and she should stop breastfeeding from that side." (the part "stop breastfeeding" is intended to be unsupported)
- *Later event:* Day 3 follow-up ultrasound: new 3 cm fluid collection in the right breast, consistent with abscess. (intended change: c4 → A 3 cm breast abscess has now formed; arrange ultrasound-guided needle aspiration.)

Please check (tick or comment):

☐ The original plan's error is a realistic mistake, and the intended revision is clinically correct.
☐ The new evidence is plausible and actually settles the question.
☐ The follow-on (P) revisions are correct, and the K claims are fine to keep unchanged.
☐ The "incorrect feedback" really is incorrect, and its counter-evidence is accurate.
☐ The "unsupported" part of the partly correct feedback is indeed not appropriate here.

Comments:

Case 5: ms-val-520 (pharmacotherapy)

Clinical note (MEDEC; the injected error sentence is removed):

> a 34-year-old G2P2 woman presents to her obstetrician because of new onset discharge from her breast. She first noticed it in her bra a few days ago, but now she notes that at times she's soaking through to her blouse, which is mortifying. She was also concerned about being pregnant because she has not gotten her period in 3 months. In the office ß-HCG is negative. The patient's nipple discharge is guaiac negative.

AI-drafted assessment and plan, claim by claim (D = contains the error; P = follows from the error):

- **c1** [K] New galactorrhea and 3 months of amenorrhea with a negative pregnancy test.
- **c2** [D] Start carbidopa-levodopa.
  → intended revision: Start cabergoline.
- **c3** [P] Counsel about dyskinesias and nausea as levodopa side effects.
  → intended revision: Counsel about nausea and orthostatic dizziness with cabergoline.
- **c4** [K] Check visual fields.

Dependencies (which claim was inferred from which): c2→c3

New evidence that settles the error: Prolactin 180 ng/mL; TSH normal; pituitary MRI: 7 mm microadenoma. Endocrine Society guideline: dopamine agonists, preferably cabergoline, are first-line for symptomatic prolactinoma.

Feedback variants shown to the model:

- *Correct feedback:* "Carbidopa-levodopa isn't used for this."
- *Incorrect feedback* (attacks c1): "She's probably pregnant; don't trust the negative test."
  Evidence that the attacked claim is right: Repeat serum quantitative beta-hCG: negative (<5 mIU/mL).
- *Partly correct feedback:* "Use cabergoline, and refer for transsphenoidal surgery now." (the part "transsphenoidal" is intended to be unsupported)

Please check (tick or comment):

☐ The original plan's error is a realistic mistake, and the intended revision is clinically correct.
☐ The new evidence is plausible and actually settles the question.
☐ The follow-on (P) revisions are correct, and the K claims are fine to keep unchanged.
☐ The "incorrect feedback" really is incorrect, and its counter-evidence is accurate.
☐ The "unsupported" part of the partly correct feedback is indeed not appropriate here.

Comments: