



Spinning Conformal Correlations

Mustafa Efe Özkara

efe.ozkara@metu.edu.tr

2372100

November 28, 2024

1. Introduction

1.1 Aim and Motivation

2. Methods

2.1 Embedding Formalism

2.2 Encoding Tensors as Functions

3. Summary

Operator Product Expansion (OPE) associativity is used only in 4-point CFT correlation functions of scalars.

Operator Product Expansion (OPE) associativity is used only in 4-point CFT correlation functions of scalars.

Aim: Extend correlation functions of scalars to spin- l operators and tensors.

Operator Product Expansion (OPE) associativity is used only in 4-point CFT correlation functions of scalars.

Aim: Extend correlation functions of scalars to spin- l operators and tensors.

Motivation: Analogy between CFT correlation functions in Mellin representations and scattering amplitudes

Points in d-dimensional space \rightarrow null vectors in d+2 dimensional space

$$P^A = \lambda(1, x^2, x^a)$$

Properties:

- Null vector on the lightcone $P^2 = 0$
- Homogeneity
- Symmetric and Traceless
- Transverse: $(P \cdot F)_{A_2 \dots A_I} \equiv P^A F_{AA_2 \dots A_I}$

Conformal groups \rightarrow Lorentz groups

$$\langle \Psi_1(P_1) \Psi_2(P_2) \Psi_3(P_3) \rangle = F(P_1, P_2, P_3)$$

1. $SO(d+1, 1)$ symmetry: F depends only on $(P_i - P_j)^2 = P_i^2 + P_j^2 - 2P_i \cdot P_j$.
2. On the null cone: $P_i^2 = 0$, hence $F(P_1, P_2, P_3) = F(P_{12}, P_{23}, P_{31})$,

$$P_{ij} = -2P_i \cdot P_j.$$

$$\langle \Psi_1(\lambda P_1) \Psi_2(P_2) \Psi_3(P_3) \rangle = \lambda^{-\Delta_1} \langle \Psi_1(P_1) \Psi_2(P_2) \Psi_3(P_3) \rangle,$$

$$F(\lambda P_{12}, P_{23}, P_{31}) = \lambda^{-\Delta_1} F(P_{12}, P_{23}, P_{31}),$$

$$F(\lambda P_{12}, \lambda^{-1} P_{23}, P_{31}) = \lambda^{\Delta_3 - \Delta_1} F(P_{12}, P_{23}, P_{31}).$$

$$F(P_{12}, P_{23}, P_{31}) = NP_{12}^a P_{23}^b P_{31}^c,$$

$$\langle \Psi_1(P_1) \Psi_2(P_2) \Psi_3(P_3) \rangle = \frac{N}{(P_{12})^{\frac{\Delta_1 + \Delta_2 - \Delta_3}{2}} (P_{23})^{\frac{\Delta_2 + \Delta_3 - \Delta_1}{2}} (P_{31})^{\frac{\Delta_3 + \Delta_1 - \Delta_2}{2}}}.$$

Projection to Physical Space: $f(x) = F(P_x), \quad P_{ij} \rightarrow x_{ij}^2.$

Tensors are converted to polynomials

$$f_{a_1 \dots a_I} \text{ symmetric } \leftrightarrow f(z) \equiv f_{a_1 \dots a_I} z^{a_1} \dots z^{a_I}$$

$$F_{A_1 \dots A_I}(P) \text{ symmetric } \leftrightarrow F(P, Z) \equiv F_{A_1 \dots A_I} Z^{A_1} \dots Z^{A_I}$$

$$f_{a_1 \dots a_l}(x) = \frac{\partial P^{A_1}}{\partial x^{a_1}} \dots \frac{\partial P^{A_l}}{\partial x^{a_l}} F_{A_1 \dots A_l}(P_x)$$



Thank You

for your attention.

Do you have any question?