

Essential Linear Algebra - Titu Andresscu

Contents

1 Matrix Algebra	2
1.1 Vectors, Matrices, and Basic Operations on Them	2
1.1.1 Summary	2
1.1.2 Exercises	3
1.2 Matrices as Linear Maps	4
1.2.1 Summary	4
1.2.2 Problems for Practice	4
1.3 Matrix Multiplication	6
1.3.1 Summary	6
1.3.2 Problems for Practice	6
1.4 Block Matrices	10
1.4.1 Summary	10
1.4.2 Problems for Practice	10
1.5 Invertible Matrices	10
1.5.1 Summary	10
1.5.2 Problems for Practice	10
1.6 The Transpose of a Matrix	13
1.6.1 Summary	13
1.6.2 Problems for Practice	13
 2 Square Matrices of Order 2	 15
2.1 The Trace and Determinant Maps	15
2.1.1 Summary	15
2.1.2 Problems for Practice	16
2.2 The Characteristic Polynomial and the Cayley-Hamilton Theorem	17
2.2.1 Summary	17
2.2.2 Problems for Practice	22
2.3 The Powers of a Square matrix of Order 2	24
2.3.1 Summary	24
2.3.2 Problems for Practice	25
2.4 Application to Linear Recurrences	26
2.4.1 Summary	26
2.4.2 Problems for Practice	27
2.5 Solving the Equation $X^n = A$	29
2.5.1 Summary	29
2.5.2 Problems for Practice	29
2.6 Application to Pell's Equations	30

2.6.1 Summary	30
2.6.2 Problems for Practice	30
3 Matrices and Linear Equations	31
3.1 Linear Systems: The Basic Vocabulary	31
3.1.1 Summary	31
3.1.2 Problems for Practice	32
3.2 The Reduced Row-Echelon Form and Its Relevance to Linear Systems	33
3.2.1 Summary	33
3.2.2 Problems for Practice	33
3.3 Solving the system $AX = b$	35
3.3.1 Sumamry	35
3.3.2 Problems for Practice	35
3.4 Computing the Inverse of a Matrix	35
3.4.1 Summary	35
3.4.2 Problems for Practice	36
4 Vector Spaces and Subspaces	36
4.1 Vector Spaces: Definintion, Basic Properties and Examples	36
4.1.1 Summary	36
4.1.2 Problems for Practice	36
4.2 Subspaces	37
4.2.1 Summary	37
4.2.2 Problems for Practice	38
4.3 Linear Combinations and Span	40
4.3.1 Summary	40
4.3.2 Problems for Practice	40
4.4 Linear Independence	40
4.4.1 Summary	40
4.4.2 Problems for Practice	41
4.5 Dimension Theory	43
5 Linear Transformations	43
5.1 Definitions and Objects Canonically Attached to a Linear Map	43
5.1.1 Summary	43
5.1.2 Problems for Practice	45

1 Matrix Algebra

1.1 Vectors, Matrices, and Basic Operations on Them

1.1.1 Summary

1.1.2 Exercises

Problem 1.1.6

Let F be a field.

- Prove that if $A, B \in M_n(\mathbb{F})$ are diagonal matrices, then $A + cB$ is a diagonal matrix for any $c \in \mathbb{F}$.
- Prove that the same result holds if we replace diagonal with upper-triangular.
- Prove that any matrix $A \in M_n(\mathbb{F})$ can be written as the sum of an upper-triangular matrix and of a lower-triangular matrix. Is there such a unique writing?

Solution.

- Since A and B are both diagonal matrices,

$$A + cB = \begin{bmatrix} a_{11} + cb_{11} & 0 & \cdots & 0 \\ 0 & a_{22} + cb_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} + cb_{nn} \end{bmatrix}$$

$\Rightarrow A + cB$ is also a diagonal matrix.

- Since A and B are both upper-triangular matrices,

$$A + cB = \begin{bmatrix} a_{11} + cb_{11} & a_{12} + cb_{12} & \cdots & a_{1n} + cb_{1n} \\ 0 & a_{22} + cb_{22} & \cdots & a_{2n} + cb_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} + cb_{nn} \end{bmatrix}$$

$\Rightarrow A + cB$ is also a diagonal matrix.

- We always have

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} = B + C,$$

where B is a lower-triangular matrix of b_{ij} -entries, C a upper-triangular matrix of c_{ij} -entries.

Now let's suppose that this way of writing isn't unique; in other words, we assume that $A = E + F$ ($E, F \in M_n(\mathbb{F})$) and E, F are upper-triangular and lower-triangular matrices, respectively. Thus $E + G = B + C \iff O_n = (B - E) + (C - G)$. This means that $b_{ij} = e_{ij}$, $c_{ij} = g_{ij}$, which implies that A can be

written as the sum of two unique matrices: one upper-triangular, the other lower-triangular.

□

Problem 1.1.7

- How many distinct matrices are there $M_{m,n}(\mathbb{F}_2)$?
- How many of these matrices are diagonal?
- How many of these matrices are upper-triangular?

Solution.

- Each entry in an $m \times n$ matrix can take on two values: 0 or 1. Since there are $m \times n$ entries, the total number of distinct $m \times n$ matrices are 2^{mn} for $m, n \in \mathbb{N}^*$. Thus the total number of distinct matrices is $\sum_{i=1}^m \sum_{j=1}^n 2^{ij}$.
- A diagonal matrix A has to be $M_n(\mathbb{F}_2)$, and $a_{ij} = 0, i \neq j$. We only need to consider the a_{ii} -entries, of which there are n entries. Using the same reasoning in (a), we get the number of $n \times n$ diagonal matrices: 2^n . Since n runs from $1 \rightarrow n$, we get the total number of distinct diagonal matrices: $\sum_{i=1}^n 2^i$.
- Similar to (b), we get the number of distinct upper-triangular matrices: $\sum_{i=1}^n 2^{\frac{i(i+1)}{2}}$.

□

1.2 Matrices as Linear Maps

1.2.1 Summary

1.2.2 Problems for Practice

Problem 1.2.2

Consider the map $f : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ defined by

$$f(x, y, z) = (x - 2y + 2z, y - z + x, x, z).$$

Prove that f is linear and describe the matrix associated with f .

Problem 1.2.3

a. Consider the map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$f(x, y) = (x^2, y^2).$$

Is this map linear?

b. Answer the same question with the field \mathbb{R} replaced with \mathbb{F}^2 .

Solution.

a. Let $X, Y \in \mathbb{R}^2 \wedge \alpha, \beta \in \mathbb{R}$. We then have

$$f(\alpha X + \beta Y) = \alpha^2 f(X) + \beta^2 f(Y) \neq \alpha f(X) + \beta f(Y)$$

$\Rightarrow f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is not a linear map.

b. If, however, $X, Y \in \mathbb{F}^2 \wedge \alpha, \beta \in \mathbb{F}$, we get

$$f(\alpha X + \beta Y) = \alpha^2 f(X) + \beta^2 f(Y) = \alpha f(X) + \beta f(Y)$$

since $\alpha, \beta \in \mathbb{F}$. It's also easy to prove that f is an injective map, thus f is linear \mathbb{F}^2 .

□

Problem 1.2.4

Consider the map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$f(x, y) = (x + 2y, x + y - 1).$$

Is the map f linear?

Problem 1.2.5

Consider the matrix $A = \begin{pmatrix} 1 & -2 & 2 & 0 \\ 2 & 0 & 4 & 1 \\ -1 & 1 & 0 & 1 \end{pmatrix}$. Describe the image of the vector $v = \begin{pmatrix} 1 \\ 1 \\ 2 \\ 2 \end{pmatrix}$ through the linear map attached to A .

Problem 1.2.6

Give an example of a map $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ which is *not* linear and for which

$$f(av) = af(v)$$

$\forall a \in \mathbb{R}$ and $v \in \mathbb{R}^2$.

1.3 Matrix Multiplication**1.3.1 Summary****1.3.2 Problems for Practice****Problem 1.3.5**

Determine all matrices $A \in M_2(\mathbb{R})$ commuting with the matrix

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}.$$

Problem 1.3.7 (Matrix representation of \mathbb{C})

Let G be the set of the matrices of the form $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ with $a, b \in \mathbb{R}$.

- Prove that the sum and product of two elements of G is in G .
- Consider the map $f : G \rightarrow \mathbb{C}$ defined by

$$f\left(\begin{bmatrix} a & -b \\ b & a \end{bmatrix}\right) = a + ib.$$

Prove that f is a bijective map satisfying $f(A + B) = f(A) + f(B)$ and $f(AB) = f(A)f(B) \forall A, B \in G$.

- Use this to compute the n th power of the matrix $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$.

Solution.

- Suppose $A, B \in G$ and $A = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}, B = \begin{bmatrix} c & -d \\ d & c \end{bmatrix}$.

$$A + B = \begin{bmatrix} a + c & -(b + d) \\ b + d & a + c \end{bmatrix} \Rightarrow A + B \in G.$$

$$AB = \begin{bmatrix} ac - bd & -(ad + bc) \\ ad + bc & ac - bd \end{bmatrix} \Rightarrow AB \in G.$$

- $f(A) + f(B) = f(A + B)$ and $f(A)f(B) = f(AB)$ are easily proven; we'll prove that f is bijective.

We'll reuse A and B from (a). Suppose that $f(A) = f(B) \Rightarrow \begin{cases} a = c \\ b = d \end{cases}$. Thus $A = B$ and f is injective.

Next, we observe that $\forall a, b \in \mathbb{R}, \exists X = \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \in G : f(X) = a + bi$. So f is surjective.

c. From (b), we get

$$\begin{aligned} f(A^n) &= [f(A)]^n = (a + bi)^n = (a^2 + b^2)^{\frac{n}{2}} \left(\frac{a}{a^2 + b^2} + \frac{b}{a^2 + b^2} i \right)^n \\ &= (a^2 + b^2)^{\frac{n}{2}} (\cos \theta + i \sin \theta)^n = (a^2 + b^2)^{\frac{n}{2}} (\cos n\theta + i \sin n\theta). \end{aligned} \quad (1.1)$$

Then, we obtain

$$A^n = (a^2 + b^2)^{\frac{n}{2}} \begin{pmatrix} \cos n\theta & -\sin n\theta \\ \sin n\theta & \cos n\theta \end{pmatrix},$$

where $\theta = \arccos \frac{a}{a^2 + b^2}$.

□

Problem 1.3.8

For any real number x let

$$A(x) = \begin{bmatrix} 1-x & 0 & x \\ 0 & 1 & 0 \\ x & 0 & 1-x \end{bmatrix}.$$

a. Prove that for all $a, b \in \mathbb{R}$, we have

$$A(a)A(b) = A(a + b - 2ab).$$

b. Given $x \in \mathbb{R}$, compute $A(x)^n$.

Problem 1.3.9

Compute A^{20} , where

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

Solution. The power of a diagonal matrix is the power of its diagonal entries, so

$$A^{20} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2^{20} & 0 \\ 0 & 0 & 3^{20} \end{bmatrix} \quad (1.2)$$

.

□

Problem 1.3.10

- a. Give a detailed proof, by induction on k , for the *binomial formula*: if $A, B \in M_n(\mathbb{F})$ commute then

$$(A + B)^k = \sum_{j=0}^k \binom{k}{j} A^{k-j} B^j.$$

- b. Give a counterexample to the binomial formula if we drop the hypothesis that A and B commute.

Solution.

- a. There's nothing to prove the base case $k = 0$, so assume that the binomial formula is true $\forall k = n \in \mathbb{N}$.

For $k = n + 1$,

$$\begin{aligned} (A + B)^{n+1} &= (A + B) \cdot (A + B)^n \\ &= \sum_{j=0}^n \binom{n}{j} A^{n-j} B^j (A + B) \\ &= A^{n+1} + (n+1)A^n B + \dots + (n+1)AB^n + B^{n+1} \\ &= \sum_{j=0}^{n+1} \binom{n+1}{j} A^{n+1-j} B^j. \end{aligned}$$

This closes the induction.

- b. The claim isn't necessarily true: let $A = \begin{bmatrix} 1 & 2 \\ 5 & 6 \end{bmatrix}, B = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$. Then $(A + B)^2 = \begin{bmatrix} 36 & 48 \\ 96 & 132 \end{bmatrix}$ and $\sum_{j=0}^2 \binom{2}{j} A^{2-j} B^j = \begin{bmatrix} 32 & 44 \\ 96 & 136 \end{bmatrix}$, so in this case, $(A + B)^2 \neq \sum_{j=0}^2 \binom{2}{j} A^{2-j} B^j$.

□

Problem 1.3.11

- a. Let

$$B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & -1 \\ 1 & 1 & 0 \end{bmatrix}.$$

Prove that $B^3 = O_3$.

- b. Let $a \in \mathbb{R}$ Using part (a) and the binomial formula, compute A^n where

$$A = \begin{bmatrix} 1 & 0 & a \\ 0 & 1 & -a \\ a & a & 1 \end{bmatrix}.$$

Problem 1.3.12

Let

$$A = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & \frac{1}{2} \\ 0 & 0 & 1 \end{bmatrix}.$$

- Prove that $(A - I_3)^3 = O_3$.
- Compute $A^n \forall n \in \mathbb{N}^*$, by using part (a) and the binomial formula.

Problem 1.3.11

- Prove that matrix

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

satisfies $(A - I_3)^3 = O_3$.

- Compute $A^n \forall n \in \mathbb{N}^*$.

Problem 1.3.16

A matrix $A \in M_n(\mathbb{R})$ is called a *permutation matrix* if each row and column of A has an entry equal to 1 and all other entries equal to 0. Prove that the product of two permutation matrices is a permutation matrix.

Problem 1.3.17

Consider the permutation σ of $1, 2, \dots, n$ that is a bijective map

$$\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}.$$

We define the associated permutation matrix P_σ as follows: the (i, j) -entry of P_σ is equal to 1 if $i = \sigma(j)$ and 0 otherwise.

- Prove that any permutation matrix is of the form P_σ for unique permutation σ .
- Deduce that there are $n!$ permutation matrices.
- Prove that

$$P_{\sigma_1} \cdot P_{\sigma_2} = P_{\sigma_1 \circ \sigma_2}.$$

for all permutations σ_1, σ_2 .

- Given a matrix $B \in M_n(\mathbb{F})$, describe the matrices $P_\sigma B$ and $B P_\sigma$ in terms of B and of the permutation σ .

1.4 Block Matrices

1.4.1 Summary

1.4.2 Problems for Practice

If $A = [a_{ij}] \in M_{m_1, n_1}(\mathbb{F})$ and $B \in M_{m_2, n_2}(\mathbb{F})$ are matrices, the *Kronecker product* or *tensor product* of A and B is the matrix $A \otimes B \in M_{m_1 m_2, n_1 n_2}(\mathbb{F})$ defined by

$$A \otimes B = \begin{bmatrix} a_{11}B & a_{12}B & \cdots & a_{1, n_1}B \\ a_{21}B & a_{22}B & \cdots & a_{2, n_1}B \\ \vdots & \vdots & \ddots & \vdots \\ a_{m_1, 1}B & a_{m_1, 2}B & \cdots & a_{m_1, n_1}B \end{bmatrix}.$$

Problem 1.4.1

Compute the Kronecker product of the matrices

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix}.$$

Solution: $A \otimes B = \begin{bmatrix} 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ 2 & 1 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 1 & -1 \end{bmatrix}.$

Problem 1.4.2

Do we always have $A \otimes B = B \otimes A$?

Problem 1.4.3

Check that $I_m \otimes I_n = I_{mn}$.

1.5 Invertible Matrices

1.5.1 Summary

1.5.2 Problems for Practice

Problem 1.5.3

Is the matrix

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix} \in M_3(\mathbb{F}_2)$$

invertible? If so, compute its inverse.

Problem 1.5.5

Consider the matrix

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \in M_5(\mathbb{R}).$$

Prove that A is invertible and compute its inverse.

Problem 1.5.6

Consider the matrix

$$A = \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}.$$

- Compute the inverse of A by solving, for each $b \in \mathbb{R}^4$, the system $AX = b$.
- Prove that $A^2 = 3I_4 + 2A$. Deduce a new way of computing A^{-1} .

Problem 1.5.7

Let A be the matrix

$$A = \begin{bmatrix} 3 & -1 & 2 \\ 5 & -2 & 3 \\ -1 & 0 & -1 \end{bmatrix}.$$

- Check that $A^3 = O_3$.
- Compute $(I_3 + A)^{-1}$.

Problem 1.5.10

Let $A, B \in M_{\mathbb{R}}$ such that

$$A + B = I_n \text{ and } A^2 + B^2 = O_n.$$

Prove that A and B are invertible and that $(A^{-1} + B^{-1})^n = 2^n I_n \forall n \in \mathbb{N}^*$.

Proof. We have

$$(A + B)^2 = A^2 + AB + BA + B^2 = AB + BA = I_n.$$

On the other hand,

$$A(A + B) = AI_n = (A + B)A = A^2 + AB = A^2 + BA.$$

Combining these two equations, we obtain $AB = BA \implies AB + BA = 2AB = 2BA = I_n \implies A$ has inverse and $A^{-1} = 2B$.

$(A^{-1} + B^{-1})^n = (A + B)^n (A^{-1} + B^{-1})^n = (2I_n + BA^{-1} + AB^{-1})^n = (2I_n + A^2 + B^2)^n = 2^n I_n$, as desired. \square

Problem 1.5.11

Let $A \in M_n(\mathbb{R})$ be invertible be an invertible matrix such that

$$A^{-1} = I_n - A.$$

Prove that $A^6 - I_n = O_n$.

Proof. So,

$$\begin{aligned} A^6 - I_n &= (A - I_n)(A + I_n)(A^2 + A + I_n)(A^2 - A + I_n) \\ &= B(A^2 - A + I_n) = BO_n = O_n, \end{aligned}$$

where $B = (A - I_n)(A^2 + A + I_n)(A + I_n)$, as desired. \square

Problem 1.5.14

Suppose that an upper-triangular matrix $A \in M_n(\mathbb{C})$ is invertible. Prove that A^{-1} is also upper-triangular.

Problem 1.5.15

Let $a, b, c \in \mathbb{R}_+$, not all of them equal and consider the matrix

$$A = \begin{bmatrix} a & 0 & b & 0 & c & 0 \\ 0 & a & 0 & c & 0 & b \\ c & 0 & a & 0 & b & 0 \\ 0 & b & 0 & a & 0 & c \\ b & 0 & c & 0 & a & 0 \\ 0 & c & 0 & b & 0 & a \end{bmatrix}.$$

Prove that A is invertible.

1.6 The Transpose of a Matrix**1.6.1 Summary****1.6.2 Problems for Practice****Problem 1.6.2**

Let $\theta \in \mathbb{R}$ and let

$$A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

- Prove that A is orthogonal.
- Find all values of θ for which A is symmetric.
- Find all values of θ for which A is skew-symmetric.

Problem 1.6.3

Which matrices $A \in M_n(\mathbb{F}_2)$ are the sum of a symmetric matrix and of a skew-symmetric matrix?

Problem 1.6.4

Write the matrix

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix}$$

and the sum of a symmetric matrix and of a skew-symmetric matrix with real entries.

Problem 1.6.5

All matrices in the following statements are square matrices of the same sizes. Prove that

- The product of two symmetric matrices is a symmetric matrix if and only if the two matrices commute.
- The product of two antisymmetric matrices is a symmetric matrix if and only if the two matrices commute.
- The product of a symmetric and a skew-symmetric matrix is a symmetric matrix if and only if the two matrices commute.

Problem 1.6.7

Consider the map $\varphi : M_3(\mathbb{R}) \longrightarrow M_3(\mathbb{R})$ defined by

$$\varphi(A) = A^T + 2A.$$

Prove that φ is linear, that is,

$$\varphi(cA + B) = c\varphi(A) + \varphi(B)$$

$$\forall A, B \in M_3(\mathbb{R}), c \in \mathbb{R}.$$

Problem 1.6.8

Let $A \in M_n(\mathbb{R})$ be a matrix such that $A \cdot A^T = O_n$. Prove that $A = O_n$.

Problem 1.6.9

Find all skew-symmetric matrices $A \in M_m(\mathbb{R})$ such that $A^2 = O_n$.

Proof. If a skew-symmetric matrix $A = [a_{ij}] \in M_n(\mathbb{R}) : A^2 = O_n$ then its diagonal entries must be zero, or

$$\begin{aligned} \sum_{i=1}^n a_{1i}^2 &= 0 \\ \sum_{i=1}^n a_{2i}^2 &= 0 \\ &\vdots \\ \sum_{i=1}^n a_{ni}^2 &= 0 \\ \Rightarrow a_{ij} &= 0 \quad \forall i, j \leq n \iff A = O_n. \end{aligned}$$

□

Problem 1.6.10

Let $a_1, \dots, a_k \in M_n(\mathbb{R})$ be matrices such that

$$\sum_{i=1}^k A_i \cdot A_i^T = O_n$$

Prove that $A_1 = \dots = A_k = O_n$.

Problem 1.6.11

- a. Let $A \in M_3(\mathbb{R})$ be a skew-symmetric matrix. Prove that there exists a non-zero vector $X \in \mathbb{R}^3 : AX = 0$.
- b. Does the result in part (a) remain true if we replace $M_3(\mathbb{R})$ with $M_2(\mathbb{R})$?

Proof.

- a. Suppose that $A \in M_3(\mathbb{R}) = \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}$. We then have four cases to $AX = 0$:
- $a = b = c = 0$: Any nonzero vector $X \in \mathbb{R}^3$ satisfies the problem.
 - One of a, b, c is 0. Then the value of the variable with the zero coefficient can be any real number, while the remaining two are 0.
 - One of a, b, c is 0. Then the value of the two variables with the zero coefficient can be any real number (but they can't be both 0), while the remaining one is 0.
 - $a, b, c \neq 0$. Then $x = \begin{bmatrix} c \\ -b \\ a \end{bmatrix}$.
- b. The result of (a) holds true only for $a = 0$; if $a \neq 0$, then $X = 0$.

□

Problem 1.6.12

Describe all upper-triangular matrices $A \in M_m(\mathbb{R})$ such that

$$A \cdot A^T = A^T \cdot A.$$

2 Square Matrices of Order 2

2.1 The Trace and Determinant Maps

2.1.1 Summary

Defintion 2.1

Consider a matrix $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in M_2(\mathbb{C})$. We define

- the *trace* of A is

$$\text{Tr}(A) = a_{11} + a_{22}.$$

- the *determinant* of A as

$$\det A = a_{11}a_{22} - a_{12}a_{21}.$$

We also write

$$\det A = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

for $\det A$.

Proposition 2.2

For all matrices $A, B \in M_2(\mathbb{C})$, $z \in \mathbb{C}$,

- (1) $\text{Tr}(A + zB) = \text{Tr}(A) + z \text{Tr}(B)$;
- (2) $\text{Tr}(AB) = \text{Tr}(BA)$;
- (3) $\text{Tr}(A^T) = \text{Tr}(A)$.

Proposition 2.4

For all matrices $A, B \in M_2(\mathbb{C})$ and all $\alpha \in \mathbb{C}$ we have

- (1) $\det(AB) = \det A \cdot \det B$;
- (2) $\det A^T = \det A$;
- (3) $\det(\alpha A) = \alpha^2 \det A$.

2.1.2 Problems for Practice**Problem 2.1.4**

Prove that $\forall A \in M_2(\mathbb{C})$, we have

$$\det A = \frac{(\text{Tr}(A))^2 - \text{Tr}(A^2)}{2}.$$

Problem 2.1.5

Prove that $\forall A, B \in M_2(\mathbb{C})$ we have

$$\det(A + B) = \det A + \det B + \text{Tr}(A) \cdot \text{Tr}(B) - \text{Tr}(AB).$$

Problem 2.1.6

Let $f : M_2(\mathbb{C}) \rightarrow \mathbb{C}$ be a map with the property that $\forall A, B \in M_2(\mathbb{C})$ and $z \in \mathbb{C}$, we have

$$f(A + zB) = f(A) + zf(B) \quad \text{and} \quad f(AB) = f(BA).$$

a. Consider the matrices

$$E_{11} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}, \quad E_{12} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad E_{21} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad E_{22} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

and define $x_{ij} = f(E_{ij})$. Check that $E_{12}E_{21} = E_{11}$ and $E_{21}E_{12} = E_{22}$, and deduce that $x_{11} = x_{22}$.

b. Check that $E_{11}E_{12} = E_{12}$ and $E_{12}E_{11} = O_2$, and deduce that $x_{12} = 0$. Using a similar argument, prove that $x_{21} = 0$.

c. Conclude that $\exists c \in \mathbb{C}$ such that

$$f(A) = c \cdot \text{Tr}(A)$$

\forall mts A .

2.2 The Characteristic Polynomial and the Cayley-Hamilton Theorem

2.2.1 Summary

The *characteristic polynomial* of A is the polynomial denoted

$$\det(XI_2 - A) = X^2 - \text{Tr}(A)X + \det A.$$

Note that AB and BA have the same characteristic polynomial for all $A, B \in M_2(\mathbb{C})$. In particular, if P is invertible, then A and PAP^{-1} have the same characteristic polynomial.

For any $z \in \mathbb{C}$, if we evaluate the characteristic polynomial of A at z , we know the value of $\det(zI_2 - A)$.

Worked Example 2.6

For any $A, B \in M_2(\mathbb{C})$, there is a complex number u such that

$$\det(A + zB) = \det A + uz + \det B \cdot z^2$$

$\forall z \in \mathbb{C}$ If A, B have integer/rational/real entries, so does u .

Proof. Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, B = \begin{bmatrix} e & f \\ g & h \end{bmatrix}$. Then,

$$\begin{aligned}
\det(A + zB) &= \det \begin{pmatrix} a + ze & b + zf \\ c + zg & d + zh \end{pmatrix} \\
&= ad - bc + z^2(eh - fg) + (ah + de - bg - cf)z \\
&= \det A + uz + \det B \cdot z^2,
\end{aligned}$$

where $u = ah + de - bg - cf$. □

Worked Example 2.7

Let $U, V \in M_2(\mathbb{R})$. Using the polynomial $\det(U + XV)$, prove that

$$\det(U + V) + \det(U - V) = 2 \det U + 2 \det V.$$

Proof. We have, from Worked Example ,

$$\begin{aligned}
&\det(U + V) + \det(U - V) \\
&= \det U + m + \det V + \det U - m + \det V \\
&= 2 \det U + 2 \det V
\end{aligned}$$

for some $m \in \mathbb{R}$. □

Worked Example 2.8

Let $A, B \in M_2(\mathbb{R})$. Using the previous problem, prove that

$$\det(A^2 + B^2) + \det(AB + BA) \geq 0.$$

Proof. From the previous problem, we can derive

$$\begin{aligned}
\det(A^2 + B^2) + \det(AB + BA) &= \det \left[\frac{(A + B)^2 + (A - B)^2}{2} \right] + \det \left[\frac{(A + B)^2 - (A - B)^2}{2} \right] \\
&= \det[(A + B)^2] + \det[(A - B)^2] \\
&= [\det(A + B)]^2 + [\det(A - B)]^2 \geq 0
\end{aligned}$$

as desired. □

Worked Example 2.9

Let $A, B \in M_2(\mathbb{R})$. Using the polynomial

$$f(X) = \det[I_2 + AB + X(BA - AB)],$$

prove that

$$\det\left(I_2 + \frac{2AB + 3BA}{5}\right) = \det\left(I_2 + \frac{3AB + 2BA}{5}\right).$$

Proof. We'll need to prove the more general result $f(X) = f(1 - X)$; the statement of the problem asks us to verify this for $X = \frac{2}{5}$.

Our claim can be proved by observing that

$$\begin{aligned} f(X) &= \det[I_2 + AB + X(BA - AB)] \\ &= 1 + u + \det[(1 - X)AB + XBA] \\ &= 1 + u + \det[AB - X(AB - BA)] \\ &= 1 + u + \det AB - Xv + X^2 \det(AB - BA) \\ &= 1 + u + \det BA - Xv + X^2 \det(BA - AB) \\ &= 1 + u + \det[(1 - X)BA + XAB] \\ &= \det[I_2 + BA + X(AB - BA)] \\ &= f(1 - X) \text{ for some } u, v \in \mathbb{R}, \end{aligned}$$

as desired. Applying this result to $X = \frac{2}{5}$ gives us the result we want. \square

Defintion 2.10 (Eigenvalues)

The *eigenvalues* of a matrix $A \in M_2(\mathbb{C})$ are the roots of its characteristic polynomial in other words, they are the complex solutions λ_1, λ_2 of the equation

$$\det(tI_2 - A) = t^2 - \text{Tr}(A)t + \det A = 0.$$

Theorem 2.11 (Cayley-Hamilton theorem)

For any $A \in M_2(\mathbb{C})$, we have

$$A^2 - \text{Tr}(A) \cdot A + (\det A) \cdot I_2 = O_2.$$

Proof. For $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we have

$$\begin{aligned}
& A^2 - \text{Tr}(A) \cdot A + (\det A) \cdot I_2 = O_2 \\
&= \begin{bmatrix} a^2 + bc & ab + bd \\ ac + dc & bc + d^2 \end{bmatrix} - \begin{bmatrix} a^2 + ad & ab + ad \\ ac + cd & ad + d^2 \end{bmatrix} - \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} \\
&= O_2.
\end{aligned}$$

□

Remark 2.12

a. In other words, the matrix A is a solution of the characteristic polynomial

$$\det(tI_2 - A) = t^2 - \text{Tr}(A)t + \det A = 0.$$

b. Expressed in terms of the of the eigenvalues λ_1, λ_2 of A , the Cayley-Hamilton theorem can written either as

$$A^2 - (\lambda_1 + \lambda_2)A + \lambda_1\lambda_2 \cdot I_2 = O_2 \quad (2.1)$$

or equivalently

$$(A - \lambda_1 I_2)(A - \lambda_2 I_2) = O_2. \quad (2.2)$$

Worked Example 2.13

Let $A \in M_2(\mathbb{C})$ have eigenvalues λ_1 and λ_2 . Prove that $\forall n \geq 1$ we have

$$\text{Tr}(A^n) = \lambda_1^n + \lambda_2^n.$$

Deduce that λ_1^n and λ_2^n are eigenvalues of A^n .

Proof. Let $x_n = \text{Tr}(A^n)$. Multiplying (2.1) by A^n and taking its trace, we get

$$x_{n+2} - (\lambda_1 + \lambda_2)x_{n+1} + \lambda_1\lambda_2 x^n = 0.$$

□

Since $x_0 = 2$ and $x_1 = \lambda_1 + \lambda_2 = \text{Tr}(A)$, a simple induction shows that $\text{Tr}(A^n) = \lambda_1^n + \lambda_2^n$.

Worked Example 2.14

Let $A \in M_2(\mathbb{C})$ be a matrix with $\text{Tr}(A) \neq 0$. Prove that a matrix $B \in M_2(\mathbb{C})$ commutes with A if and only if B commutes with A^2 .

Worked Example 2.15

Prove that for any matrices $A, B \in M_2(\mathbb{R})$ there is a real a such that $(AB - BA)^2 = aI_2$.

Worked Example 2.16

Let $X \in M_2(\mathbb{R})$ be a matrix such that $\det(X^2 + I_2) = 0$. Prove that $X^2 + I_2 = O_2$.

Worked Example 2.17

A matrix $A \in M_2(\mathbb{C})$ is invertible if and only if $\det A \neq 0$. If this is the case, then

$$A^{-1} = \frac{1}{\det A} [\text{Tr}(A) \cdot I_2 - A].$$

Worked Example 2.19

Let $A, B \in M_2(\mathbb{C})$ be two matrices such that $AB = I_2$. Then A is invertible and $B = A^{-1}$. In particular, we have $BA = I_2$.

Theorem 2.20

If $A \in M_2(\mathbb{C})$ and $z \in \mathbb{C}$, then the following assertions are equivalent:

- z is an eigenvalue of A ;
- $\det(zI_2 - A) = 0$;
- There is a nonzero vector $v \in \mathbb{C}^2$ such that $Av = zv$.

Worked Example 2.21

Let $A \in M_2(\mathbb{C})$ have two distinct λ_1, λ_2 . Prove that we can find an invertible matrix $P \in \text{GL}_2(\mathbb{C})$ such that

$$A = P \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} P^{-1}.$$

Worked Example 2.22

Solve in $M_2(\mathbb{C})$ the following equations:

- $A^2 = O_2$;
- $A^2 = I_2$;
- $A^2 = A$.

Worked Example 2.23

Let $A \in M_2(\mathbb{C})$ be a matrix. Prove that the following statements are equivalent:

- a. $\text{Tr}(A) = \det A = 0$.
- b. $A^2 = O_2$.
- c. $\text{Tr}(A) = \text{Tr}(A^2) = 0$.
- d. $\exists n \geq 2 : A^n = O_2$.

Worked Example 2.24

Find all matrices $X \in M_2(\mathbb{R}) : X^3 = I_2$.

2.2.2 Problems for Practice**Problem 2.2.1**

Let $A, B \in M_2(\mathbb{R})$ be commuting matrices. Prove that

$$\det(A^2 + B^2) \geq 0$$

Proof.

$$\begin{aligned} \det(A^2 + B^2) &= \det(A + iB) \cdot \det(A - iB) \\ &= (\det A - \det B + ui)(\det A - \det B - ui) \\ &= (\det A - \det B)^2 + u^2 \text{ for some } u \in \mathbb{R} \end{aligned}$$

□

The last equation clearly shows that $\det(A^2 + B^2) \geq 0$.

Problem 2.2.2

Let $A, B \in M_2(\mathbb{R})$ such that $AB = BA$ and $\det(A^2 + B^2) = 0$. Prove that $\det A = \det B$.

Hint: use the hint of the previous problem and consider the polynomial $\det(A + XB)$.

Proof. Since $\det(A + iB) = \overline{\det(A - iB)}$ and $\det(A^2 + B^2) = 0$, this means that $\det(A + iB) = \det(A - iB) = 0 \implies \det A + iu - \det B = \det A - iu - \det B = 0 \implies \det A - \det B = 0 \implies \det A = \det B$.

□

Problem 2.2.3

Let $A, B, C \in M_2(\mathbb{R})$ be pairwise commuting matrices and let

$$f(X) = \det(A^2 + B^2 + C^2 + X[AB + BA + CA]).$$

- Prove that $f(2) \geq 0$.
- Prove that $f(-1) \geq 0$.
- Deduce that

$$\det(A^2 + B^2 + C^2) + 2 \det(AB + BC + CA) \geq 0.$$

Solution.

a.

□

Problem 2.2.4

Let $A, B \in M_2(\mathbb{C})$ be matrices with $\text{Tr}(AB) = 0$. Prove that $(AB)^2 = (BA)^2$.

Problem 2.2.5

Let $A \in M_2(\mathbb{Q})$ with the property that

$$\det(A^2 - 2I_2) = 0.$$

Prove that $A^2 = 2I_2$ and $\det A = -2$.

Problem 2.2.6

Let $x \in \mathbb{R}_+$ and let $A \in M_2(\mathbb{R})$ such that $\det(A^2 + xI_2) = 0$. Prove that

$$\det(A^2 + A + xI_2) = x.$$

Problem 2.2.7

Let $A, B \in M_2(\mathbb{R})$ be such that $\det(AB - BA) \leq 0$. Consider the polynomial

$$f(X) = \det(I_2 + [1 - X]AB + XBA).$$

- Prove that $f(0) = f(1)$.
- Deduce that

$$\det(I_2 + AB) \leq \det\left(I_2 + \frac{1}{2}[AB + BA]\right).$$

Problem 2.2.8

Let $n \geq 3$ be an integer. Let $X \in M_2(\mathbb{R})$ be such that

$$X^n + X^{n-2} = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}.$$

- a. Prove that $\det X = 0$.
- b. Let $t = \text{Tr}(X)$. Prove that

$$t^n + t^{n-2} = 2.$$

- c. Find all possible matrices X satisfying the original equation.

2.3 The Powers of a Square matrix of Order 2**2.3.1 Summary****Theorem 2.25**

Let $A \in M_2(\mathbb{C})$ and let λ_1, λ_2 be its eigenvalues.

- a. If $\lambda_1 \neq \lambda_2$, then $\forall n \geq 1$ we have $A^n = \lambda_1^n B + \lambda_2^n C$, where

$$B = \frac{1}{\lambda_1 - \lambda_2}(A - \lambda_2 I_2) \text{ and } C = \frac{1}{\lambda_2 - \lambda_1}(A - \lambda_1 I_2).$$

- b. If $\lambda_1 = \lambda_2$, then $\forall n \geq 1$ we have $A^n = \lambda_1^n I_2 + n\lambda_1^{n-1}(A - \lambda_1 I_2)$.

Worked Example 2.26

Compute A^n , where $A = \begin{bmatrix} 1 & 3 \\ -3 & -5 \end{bmatrix}$.

Corollary 2.27

For any matrix $A \in M_2(\mathbb{C})$ there are sequences $(x_n)_{n \geq 0}, (y_n)_{n \geq 0}$ of complex numbers such that

$$A^n = x_n A + y_n I_2.$$

Worked Example 2.28

Let m, n be positive integers and let $A, B \in M_2(\mathbb{C})$ be two matrices such that $A^m B^n = B^n A^m$. If A^m and B^n are not scalar, prove that $AB = BA$.

Worked Example 2.29

Let $t \in \mathbb{R}$ and let

$$A_t = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}.$$

Compute A_t^n for $n \geq 1$.

2.3.2 Problems for Practice**Problem 2.3.1**

Consider the matrix

$$\begin{bmatrix} 2 & 3 \\ 3 & 2 \end{bmatrix}.$$

- a. Let n be a positive integer. Prove the existence of a unique pair of integers $(x_n, y_n) \in \mathbb{Z}$ such that

$$A^n = x_n A + y_n I_2.$$

- b. Compute $\lim_{n \rightarrow \infty} \frac{x_n}{y_n}$.

Problem 2.3.2

Given a positive integer n , compute the n th power of the matrix

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.$$

Problem 2.3.3

Let $a, b \in \mathbb{R}$ and $n \in \mathbb{N}^*$. Compute the n th power of the matrix $\begin{bmatrix} a & b \\ 0 & a \end{bmatrix}$.

Problem 2.3.4

Let $x \in \mathbb{R}$ and let

$$A = \begin{bmatrix} \cos x + \sin x & 2 \sin x \\ -\sin x & \cos x - \sin x \end{bmatrix}.$$

Compute A^n for all positive integers n .

2.4 Application to Linear Recurrences

2.4.1 Summary

Let a, b, c, d, x_0, y_0 be complex numbers and consider the sequences x_n, y_n defined by

$$\begin{cases} x_{n+1} = ax_n + by_n \\ y_{n+1} = cx_n + dy_n \end{cases}, n \geq 0.$$

Noting that $\begin{bmatrix} x_{n+1} \\ y_{n+1} \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x_n \\ y_n \end{bmatrix} \forall n \geq 0$, we yield, via induction,

$$\begin{bmatrix} x_n \\ y_n \end{bmatrix} = A^n \begin{bmatrix} x_0 \\ y_0 \end{bmatrix}$$

where $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

For second-order linear recurrences with constant coefficients

$$x_{n+1} = ax_n + bx_{n-1} \quad n \geq 1,$$

we can reduce it back to the first case by letting $y_n = x_{n-1}$ for $n \geq 1$ and $y_0 = b^{-1}(x_1 - ax_0)$ if $b \neq 0$. Indeed, we then get the following system

$$\begin{cases} x_{n+1} = ax_n + by_n \\ y_{n+1} = x_n \end{cases}, \quad n \geq 0.$$

Finding x_n is now a matter of computing the powers of the matrix

$$A = \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix}.$$

The characteristic polynomial of this matrix is $\lambda^2 - a\lambda - b = 0$. If λ_1, λ_2 are the roots of this equation, then Theorem 2.25 allows us to yield the following:

- If $\lambda_1 \neq \lambda_2$, we can find constants u, v such that

$$x_n = u\lambda_1^n + v\lambda_2^n \quad \forall n.$$

They can be easily computed by solving for u, v in

$$\begin{cases} x_0 = u + v \\ x_1 = u\lambda_1 + v\lambda_2 \end{cases}.$$

- If $\lambda_1 = \lambda_2$, we can find u, v such that $\forall n \geq 0$,

$$x_n = (un + v)\lambda_1^n$$

and u, v can be found by solving

$$\begin{cases} x_0 = v \\ x_1 = (u + v)\lambda_1 \end{cases}.$$

2.4.2 Problems for Practice

Problem 2.4.1

Find the general term of the sequence $(x_n)_{n \geq 0}$ defined by $x_1 = 1, x_2 = 0$ and for $n \geq 1$,

$$x_{n+2} = 4x_{n+1} - x_n.$$

Solution. Letting $y_{\{n+1\}} = x_n$, we obtain

$$\begin{cases} x_{n+1} = 4x_n - x_{n-1} \\ y_{n+1} = x_n \end{cases}$$

Calculating the eigenvalues of the matrix $A = \begin{bmatrix} 4 & -1 \\ 1 & 0 \end{bmatrix}$, we get

$$\lambda_1 = 2 + \sqrt{3}, \lambda_2 = 2 - \sqrt{3}.$$

Thus there are constants u, v such that $x_n = u\lambda_1^n + v\lambda_2^n$ and

$$\begin{cases} x_0 = -4 = u + v \\ x_1 = 1 = u\lambda_1 + v\lambda_2 \end{cases}$$

$$\Rightarrow u = \frac{3\sqrt{3} - 4}{2} - 2, v = \frac{-3\sqrt{3} - 4}{2}$$

$$\Rightarrow x_n = \frac{(3\sqrt{3} - 4)(2 + \sqrt{3})^n}{2} - \frac{(3\sqrt{3} + 4)(2 - \sqrt{3})^n}{2}$$

□

Problem 2.4.2

Consider the sequence $(x_n)_{n \geq 0}$ defined by $x_0 = 1, x_1 = 2$ and for $n \geq 1$,

$$x_{n+2} = x_{n+1} - x_n.$$

Is this series periodical? If so, find its minimal period.

Solution. Setting $y_n = x_{n+1}$, we get

$$\begin{cases} x_{n+1} = x_n - x_{n-1} \\ y_{n+1} = x_n \end{cases}.$$

Solving for the eigenvalues of the matrix $A = \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}$, we get $\lambda_1 =$

$\frac{1 + \sqrt{3}i}{2}, \lambda_2 = \frac{1 - \sqrt{3}i}{2}$. This means that there exists constants, u, v such that

$$x_n = u\lambda_1^n + v\lambda_2^n.$$

. Solving the system $\begin{cases} 1 = u + v \\ 2 = u\lambda_1 + v\lambda_2 \end{cases}$, we obtain $u = \frac{1 - \sqrt{3}i}{2}$, $v = \frac{1 + \sqrt{3}i}{2}$. Thus

$$x_n = \frac{(1 - \sqrt{3}i)(1 + \sqrt{3}i)^n}{2^{n+1}} + \frac{(1 + \sqrt{3}i)(1 - \sqrt{3}i)^n}{2^{n+1}}.$$

□

Problem 2.3.3

Find the general terms of the sequences $(x_n)_{n \geq 0}$, $(y_n)_{n \geq 0}$ satisfying $x_0 = y_0 = 1$, $x_1 = 1$, $y_1 = 2$ and

$$x_{n+1} = \frac{2x_n + 3y_n}{5}, y_{n+1} = \frac{2y_n + 3x_n}{5}.$$

Problem 2.4.4

A sequence $(x_n)_{n \geq 0}$ satisfies $x_0 = 2$, $x_1 = 3$ and $\forall n \geq 1$,

$$x_{n+1} = \sqrt{x_{n-1}x_n}.$$

Find the general term of this sequence.

Solution. Taking the logarithm base 2 of this recurrence relation gives us

$$\lg x_{n+1} = \frac{1}{2} \lg x_n + \frac{1}{2} \lg x_{n-1}.$$

Similarly to the above questions, if we take $y_n = \lg x_{n-1}$ and $z_n = \lg x_n$, we get

$$\begin{cases} z_{n+1} = 0.5z_n + 0.5z_{n-1} \\ y_{n+1} = z_n \end{cases}$$

□

The eigenvalues of the matrix $A = \begin{bmatrix} 0.5 & 0.5 \\ 1 & 0 \end{bmatrix}$ are $\lambda_1 = \frac{1 + \sqrt{7}i}{4}$, $\lambda_2 = \frac{1 - \sqrt{7}i}{4}$. Then, we must have

$$x_n = u\lambda_1^n + v\lambda_2^n.$$

Solving for $\begin{cases} 2 = u + v \\ 3 = \lambda_1 u + \lambda_2 v \end{cases}$, we obtain $u = 1 - \frac{5\sqrt{7}}{7}i$, $v = 1 + \frac{5\sqrt{7}}{7}i$.

Problem 2.4.5

Consider a map $f : (0, \infty) \rightarrow (0, \infty)$ such that

$$f(f(x)) = 6x - f(x)$$

$\forall x > 0$. Let $x > 0$ and define a sequence $(z_n)_{n \geq 0}$ by $z_0 = x$ and $z_{n+1} = f(z_n)$ for $n \geq 0$.

a. Prove that

$$z_{n+2} + z_{n+1} - 6z_n = 0 \quad \forall n \geq 0.$$

b. Deduce the existence of $a, b \in \mathbb{R}$ such that

$$z_n = a \cdot 2^n + b \cdot (-3)^n \quad \forall n \geq 0.$$

c. Using the fact that $z_n > 0 \quad \forall n$, prove that $b = 0$ and conclude that $f(x) = 2x \quad \forall x > 0$.

2.5 Solving the Equation $X^n = A$ **2.5.1 Summary****2.5.2 Problems for Practice****Problem 2.5.1**

Let $n > 1 \in \mathbb{N}^*$. Prove that the equation

$$X^n = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

has no solutions in $M_2(\mathbb{C})$.

Problem 2.5.2

Solve in $M_2(\mathbb{C})$ the binomial equation

$$X^4 = \begin{bmatrix} -1 & -2 \\ 1 & 2 \end{bmatrix}.$$

Problem 2.5.1

Let $n > 1 \in \mathbb{N}^*$. Prove that the equation

$$X^n = \begin{bmatrix} 3 & -1 \\ 0 & 0 \end{bmatrix}$$

has no solutions in $M_2(\mathbb{Q})$.

Problem 2.5.4

Find all matrices $X \in M_2(\mathbb{R})$ such that

$$X^3 = \begin{bmatrix} 4 & 3 \\ -3 & -2 \end{bmatrix}.$$

Problem 2.5.5

Find all matrices $A, B \in M_2(\mathbb{C})$ such that

$$AB = O_2 \quad \text{and} \quad A^5 + B^5 = O_2.$$

Problem 2.5.6

Solve in $M_2(\mathbb{R})$ the binomial equation

$$X^n = \begin{bmatrix} 7 & -5 \\ -15 & 12 \end{bmatrix}.$$

Problem 2.5.7

Solve in $M_2(\mathbb{R})$ the binomial equation

$$X^n = \begin{bmatrix} -6 & -2 \\ 21 & 7 \end{bmatrix}.$$

2.6 Application to Pell's Equations**2.6.1 Summary****2.6.2 Problems for Practice****Problem 2.6.1**

A triangular number is a number of the form $1 + 2 + \dots + n$ for some positive integer n . Find all triangular numbers which are perfect squares.

Problem 2.6.2

Find all positive integers n such that $n + 1$ and $3n + 1$ are simultaneously perfect squares.

Problem 2.6.3

Find all integers a, b such that $a^2 + b^2 = 1 + 4ab$.

Problem 2.6.4

The difference of two consecutive cubes equals n^2 for some positive integer n . Prove that $2n - 1$ is a perfect square.

Problem 2.6.5

Find all triangles whose side lengths are consecutive integers and whose area is also an integer.

3 Matrices and Linear Equations

3.1 Linear Systems: The Basic Vocabulary

3.1.1 Summary

A linear equation in the variables x_1, \dots, x_n is an equation of the form

$$\sum_{i=1}^n a_i x_i = b,$$

where $a_1, \dots, a_n, b \in \mathbb{F}$ are given scalars and n is a given positive integer. The unknowns x_1, \dots, x_n are supposed to be elements in \mathbb{F} . A *linear system in the variables* x_1, \dots, x_n is a family of linear equations. It can also be written as

$$AX = b \tag{1}$$

where $A = \begin{bmatrix} x_{11} & \dots & x_{1n} \\ \vdots & \ddots & \vdots \\ x_{n1} & \dots & x_{nn} \end{bmatrix}$, $X = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$, $b = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$. If C_1, \dots, C_n are the columns of A , (1) is equivalent to

$$\sum_{i=1}^n x_i C_i = b. \tag{2}$$

Defintion 3.1

- The linear system is called *homogenous* if $b_1 = \dots = b_m = 0$.
- The homogeneous linear system associated with the system (1) is the system $AX = 0$.

Proposition 3.2 (Superposition principle)

Let $A \in M_{m,n}(F)$ and $b \in \mathbb{F}^m$. Let $S \subset \mathbb{F}^n$ be the set of solutions of the homogeneous linear system $AX = 0$. If the system $AX = b$ has a solution X_0 , then the set of solutions of this systems is $X_0 + S$.

Defintion 3.13

A linear system is called *consistent* if it has at least one solution and is *inconsistent* otherwise.

Defintion 3.4

- Two linear systems are *equivalent* if they have exactly the same set of solutions.
- If A, B are matrices of the same size and are equivalent, we write $A \sim B$.

3.1.2 Problems for Practice**Problem 3.1.3**

Let $a, b \in \mathbb{R}$, not both equal to 0.

- Prove that the system

$$\begin{cases} ax_1 + bx_2 = 0 \\ -bx_1 + ax_2 = 0 \end{cases}$$

has only the trivial solution.

- Prove that $\forall c, d \in \mathbb{R}$, the system

$$\begin{cases} ax_1 + bx_2 = c \\ -bx_1 + ax_2 = d \end{cases}$$

has a unique solution and find this solution in terms of a, b, c, d .

Solution:

- Beacause the system is homogeneous, it has the trivial solution.

Supposing that this system has at least one non-trivial solution.

If one a, b is equal to 0 (without loss of generality, let $b = 0$), then $ax_1 = ax_2 = 0 \implies x_1 = x_2 = 0$. If $a, b \neq 0$, then we have $x_2/x_1 = \frac{b}{a} = -\frac{a}{b}$, a contradiction. In either case, each result gives a contradiction. Thus $\begin{cases} ax_1 + bx_2 = 0 \\ -bx_1 + ax_2 = 0 \end{cases}$ only has the trivial solution.

- The system can be written as $\begin{bmatrix} a & b \\ -b & a \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} c \\ d \end{bmatrix}$. $\det A > 0 \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \frac{ac-bd}{a^2+b^2} \\ \frac{ad+bc}{a^2+b^2} \end{bmatrix}$.

Problem 3.1.4

Let $A \in M_2(\mathbb{C})$ be a matrix and consider the homogeneous system $AX = 0$. Prove that the following statements are equivalent:

- This system has only the trivial solution.
- A is invertible.

Solution. Since A is invertible, we get $X = A^{-1}AX = 0$, so the homogeneous system only has the trivial solution. Now assume that A only has the trivial solution but is also not invertible. Writing $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we must have $ad - bc = 0$. If $a = 0 \implies bc = 0$. If $c = 0$ then $X = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a non-trivial solution to $AX = 0$. If $c \neq 0 \implies b = 0 \implies X = \begin{bmatrix} -d \\ c \end{bmatrix}$.

If $a \neq 0 \implies d = \frac{bc}{a} \implies A = \begin{bmatrix} -b & a \\ a & a \end{bmatrix}$. As such, for each case we have a non-trivial solution to $AX = 0$, a contradiction to our assumption. Hence A is invertible. \square

Problem 3.1.5

Let A, B be $n \times n$ matrices such that the system $ABX = 0$ has only the trivial solution. Show that the system $BX = 0$ also has only the trivial solution.

Problem 3.1.6

Let C, D be $n \times n$ such that the system $CDX = b$ is consistent for every choice of $b \in \mathbb{R}^n$. Show that the system $CY = b$ is consistent for every choice of $b \in \mathbb{R}^n$.

3.2 The Reduced Row-Echelon Form and Its Relevance to Linear Systems

3.2.1 Summary

3.2.2 Problems for Practice

Problem 3.2.3

Determine the fundamental solutions of the homogeneous linear system of equations $AX = 0$, where

$$A = \begin{bmatrix} 1 & -2 & 1 & 0 \\ -2 & 4 & 0 & 2 \\ -1 & 2 & 1 & 2 \end{bmatrix}.$$

Solution. $A = \begin{bmatrix} 2a+b \\ -a \\ -b \\ b \end{bmatrix}$, where $a, b \in \mathbb{R}$. □

Problem 3.2.4

- a. Write the solutions of the homogeneous system of equations $AX = 0$ in parametric vector form, where

$$A = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & -1 \\ -1 & 1 & 4 \end{bmatrix}.$$

- b. Find a solution to the system for which the sum of the first two coordinates is 1.

Solution.

a. $X = \begin{bmatrix} -3a \\ -7a \\ a \end{bmatrix}.$

b. $-10a = 1 \Rightarrow a = -0.1 \Rightarrow X = \begin{bmatrix} 0.3 \\ 0.7 \\ -0.1 \end{bmatrix}.$ □

Problem 3.2.7

Let $n > 2 \in \mathbb{Z}$. Solve in \mathbb{R} the linear system

$$x_2 = \frac{x_1 + x_3}{2}, x_3 = \frac{x_2 + x_4}{2}, \dots, x_{n-1} = \frac{x_{n-2} + x_n}{2}.$$

Solution. We'll prove the following:

Claim

If $X = \begin{bmatrix} a_1 \\ a_2 \\ x_3 \\ \vdots \\ x_{n-1} \\ x_n \end{bmatrix}$, then

$$x_n = (n-1)a_2 - (n-2)a_1 \quad \text{for } n \geq 3.$$

Indeed, $x_3 = 2a_2 - a_1$, so assume the claim is true $\forall n \geq 3$. For $n+1$, $x_n = \frac{x_{n-1} + x_{n+1}}{2} \implies x_{n+1} = 2x_n - x_{n-1} = 2[(n-1)a_2 - (n-2)a_1] - (n-2)a_2 + (n-3)a_1 = na_2 - (n-1)a_1$. Thus we've proved our claim via induction, and

$$A = \begin{bmatrix} a_1 \\ a_2 \\ 2a_2 - a_1 \\ \vdots \\ (n-1)a_2 - (n-2)a_1 \end{bmatrix} \quad (a_1, a_2 \in \mathbb{R}).$$

□

3.3 Solving the system $AX = b$

3.3.1 Summary

3.3.2 Problems for Practice

Problem 3.2.6

Find all possible values of h and k such that the system with augmented matrix

$$\left[\begin{array}{cc|c} 1 & 2 & h \\ 2 & k & 12 \end{array} \right]$$

has

- a. a unique solution.
- b. infinitely many solutions.
- c. no solution.

Problem 3.3.8

Let $a, b \in \mathbb{R}$. Solve in real numbers the system

$$\begin{cases} x + y = a \\ y + z = b \\ z + t = a \\ t + x = b \end{cases}.$$

3.4 Computing the Inverse of a Matrix

3.4.1 Summary

3.4.2 Problems for Practice

Problem 3.4.2

For which $x \in \mathbb{R}$ is the matrix

$$A = \begin{bmatrix} 1 & x & 1 \\ 0 & 1 & x \\ 1 & 0 & 1 \end{bmatrix}$$

invertible? For any such number x , compute the inverse of A .

Problem 3.3.3

Let $x, y, z \in \mathbb{R}$. Compute the inverse of the matrix

$$A = \begin{bmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix}.$$

Problem 3.4.4

Determine the inverse of the matrix

$$A = \begin{bmatrix} n & 1 & 1 & \cdots & 1 \\ 1 & n & 1 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & \cdots & n \end{bmatrix} \in M_n(\mathbb{R}).$$

Problem 3.4.5

Let $a \in \mathbb{R}$. Determine the inverse of the matrix

$$A = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ a & 1 & 0 & \cdots & 0 & 0 \\ a^2 & a & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ a^{n-2} & a^{n-3} & a^{n-4} & \cdots & 1 & 0 \\ a^{n-1} & a^{n-2} & a^{n-3} & \cdots & a & 1 \end{bmatrix} \in M_m(\mathbb{R}).$$

4 Vector Spaces and Subspaces

4.1 Vector Spaces: Definition, Basic Properties and Examples

4.1.1 Summary

4.1.2 Problems for Practice

Problem 4.1.1

Consider the set $V = \mathbb{R}^2$ endowed with addition rule defined by

$$(x, y) + (x', y') = (x + x', y + y')$$

and with a multiplication rule by elements λ of \mathbb{R} as follows:

$$\lambda \cdot (x, y) = (2x, 0).$$

Is V endowed with these operations a vector space over \mathbb{R} ?

Solution. No V isn't a \mathbb{R} -vector space as $\forall v = (x, y) \in V, 1(x, y) = (x, 0) \neq (x, y) = v$. □

Problem 4.1.2

Define an operation $+$ on $(0, \infty)$ by

$$a +_* b = ab$$

for $a, b \in (0, \infty)$ and an external multiplication by real numbers as follows:

$$a \cdot_* b = b^a$$

for $a \in \mathbb{R}, b \in (0, \infty)$. Does $(0, \infty)$ endowed with this new addition and scalar multiplication become a vector space over \mathbb{R} ?

Problem 4.1.3 (Complexification of a real vector space)

Let V be a vector space over \mathbb{R} . Let $V_{\mathbb{C}}$ be the Cartesian product $V \times V$ endowed with the following addition rule:

$$(x, y) + (x', y') = (x + x', y + y').$$

Also, for each $z = a + bi \in \mathbb{C}$, consider the "multiplication by z rule"

$$z \cdot (x, y) = (ax - by, ay + bx)$$

on $V_{\mathbb{C}}$. Prove that $V_{\mathbb{C}}$ endowed with these operations becomes a \mathbb{C} -vector space (this space is called the *complexification* of the vector space V).

4.2 Subspaces**4.2.1 Summary**

Defintion 4.5

Let V be an \mathbb{F} -vector space. A *subspace* of V is a nonempty subset W of V which is stable under the operations of addition and scalar multiple: $v + w \in W$ and $cv \in W \forall v, w \in W$ and $c \in \mathbb{F}$.

Remark 4.7

- a. Note that a subspace must contain the zero vector.
- b. If W is a subspace of V , then W becomes an \mathbb{F} -vector space itself, by limiting the operations in V to W .
- c. A nonempty subset W is a subspace if and only if $av + bw \in W, v, w \in W, a, b, \in \mathbb{F}$.
- d. If $(W_i)_{i \in I}$ is a family of subspaces of V , then

$$W := \bigcap_{i \in I} W_i.$$

is also a subset of V .

4.2.2 Problems for Practice**Problem 4.2.1**

Show that none of the following sets of vectors is a subspaces pf \mathbb{R}^3 :

- a. The set U of vectors $x = (x_1, x_2, x_3)$ such that $\sum_{i=1}^3 x_i^2 = 1$.
- b. The set V of vectors in \mathbb{R}^3 all of whose coordinates are integers.
- c. Thet set W of vectors \mathbb{R}^3 that have at least one coordinate equal to 0.

Problem 4.2.2

Determine if U is a subspace of $M_2(\mathbb{R})$, where

- a. U is the set of 2×2 matrices such that the sum of the entries in the first column is 0.
- b. U is the set of 2×2 matrices such that the product of the product of the entries in the first column is 0.

Problem 4.2.6

Let V be the set of twice differerentiable functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that for all x we have

$$f''(x) + x^2 f'(x) - 3f(x) = 0.$$

Is V a subspace of the space of all maps $f : \mathbb{R} \rightarrow \mathbb{R}$?

Problem 4.2.7

Let V be the set of differentiable functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x$ we have

$$f'(x) - f(x)^2 = x$$

Is V a subspace of the space of all maps $f : \mathbb{R} \rightarrow \mathbb{R}$?

Problem 4.2.8

- Is the set of bounded sequences of real numbers a vector space of the space of all sequences of real numbers?
- Answer the question if instead of bounded sequences we consider monotonic sequences.

Problem 4.2.9

Let V be the set of all sequences $(x_n)_{n \geq 0}$ of real numbers such that

$$x_{n+2} + nx_{n+1} - (n-1)x_n = 0 \quad n \geq 0.$$

Prove that V is a subspace of the space of all sequence of real numbers.

Problem 4.2.10

Let V be the space of real-valued maps on \mathbb{R} and let W be the \subset of V consisting of maps f such that $f(0) + f(1) = 0$.

- Check that W is a subspace of V .
- Find a subspace S of V such that $V = W \oplus S$.

Problem 4.2.12

Let V be the space of convergent sequences of real numbers. Let W be the \subset of V consisting of sequences converging to 0 and let Z be the \subset of V consisting of constant sequences. Prove or disprove that W, Z are subspaces of V and $W \oplus Z = V$.

Problem 4.2.13 (Quotient space)

Let V be a vector space over \mathbb{F} and let $W \subset V$ be a subspace. For a vector $v \in V$, let $[v] = \{v + w : w \in W\}$. Note that $[v_1] = [v_2]$ if $v_1 - v_2 \in W$. Define the *quotient space* V/W to be $\{[v] : v \in V\}$. Define an addition and scalar multiplication on V/W by $[u] + [v] = [u + v]$ and $a[v] = [av]$. Prove that the addition and multiplication above are well-defined and V/W equipped with these operations is vector space.

4.3 Linear Combinations and Span**4.3.1 Summary****4.3.2 Problems for Practice****Problem 4.3.1**

Show that the vector $(1 \ 1 \ 1)$ can't be expressed as a linear combination of

$$a_1 = (1 \ -1 \ 0), \quad a_2 = (1 \ 0 \ -1) \quad \text{and} \quad a_3 = (0 \ 1 \ -1).$$

Problem 4.3.3

Let W be the \subset of \mathbb{R}^n consisting of vectors whose sum of coordinates equals 0. Let $Z = \text{Span}(1, \dots, 1)$ in \mathbb{R}^n . Prove or disprove that $W \oplus Z = \mathbb{R}^n$.

Problem 4.3.4

Let $P = \text{Span}[(1 \ 1 \ 1), (1 \ 1 \ -1)]$ in \mathbb{R}^3 , and let $D = \text{Span}(0, 1, -1)$. Is it true that $P \oplus D = \mathbb{R}^3$?

Problem 4.3.6

Let V be the vector space of real-valued on \mathbb{R} and let f_n (respectively g_n) be the map $x \mapsto nx$ (respectively $\cos^n x$). Prove or disprove that

$$\text{Span}(\{f_n \text{ bar } n \geq 0\}) = \text{Span}(\{g_n \text{ bar } n \geq 0\}).$$

4.4 Linear Independence**4.4.1 Summary**

Defintion 4.4.1 (Linear (in)dependence)

a. Vectors v_1, \dots, v_n in some vector space are *linearly dependent* if

$$\sum_{i=1}^n c_i v_i = 0$$

forces at least one of c_1, \dots, c_n to be nonzero.

b. Vectors v_1, \dots, v_n in some vector space v are *linearly independent* if

$$\sum_{i=1}^n c_i v_i = 0$$

forces $c_1 = \dots = c_n = 0$.

Remark 4.4.2

- a. A subfamily of a linearly independent family is linearly independent.
- b. If two vectors in a family of vectors are equal, then this family is automatically linearly dependent.

Remark 4.4.3

Given some vectors

4.4.2 Problems for Practice**Problem 4.4.1**

Are the vectors

$$v_1 = (1 \ 2 \ 1), \quad v_2 = (-3 \ 4 \ 5), \quad v_3 = (0 \ 2 \ -3)$$

linearly independent in \mathbb{R}^3 ?

Problem 4.4.2

Consider the vectors

$$v_1 = (1 \ 2 \ 1 \ 3), \quad v_2 = (1 \ -1 \ 1 \ -1), \quad v_3 = (3 \ 0 \ 3 \ 1)$$

.

- a. Prove that v_1, v_2, v_3 are linearly dependent.
- b. Express one of these vectors as a linear combination of two other vectors.

Problem 4.4.3

Let V be the vector space of polynomials with real coefficients whose degree does not exceed 3. Are the following vectors

$$1 + 3X + x^2, \quad X^3 - 3X + 1, \quad 3X^3 - X^2 - X - 1$$

linearly independent in V ?

Problem 4.4.4

Let V be the space of all real-valued on \mathbb{R} .

a. If $a_1 < \dots < a_n \in \mathbb{R}$, compute

$$\lim_{x \rightarrow \infty} \sum_{i=1}^n e^{(a_i - a_n)x}.$$

b. Prove that the family of maps $(x \mapsto e^{ax})_{a \in \mathbb{R}}$ is linearly independent in V .

Problem 4.4.5

Let V be the space of all maps $\varphi : [0, \infty) \rightarrow \mathbb{R}$. For each $a \in (0, \infty)$ consider the map $f_a \in V$ defined by

$$f_a(x) = \frac{1}{x + a}.$$

a. Let $a_1 < \dots < a_n$ be positive real numbers and suppose that $\alpha_1, \dots, \alpha_n$ are real numbers such that

$$\sum_{i=1}^n \alpha_i f_{a_i}(x) = 0 \quad \forall x \geq 0.$$

Prove that $\forall x \in \mathbb{R}$, we have

$$\sum_{i=1}^n \alpha_i \cdot \prod_{j \neq i} (x + a_j) = 0.$$

By making suitable choices of x , deduce that $\alpha_1 = \dots = \alpha_n = 0$.

b. Prove that the family that $(f_a)_{a>0}$ is linearly independent in V .

Problem 4.4.6

Consider the $V = \mathbb{R}$, seen as a vector space over $F = \mathbb{Q}$.

a. Prove the $1, \sqrt{2}, \sqrt{3}$ is a linearly independent set in V .

b. Prove that the set of numbers $\ln p$, where p runs over the prime numbers, is linearly independent in V .

Problem 4.4.7

a. If $m, n \neq 0 \in \mathbb{Z}$, compute

$$\int_0^{2\pi} \cos(mx) \cos(nx) dx.$$

b. Deduce that the maps $x \mapsto \cos nx$ with $n \neq 0 \in \mathbb{Z}$, form a linearly independent set in the space of all real-valued maps on \mathbb{R} .

c. Let v_1, \dots, v_n be linearly independent vectors in \mathbb{R}^n . Is it always the case that $v_1, v_1 + v_2, \dots, v_1 + v_2 + \dots + v_n$ are linearly independent?

4.5 Dimension Theory**Problem 1**

Given an irrep V of a finite group G , it is known that $\dim(V) \mid |G|$.

Using this, if $Z(G)$ denotes the center of the group, prove that $\dim(V) \mid |G/Z(G)|$.

5 Linear Transformations**5.1 Definitions and Objects Canonically Attached to a Linear Map****5.1.1 Summary****Defintion 5.1**

Let V, W be vector spaces over F . A *linear map* (or *linear transformation* or *homomorphism*) between V and W is a map $T : V \longrightarrow W$ satisfying the following two properties:

- 1) $T(v_1 + v_2) = T(v_1) + T(v_2)$ for vectors $v_1, v_2 \in V$;
- 2) $T(cv) = cT(v) \forall v \in V, c \in F$.

In practice, instead of checking separately that T respects addition and scalar multiplication, it may be advantageous to prove directly that

$$T(v_1 + cv_2) = T(v_1) + cT(v_2) \forall v_1, v_2 \in V, c \in \mathbb{F}.$$

Worked Example 5.2

If $T : V \longrightarrow W$ is a linear transformation, then $T(0) = 0 \wedge T(-v) = -T(v) \forall v \in V$.

Defintion 5.4

Let $T : V \rightarrow V$ be a linear map on a vector space V . A subspace W of V is called *stable under T* or *T -stable* if $T(W) \subset W$.

Worked Example 5.5

Consider the map $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ sending $(x_1, x_2) \rightarrow (x_2, -x_1)$. Find all subspaces of \mathbb{R}^2 which are stable under T .

Solution. Let W be a subspace of \mathbb{R}^2 that's T -stable. Obviously \mathbb{R}^2 and $\{0\}$ are stable under W , so let's assume W is neither of those spaces. Then $\dim W$ is necessarily 1, that is, $W = \mathbb{R}v$ for $v = (x_1, x_2)$. Since W is T -stable, this means that $T(v) = c(v)$ for some $c \in \mathbb{R} \implies (cx_1, cx_2) = (x_2, -x_1) \implies x_1 = -cx_1^2 \iff x_1 = x_2 = 0 (c \in \mathbb{R}) \iff v = 0$, an obvious contradiction. Thus the only subspaces stable under T are \mathbb{R}^2 and $\{0\}$. \square

Worked Example 5.7

Let V be a vector space over some field F and let $T : V \rightarrow V$ be a linear transformation. Suppose that all lines in V are stable subspaces under T . Prove that there is a scalar $c \in F$ such that $T(x) = cx \forall x \in V$.

Proposition 5.8

Let V, W be vector spaces. The set $\text{Hom}(V, W)$ of linear transformations between V and W is a subspace of $M(V, W)$.

Defintion 5.9

The *kernel* (or *null space*) of a linear transformation $T : V \rightarrow W$ is

$$\ker T = \{v \in V, T(v) = 0\}.$$

The *image* (or *range*) $\text{Im}(T)$ of T is the set

$$\text{Im}(F) = \{T(v) \mid v \in V\} \subset W.$$




Figure 1: Image and kernel

Proposition 5.10

If $T : V \longrightarrow W$ is a linear transformation, then T is injective if and only if $\ker T = \{0\}$.

Proposition 5.14

If $T : V \longrightarrow W$ is a linear transformation, then $\ker T$ and $\text{Im}(T)$ are subspaces of V , respectively W . Moreover, $\ker T$ is stable under T , and if $V = W$ then $\text{Im}(T)$ is stable under T .

5.1.2 Problems for Practice

In the next problems \mathbb{F} is a field.

Problem 5.1.2

Consider the map $f : \mathbb{R}^4 \longrightarrow \mathbb{R}^3$ defined by

$$f(x_1, x_2, x_3, x_4) = (x_1 + x_2 + x_3 + x_4; 2x_1 + x_2 - x_3 + x_4; x_1 - x_2 + x_3 - x_4).$$

- Prove that f is a linear map.
- Give a basis for $\ker F$.

Solution.

- f is a linear map because $\forall X = (x_1, \dots, x_4), Y = (y_1, \dots, y_4)$, we have

$$\begin{aligned} f(X + Y) &= (x_1 + y_1 + x_2 + y_2 + x_3 + y_3 + x_4 + y_4, \\ &\quad 2x_1 + 2y_1 + x_2 + y_2 - x_3 - y_3 + x_4 + y_4, \\ &\quad x_1 + y_1 - x_2 - y_2 + x_3 + y_3 - x_4 - y_4) \\ &= f(X) + f(Y) \end{aligned}$$

and $T(cX) = cT(X) \forall c \in \mathbb{R}$.

- $\ker f$ satisfies

$$f(X) = 0 \forall X \in \mathbb{R}^4.$$

Writing down the coordinates of $f(X)$ in matrix form gives us

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 1 & -1 & 1 \\ 1 & -1 & 1 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \implies \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = a \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}.$$

So a basis for $\ker f$ is $(0, 1, 0, -1)$. □

Problem 5.1.3

Let V be the space of polynomials with real coefficients whose degree does not exceed 3, and let the map $f : V \rightarrow \mathbb{R}^4$ be defined by

$$f(P) = [P(0), P(1), P(-1), P(2)].$$

- Prove that f is a linear map.
- Is f injective?

Solution.

- $\forall P, Q \in \mathbb{R}_3[X]$, we have $f(P + Q) = f(P) + f(Q)$ and $f(cP) = cf(P)$ for some $c \in \mathbb{R}$ and f is a linear map by definition.
- Let $P(x) = ax^3 + bx^2 + cx + d$. $\ker f$ satisfies

$$f(P) = (0, 0, 0, 0)$$

$$\Rightarrow a = b = c = 0 \Rightarrow \ker f = \{0\}. \text{ Thus } f \text{ is injective.}$$

□

Problem 5.1.4

Let $n \in \mathbb{N}^*$ and let V be the space of real polynomials whose degrees is not greater than n . Consider the map

$$f : V \rightarrow V, \quad f[P(X)] = P(X) + (1 - X)P'(X),$$

where $P'(X)$ is the derivative of P .

- Explain why f is a well-defined linear map.
- Give a basis for $\ker f$.

Problem 5.1.5

Find all subspaces of \mathbb{R}^2 which are stable under the linear transformation

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad T(x, y) = (x + y, -x + 2y).$$

Problem 5.1.8

Let V be a vector space over F and let $T_1, \dots, T_n : V \rightarrow V$ be linear transformation. Prove that

$$\bigcap_{i=1}^n \ker T_i \subseteq \ker \left(\sum_{i=1}^n T_i \right).$$

Problem 5.1.11

For each of the following maps $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, check that T is linear and then check whether $\ker T$ and $\text{Im}(T)$ are in direct sum.

- $T(x, y, z) = (x - 2y + z, x - z, x - 2y + z)$;
- $T(x, y, z) = (3x + 3y + 3z, 0, x + y + z)$.

Problem 5.1.12

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a map such that $f(x + y) = f(x) + f(y) \forall x, y \in \mathbb{R}$. Prove that f is a linear map of \mathbb{Q} -vector spaces between \mathbb{R} and itself.

Problem 5.1.13 (Quotient space)

Let V be a finite-dimensional vector space over F and let $W \subset V$ be a subspace. For a vector, let

$$[v] = \{v + w : w \in W\}.$$

Note that $[v_1] = [v_2]$ if $v_1 - v_2 \in W$. Define the quotient space V/W to be $\{[v] : v \in V\}$. Define an addition and scalar multiplication on V/W by $[u] + [v] = [u + v]$ and $a[v] = [av]$. We recall that the addition and multiplication above are well defined and V/W equipped w/ these operations is a vector space.

- Show that the map $\pi : V \rightarrow V/W$ defined by $\pi(v) = [v]$ is linear with kernel W .
- Show that $\dim W + \dim(V/W) = \dim V$.
- Show that $U \subset V$ is any subspace with $W \oplus U = V$. Show that $\pi|_U : U \rightarrow V/W$ is an isomorphism.
- Let $T : V \rightarrow U$ be a linear map, let $W \subset \ker T$ be a subspace, and $\pi : V \rightarrow V/W$ be the projection onto the quotient space. Show that there is a unique linear map $S : V/W \rightarrow U$ such that $T = S \circ \pi$.