

GCE January, 2019

Jiamin JIAN

Exercise 1:

Let $E := [0, 1] - S_{\mathbb{Q}} = [0, 1] \cap (S_{\mathbb{Q}})^c$ where $S_{\mathbb{Q}} := \{x \in [0, 1] | x = \frac{\sqrt{p}}{q} \text{ for some } p, q \in \mathbb{Z}^+\}$. Prove or disprove: There exists a closed, uncountable subset $F \subset E$.

Solution:

This proposition is true. Since $S_{\mathbb{Q}}$ is a countable set, there exists a bijection between $S_{\mathbb{Q}}$ and the positive rational number in the interval $[0, 1]$, so we can enumerate the set $S_{\mathbb{Q}}$ as $\{a_n | n \in \mathbb{N}\}$. That is to say we have $S_{\mathbb{Q}} = \{a_n | n \in \mathbb{N}\}$. And then we consider the union

$$\bigcup_{n=1}^{+\infty} (a_n - \frac{\epsilon}{2^n}, a_n + \frac{\epsilon}{2^n}),$$

it is an open set, we denote it as A , then $A = \bigcup_{n=1}^{+\infty} (a_n - \frac{\epsilon}{2^n}, a_n + \frac{\epsilon}{2^n})$. And when $\epsilon \rightarrow 0$, we know that $A \subset [0, 1]$ and $S_{\mathbb{Q}} \subset A$.

Since A is an open set, then $[0, 1] \cap (A)^c$ is a closed set. We denote $F = [0, 1] \cap (A)^c$, since the measure of set A is

$$m(A) = 2 \sum_{n=1}^{+\infty} \frac{\epsilon}{2^n} = 2\epsilon,$$

then we have $m(F) = 1 - 2\epsilon > 0$, so, the set F is uncountable. Since $F \subset E$ and it is both closed and uncountable, then the proposition is true.

For any countable set S , $S \subset [0, 1]$, let $E = [0, 1] - S$, we can find a closed, uncountable subset $F \subset E$, and we have the supremum of the measure of F is 1.

Exercise 2:

For x in $[-1, 1]$ set $P_n(x) = c_n(1 - x^2)^n$ where c_n is such that $\int_{-1}^1 P_n = 1$.

(i) Show that there is a positive constant C such that $c_n \leq C\sqrt{n}$.

(ii) Let f be a real valued continuous function on $[0, 1]$ such that $f(0) = f(1) = 0$.

Set for x in $[0, 1]$

$$f_n(x) = \int_0^1 P_n(x-t)f(t) dt$$

Show that f_n is uniformly convergence to f .

(iii) Let g be in $L^1((0, 1))$. Defining $g_n(x) = \int_0^1 P_n(x-t)g(t) dt$, is g_n uniformly convergence to g in $(0, 1)$? Does g_n converge to g in $L^1((0, 1))$?

Solution:

(i) Method 1:

Since $\int_{-1}^1 c_n(1-x^2)^n dx = 1$, then we have

$$c_n = \frac{1}{2 \int_0^1 (1-x^2)^n dx}.$$

Next we need to find a lower bound of the integral term $\int_0^1 (1-x^2)^n dx$. Since for $n > 1$,

$$\begin{aligned} \int_0^1 (1-x^2)^n dx &\geq \int_0^{\frac{1}{\sqrt{n}}} (1-x^2)^n dx \\ &\geq \frac{1}{\sqrt{n}} \left(1 - \frac{1}{n}\right)^n, \end{aligned}$$

then we have $c_n \leq \frac{\sqrt{n}}{2(1-\frac{1}{n})^n}$. We just need to find a lower bound of $(1 - \frac{1}{n})^n$. Since $(1 - \frac{1}{n})^n = 1 - C_{n-1}^1 \frac{1}{n} + C_{n-2}^2 \frac{1}{n^2} - \dots + (-1)^{n-1} \frac{1}{n^{n-1}} > \frac{1}{3} - \frac{2}{6n^2} > \frac{1}{4}$ as $n > 1$, then we set $C = 2$, we have $c_n \leq C\sqrt{n}$ for $n > 1$. For $n = 1$, we get $c_1 = \frac{3}{4} < 2$, then when $C = 2$, we have $c_n \leq C\sqrt{n}$ holds.

Method 2:

We change the element and define $x = \sin y$, then we have $\int_0^{\frac{\pi}{2}} c_n \cos^{2n+1} y dy = \frac{1}{2}$.

Since

$$\int_0^{\frac{\pi}{2}} \cos^{2n+1} y dy = 2n \int_0^{\frac{\pi}{2}} \cos^{2n-1} y dy - 2n \int_0^{\frac{\pi}{2}} \cos^{2n+1} y dy,$$

we denote $I_{2n+1} = \int_0^{\frac{\pi}{2}} \cos^{2n+1} y dy$, then we have $(2n+1)I_{2n+1} = 2nI_{2n-1}$. Since $I_1 = \int_0^{\frac{\pi}{2}} \cos y dy = 1$, we have $\int_0^{\frac{\pi}{2}} \cos^{2n+1} y dy = \frac{(2n)!!}{(2n+1)!!}$. And since

$$\begin{aligned} \frac{(2n)!!}{(2n+1)!!} &= \frac{2n(2n-2)\cdots 2}{(2n+1)(2n-1)\cdots 3} \\ &\geq \frac{\sqrt{2n+1}\sqrt{2n-1}\sqrt{2n-3}\cdots \sqrt{3}\sqrt{1}}{(2n+1)(2n-1)\cdots 3} \\ &= \frac{1}{\sqrt{2n+1}}, \end{aligned}$$

then we have $c_n \leq \frac{\sqrt{2n+1}}{2}$. We set $C = 1$, then we have $c_n \leq C\sqrt{n}$.

(ii) Firstly we extend $f(x)$ to a function from \mathbb{R} to \mathbb{R} by zero. Then we have

$$f_n(x) = \int_0^1 P_n(x-t)f(t) dt = \int_{\mathbb{R}} P_n(x-t)f(t) dt,$$

then we change the element as $x-t=y$, we have

$$f_n(x) = \int_{\mathbb{R}} P_n(y)f(x-y) dy.$$

Then we know that

$$\begin{aligned}
|f_n(x) - f(x)| &= \left| \int_{\mathbb{R}} P_n(y) f(x-y) dy - \int_{-1}^1 P_n(y) f(x) dy \right| \\
&= \left| \int_{-1}^1 P_n(y) (f(x-y) - f(x)) dy + \int_{([-1,1])^c} P_n(y) f(x-y) dy \right| \\
&\leq \int_{-1}^1 P_n(y) |(f(x-y) - f(x))| dy + \int_{([-1,1])^c} |P_n(y) f(x-y)| dy.
\end{aligned}$$

Since when $x \in [0, 1]$ and $y \in ([-1, 1])^c$, we have $x - y > 1$ or $x - y < 0$, then we have $f(x - y) = 0$, so we have

$$|f_n(x) - f(x)| \leq \int_{-1}^1 P_n(y) |(f(x-y) - f(x))| dy.$$

And by the definition of continuous, we have $\forall \epsilon > 0$, there $\exists \delta$, when $|x - y - x| < \delta$, we have $|f(x - y) - f(x)| < \epsilon$. We denote $S = [-1, 1] \cap [-\delta, \delta]$, since $f(x)$ is continuous in \mathbb{R} , we denote $\sup_{x \in [0, 1]} f(x) = M$, then we have $M < +\infty$ and

$$\begin{aligned}
|f_n(x) - f(x)| &\leq \int_{-\delta}^{\delta} P_n(y) |(f(x-y) - f(x))| dy + \int_S P_n(y) |(f(x-y) - f(x))| dy \\
&\leq \epsilon \int_{-\delta}^{\delta} P_n(y) dy + 2M \int_S P_n(y) dy \\
&\leq \epsilon + 2M \int_S c_n(1 - y^2)^n dy \\
&\leq \epsilon + 4MC\sqrt{n} \int_{\delta}^1 (1 - y^2)^n dy \\
&\leq \epsilon + 4MC\sqrt{n}(1 - \delta)(1 - \delta^2)^n.
\end{aligned}$$

Since $\lim_{n \rightarrow +\infty} 4MC\sqrt{n}(1 - \delta)(1 - \delta^2)^n = 0$, then we can say that there exists a $N \in \mathbb{N}$, when $n > N$, we have $4MC\sqrt{n}(1 - \delta)(1 - \delta^2)^n < \epsilon$. Overall, we know that $\forall x \in [0, 1], \forall \epsilon > 0$, there exists a $N \in \mathbb{N}$, when $n > N$, we have $|f_n(x) - f(x)| < 2\epsilon$, so that f_n is uniformly converges to f .

(iii) Firstly, the $g_n(x)$ is not uniformly convergent to g in $(0, 1)$, we can give an counter example as following. We define

$$g(x) = \begin{cases} 1, & x = \frac{1}{2} \\ 0, & x \in (0, \frac{1}{2}) \cup (\frac{1}{2}, 1), \end{cases}$$

obviously $g(x)$ is not continuous in $(0, 1)$, but we have $g_n(x) = \int_0^1 P_n(x-t)g(t) dt = 0, \forall x \in (0, 1)$. Then $g_n(x)$ is continuous in $[0, 1]$. Since $g(x)$ is not continuous in $(0, 1)$, we can say that $g_n(x)$ is not uniformly convergent to $g(x)$ in $(0, 1)$.

Secondly, we can show that $g_n(x)$ convergent to $g(x)$ in $L^1((0, 1))$. Since the continuous functions with compact support are dense in L^1 space, then for all $\epsilon > 0$, there exist

a continuous function $f(x) \in C_c([0, 1])$, such that $\|f - g\|_1 < \epsilon$. We define the $f_n(x)$ as the section (ii), then we have

$$\|g - g_n\|_1 \leq \|g - f\|_1 + \|f - f_n\|_1 + \|f_n - g_n\|_1.$$

Since f_n is uniformly converges to f , for all $\epsilon > 0$, there exists a $N \in \mathbb{N}$, when $n > N$, we have $\|f - f_n\|_1 < \epsilon$. And for the same ϵ , by the property that continuous function is dense in L^1 space, we have $\|f - g\|_1 < \epsilon$. Next we verify that $\|f_n - g_n\|_1 < \epsilon$. Since

$$\begin{aligned} \|f_n - g_n\|_1 &= \int_0^1 \left| \int_0^1 P_n(x-t)g(t) - \int_0^1 P_n(x-t)f(t) dt \right| dx \\ &= \int_0^1 \left| \int_0^1 P_n(x-t)(g(t) - f(t)) dt \right| dx \\ &\leq \int_0^1 \int_0^1 P_n(x-t)|g(t) - f(t)| dt dx, \end{aligned}$$

and $P_n(x-t)$ is continuous for $t \in [0, 1]$, then we can find the upper bound for $P_n(x-t)$, we denote it as C , then we have

$$\begin{aligned} \|f_n - g_n\|_1 &\leq \int_0^1 \int_0^1 P_n(x-t)|g(t) - f(t)| dt dx \\ &\leq C \int_0^1 \int_0^1 |g(t) - f(t)| dt dx \\ &= C \int_0^1 |g(t) - f(t)| dt \\ &= C\|g - f\|_1. \end{aligned}$$

Since $\|g - f\|_1 < \epsilon$, we have $\|g - g_n\|_1 < (2 + \frac{1}{C})\epsilon$ for all $\epsilon > 0$. So, we know that $g_n(x)$ convergent to $g(x)$ in $L^1((0, 1))$.

Exercise 3:

Give an example of $f_n, f : \mathbb{R} \mapsto [0, \infty)$ such that $f_n \in L^1(\mathbb{R})$ for every $n \in \mathbb{N}$, $f \in L^2(\mathbb{R})$, $f_n \leq f$ for every $n \in \mathbb{N}$, $f_n \rightarrow 0$ a.e., and $\int f_n \nrightarrow 0$.

Solution:

We define the $f(x) = \frac{1}{x}\mathbb{I}_{[1, +\infty)}$ and $f_n(x) = \frac{1}{x}\mathbb{I}_{[n, n^2]}$. For a fixed n , we have

$$\int_{\mathbb{R}} |f_n(x)| dx = \int_n^{n^2} \frac{1}{x} dx = \ln n,$$

so we have $f_n \in L^1(\mathbb{R})$ for every $n \in \mathbb{N}$. And since

$$\int_{\mathbb{R}} |f(x)|^2 dx = \int_1^{+\infty} \frac{1}{x^2} dx = 1,$$

so we know that $f \in L^2(\mathbb{R})$. Since for all n , f_n is just a part of f and $f > 0$, then we have $f_n \leq f$ for every $n \in \mathbb{N}$. When $n \rightarrow +\infty$, we have $f_n(x) \leq \frac{1}{n}$, so that $f_n \rightarrow 0$ almost everywhere. And we calculate the integral of f_n , we have

$$\int_{\mathbb{R}} f_n(x) dx = \int_n^{n^2} \frac{1}{x} dx = \ln n,$$

when $n \rightarrow +\infty$, $\ln n \rightarrow +\infty$, so we can get $\int f_n \nrightarrow 0$.