

A Mean Field Game - Hybrid Systems Approach to Optimal Execution Problems in Finance with Stopping Times

Dena Firoozi, *Member, IEEE*, Ali Pakniyat, *Member, IEEE* and Peter E. Caines, *Life Fellow, IEEE*.

Abstract—Following standard financial models, the stock market is studied in this paper as a large population non-cooperative game where each trader has stochastic linear dynamics with quadratic costs. We consider the case where there exists one major trader with significant influence on market movements together with a large number of minor traders, each with individually asymptotically negligible effect on the market. The traders are coupled in their dynamics and cost functions by the market's average trading rate (a component of the system mean field). In this work, optimal stopping time strategies together with best response trading policies for all agents are established employing a hybrid systems approach in an MFG framework where each agent attempts to (i) maximize its wealth, and avoid (ii) large execution prices and (iii) large trading accelerations, where (ii) and (iii) are associated with market instability.

I. INTRODUCTION

Mean Field Game (MFG) systems theory establishes the existence of approximate Nash equilibria together with the corresponding individual strategies for stochastic dynamical systems in games involving a large number of agents. The equilibria are termed ϵ -Nash equilibria and are generated by the local, limited information feedback control actions of each agent in the population, where the feedback control actions constitute the best response of each agent with respect to the precomputed behaviour of the mass of agents and where the approximation error converges to zero as the population size goes to infinity.

The analysis of this set of problems originated in [1]–[3] (see [4]), and independently in [5], [6]. In [7] and [8] the authors analyse and solve the linear quadratic systems case where there is a major agent (i.e. non-asymptotically vanishing as the population size goes to infinity) together with a population of minor agents (i.e. individually asymptotically negligible). The existence of ϵ -Nash equilibria is established together with the individual agents' control laws that yield the equilibria [8]. The partially observed MFG theory for nonlinear and linear quadratic systems with major and minor agents has been developed in [9]–[15].

The notion of Hybrid Systems (HS) is used to describe control systems for which (i) the state has continuous and discrete components and (ii) the continuous component of the state evolves in “continuous time” and the discrete component evolves in “discrete time”. While the optimal

control of deterministic hybrid systems has been extensively studied in the literature (see e.g. [16]–[27]), the optimal control of stochastic hybrid systems, i.e. control systems that involve the interaction of continuous dynamics, discrete dynamics and stochastic diffusions, has been the subject of a limited number of studies [28]–[31]. In [30], in particular, first order variational analysis is performed on the stochastic hybrid optimal control problem via the needle variation methodology and the necessary optimality conditions are established in the form of the Stochastic Hybrid Minimum Principle (SHMP). The results are specialized in [31] to a class of Linear Quadratic Gaussian Hybrid Optimal Control Problems (LQG-HOCP) for which the Hamiltonian boundary conditions are path-independent and therefore, the corresponding stochastic Riccati equations are independent from the realization of stochastic diffusion terms.

Optimal execution problems have been addressed in the literature (see e.g. [32]–[36]) in which an agent must liquidate or acquire a certain amount of shares over a pre-specified time horizon while it balances the price impact and the price uncertainty, and maximizes its final wealth. This problem with the linear models in [32] was formulated as for the nonlinear major minor (MM) MFG model in [37], and partially observed MM LQG MFG theory was first applied in [38], [39] for the case in which traders have partial observations of the market states. In this work, the institutional investor, interpreted as a major agent, aims to liquidate a specific amount of shares. Furthermore, there is a large population high frequency traders (HFTs), interpreted as minor agents, who wish to liquidate or acquire a certain amount of shares within a specific time horizon. Optimal stopping time strategies together with best response trading policies for all agents is established employing a hybrid systems approach in an MFG framework where each agent attempts to (i) maximize its wealth, and avoid (ii) large execution prices and (iii) large trading accelerations, where (ii) and (iii) are associated with market instability, and are appropriately weighted in the agent's performance function.

We note major trader (respectively, minor trader), and institutional trader (respectively, HFT) are used interchangeably in this paper.

II. TRADING DYNAMICS OF AGENTS IN THE MARKET

As stated in the Introduction, the institutional investor is considered as a major agent in the mean field model of the market which liquidates its shares and the HFTs are considered as minor agents, where two types of them are considered: acquirers \mathcal{A}_a with the population of N_a

This work was supported by grants from NSERC, Canada.

D. Firoozi, A. Pakniyat and P. E. Caines are with the Centre for Intelligent Machines (CIM) and the Department of Electrical and Computer Engineering (ECE), McGill University, Montreal, QC, Canada.

dena.firoozi@mail.mcgill.ca,

ali.pakniyat@mail.mcgill.ca, peterc@cim.mcgill.ca

and liquidators \mathcal{A}_l with the population of N_l , such that $N_a + N_l = N$. All agents trade over the interval $[0, T]$, and minor agents are allowed to stop trading at an optimal time $t_s^i \leq T$. It will be shown in Section III that the optimal stopping time policy for each agent is \mathcal{F}_t -independent, and depends only on its dynamical parameters. In this paper, for simplicity of exposition the dynamical parameters for all minor traders in their respective type are the same, and hence the stopping times are the same for all agents of each population. Employing the trading model in [32], the trading dynamics of the major agent and any generic minor agent in the market are described by the linear time evolution of the (i) inventories, (ii) trading rates and (iii) prices while the bilinear cash process appears in the quadratic performance function for each agent.

A. Inventory Dynamics

It is assumed that the institutional investor liquidates its inventory of shares, $Q_0(t)$, by trading at a rate $\nu_0(t)$ during the trading period $[0, T]$. Hence the major agent's inventory dynamics is given by

$$dQ_0(t) = \nu_0(t)dt + \sigma_0^Q dw_0^Q, \quad 0 \leq t \leq T,$$

where w_0^Q is a Wiener process modeling the noise in the inventory information that the institutional trader collects from its branches in different locations; σ_0^Q is a positive scalar and we assume that $Q_0(0) \gg 1$. The same dynamical model is adopted for the trading dynamics of a generic HFT

$$dQ_i(t) = \nu_i(t)dt + \sigma_i^Q dw_i^Q,$$

where for a minor acquirer trader $\mathcal{A}_i \in \mathcal{A}_a$, $0 \leq t \leq t_s^a$, and correspondingly for a minor liquidator $\mathcal{A}_i \in \mathcal{A}_l$, $0 \leq t \leq t_s^l$. The Wiener process w_i^Q models the HFT's information noise, σ_i^Q is a positive scalar, $\nu_i(t)$ is the agent's rate of trading which can be positive or negative depending on whether the agent is acquirer or liquidator, respectively; $Q_i(t)$ is the minor liquidator's remaining shares at time t , or the shares the minor acquirer has bought until time t . However, the initial share stock of the HFTs, $\{Q_i(0), 1 \leq i \leq N_a + N_l\}$, are not considered to be large, furthermore they are not motivated to retain shares and are assumed to trade them quickly.

We assume that the trading rate of the major agent is controlled via $u_0(t)$ as

$$d\nu_0(t) = u_0(t)dt, \quad 0 \leq t \leq T,$$

where the trading strategy $u_0(t)$ can be seen to be the trading acceleration of the major trader. Correspondingly, $u_i(t)$ controls the trading rate of minor agent, \mathcal{A}_i , by

$$d\nu_i(t) = u_i(t)dt,$$

where again for a minor acquirer trader $\mathcal{A}_i \in \mathcal{A}_a$, $0 \leq t \leq t_s^a$, and correspondingly for a minor liquidator $\mathcal{A}_i \in \mathcal{A}_l$, $0 \leq t \leq t_s^l$, and $u_i(t)$ is the trading acceleration of the minor acquirer or liquidator.

B. Price Dynamics

The trading rate of the major agent and the average trading rate of the minor agents give rise to the fundamental asset price which models the permanent effect of agents' trading rates on the market price. Further, each agent has a temporary effect on the asset price which only persists during the action of the trade and which determines the execution price, that is to say the price at which each agent can trade.

1) *Fundamental Asset Price:* We model the dynamics of the fundamental asset price, as seen from the major agent's viewpoint, by

$$dF_0(t) = (\lambda_0 \nu_0(t) + \lambda \nu^{N_t}(t))dt + \sigma dw_0^F(t), \quad 0 \leq t \leq T,$$

where N_t is the number of minor agents trading at time t , $\nu^{N_t}(t) = \frac{1}{N_t} \sum_{i=1}^{N_t} \nu_i(t)$ is the average trading rate of the minor agents trading at time t . The Wiener process $w_0^F(t)$ models the aggregate effect of all traders in the market which - unlike the major and minor agents $\mathcal{A}_0, \mathcal{A}_i$, - have no complete or partial observations on any of the state variables appearing in the dynamical market model (these are termed uninformed traders). Further, σ denotes the intensity of the market volatility and $\lambda_0, \lambda \geq 0$ denote the strength of the linear permanent impact of the major and minor agents' trading on the fundamental asset price, respectively. Similarly, we model the fundamental asset price dynamics, as seen by a minor agent \mathcal{A}_i , by

$$dF_i(t) = (\lambda_0 \nu_0(t) + \lambda \nu^{N_t}(t))dt + \sigma dw_i^F(t),$$

where $0 \leq t \leq t_s^a$, for $\mathcal{A}_i \in \mathcal{A}_a$, and $0 \leq t \leq t_s^l$, for $\mathcal{A}_i \in \mathcal{A}_l$, $\nu^{N_t}(t) = \frac{1}{N_t} \sum_{i=1}^{N_t} \nu_i(t)$ is again the average trading rate of the N_t minor agents trading at t , and the Wiener process, $w_i^F(t)$, represents the mass effect of all uninformed traders in the market. The time differences between agents in getting data from fast changing limit order book make the Wiener processes, w_i^F , $0 \leq i \leq N_a + N_l$ independent.

2) *Execution Price:* The major agent's execution price $S_0(t)$ evolution is assumed to be given by

$$dS_0(t) = dF_0(t) + a_0 d\nu_0(t), \quad 0 \leq t \leq T, \quad (1)$$

where $a_0 \geq 0$ is the temporary impact strength of the major agent on fundamental asset price. Likewise, a minor agent's execution price, $S_i(t)$, is assumed to evolve by

$$dS_i(t) = dF_i(t) + a d\nu_i(t), \quad (2)$$

where $0 \leq t \leq t_s^a$, for $\mathcal{A}_i \in \mathcal{A}_a$, and $0 \leq t \leq t_s^l$, for $\mathcal{A}_i \in \mathcal{A}_l$, and a models the temporary impact of a minor agent's trading on its execution price.

C. Cash Process

The cash processes for the major agent and a generic minor agent, $Z_0(t)$, $Z_i(t)$, respectively, are given by

$$dZ_0(t) = -S_0(t)dQ_0(t), \quad 0 \leq t \leq T, \quad (3)$$

$$dZ_i(t) = -S_i(t)dQ_i(t), \quad \begin{array}{l} \text{for } \mathcal{A}_i \in \mathcal{A}_a, \quad 0 \leq t \leq t_s^a \\ \text{for } \mathcal{A}_i \in \mathcal{A}_l, \quad 0 \leq t \leq t_s^l \end{array}, \quad (4)$$

where $Z_0(t)$, and $Z_i(t)$ for $\mathcal{A}_i \in \mathcal{A}_l$ are the cash obtained through liquidation of shares, and $Z_i(t)$, for $\mathcal{A}_i \in \mathcal{A}_a$ is the cash paid for acquisition of shares up to time t . We note that the value of $dQ_0(t)$ in a stock sale (respectively, buy) is negative (respectively, positive) and hence for positive $S_0(t)$, $Z_0(t)$ increases (respectively, decreases).

D. Performance Function

1) *Major Liquidator*: The objective for the major trader is to liquidate \mathcal{N}_0 shares and maximize the cash it holds at the end of the trading horizon, i.e. maximize $Z_0(T)$, and if the remaining inventory at the final time T is $Q_0(T)$, it can liquidate it at a lower price than the market asset price reflected at cost function by $Q_0(T)(F_0(T) - \alpha Q_0(T))$. Further, the major trader's utility in minimizing the inventory over the period $[0, T]$ is modeled by including the penalty $\phi \int_0^T Q_0^2(s) ds$ in its objective function, and the utility of avoiding very high execution prices, large trading intensities and large trading accelerations by including the terms $\epsilon S_0^2(T)$, $\int_0^T \delta S_0^2(s) ds$, $\beta \nu_0^2(T)$, $\int_0^T \theta \nu_0^2(s) ds$ and $\int_0^T R_0 u_0^2(s) ds$ in the objective function. Therefore, its cost function to be minimized is given by

$$J_0(u_0, u_{-0}) = \mathbb{E} \left[-r Z_0(T) - p Q_0(T) (F_0(T) - \alpha Q_0(T)) + \epsilon S_0^2(T) + \beta \nu_0^2(T) + \int_0^T (\phi Q_0^2(s) + \delta S_0^2(s) + \theta \nu_0^2(s) + R_0 u_0^2(s)) ds \right], \quad (5)$$

where $r, p, \alpha, \epsilon, \beta, \phi, \delta, \theta$, and R_0 are positive scalars, and $u_{-0} := (u_1, u_2, \dots, u_N)$ are trading strategies of the minor traders. Note that for larger values of ϕ the trader attempts to liquidate its inventory more quickly.

2) *Minor Liquidator*: In a similar way, the objective function to be minimized for a liquidator HFT who wants to liquidate \mathcal{N}_l shares over the interval $[0, T]$ with the stopping time $0 \leq t_s^l \leq T$ is given by

$$J_l(u_l, u_{-l}) = \mathbb{E} \left[-r_l Z_l(t_s^l) - p_l Q_l(t_s^l) (F_l(t_s^l) - \psi_l Q_l(t_s^l)) + \xi_l S_l^2(t_s^l) + \mu_l \nu_l^2(t_s^l) + \int_0^{t_s^l} (\kappa_l Q_l^2(s) + \gamma_l S_l^2(s) + \varrho_l \nu_l^2(s) + R_l u_l^2(s)) ds \right], \text{ for } \mathcal{A}_l \in \mathcal{A}_l \quad (6)$$

where $r_l, p_l, \psi_l, \xi_l, \mu_l, \kappa_l, \gamma_l, \varrho_l$ and R_l are positive scalars, and $u_{-l} := (u_0, u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_N)$. Note that $\mathcal{N}_l \ll \mathcal{N}_0$.

3) *Minor Acquirer*: The objective for a minor acquirer is to buy \mathcal{N} shares during the trading horizon $[0, T]$. Given that it stops trading at $t_s^a \leq T$, it also wishes to minimize the execution cost including the cash $Z_i(t_s^a)$ paid up to time t_s^a , and the cash must be paid at time t_s^a to buy the remaining shares at once at a higher price than the market's asset price, i.e. $(\mathcal{N} - Q_i(t_s^a))(F_i(t_s^a) + \psi_a(\mathcal{N} - Q_i(t_s^a)))$. It is also intended to avoid high execution prices, large trading intensities and large trading accelerations modeled by including

$\xi_a S_i^2(t_s^a) + \mu_a \nu_i^2(t_s^a) + \int_0^{t_s^a} (\gamma_a S_i^2(s) + \varrho_a \nu_i^2(s) + R_a u_i^2(s)) ds$ in its objective function

$$J_i(u_i, u_{-i}) = \mathbb{E} \left[p_a (\mathcal{N} - Q_i(t_s^a)) (F_i(t_s^a) + \psi_a (\mathcal{N} - Q_i(t_s^a))) + r_a Z_i(t_s^a) + \xi_a S_i^2(t_s^a) + \mu_a \nu_i^2(t_s^a) + \int_0^{t_s^a} (\kappa_a (\mathcal{N} - Q_i(s))^2 + \gamma_a S_i^2(s) + \varrho_a \nu_i^2(s) + R_a u_i^2(s)) ds \right], \quad \mathcal{A}_i \in \mathcal{A}_a, \quad (7)$$

where $\int_0^{t_s^a} \kappa_a (\mathcal{N} - Q_i(s))^2 ds$ is to penalize the agent for the remaining shares to be bought up to t_s^a and to expedite the acquisition. The parameters $p_a, \psi_a, r_a, \xi_a, \mu_a, \kappa_a, \gamma_a, \varrho_a$, and R_a are positive scalars and $u_{-i} := (u_0, u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_N)$.

III. HS-MFG FORMULATION OF THE OPTIMAL EXECUTION PROBLEM

In this section we formulate the optimal execution problem in the MM LQG MFG framework.

A. Discrete state association

In order to present the trading dynamics of the stock market in the stochastic hybrid systems framework of [30], [31], the discrete states q_j , $j = 0, 1, 2$ are introduced, which correspond to the evolution of the market in the intervals $[t_j, t_{j+1})$, where $t_0 = 0$ is the initial time, t_1 and t_2 denote the stopping times of the first population and the second population respectively, and $t_3 = T$ is the terminal time.

We remark that the HS-MFG problems studied in this paper lie within the class of hybrid LQG problems in [31] for which optimal switching strategies are \mathcal{F}_t -independent, and therefore, optimal stopping strategies depend only on the dynamical parameters of each population.

We associate the discrete state q_0 to the initial case where both the liquidator and acquirer populations are trading together with the major agent over the interval $[0, t_1)$.

The discrete state q_1 corresponds to the interval $[t_1, t_2)$ for which two situations can be considered: (i) the liquidator population stops at t_1 while the acquirer population is still trading, in which case $q_1 = q_a$, and (ii) the acquirer population stops at t_1 while the liquidator population is trading, which corresponds to $q_1 = q_l$.

The discrete state q_2 represents the system over the interval $[t_2, T]$ after the second population of HFTs stops at t_2 , and hence the major agent is trading in the absence of both populations.

The above discrete state association is summarized in the following table.

Discrete State		\mathcal{A}_0	\mathcal{A}_a	\mathcal{A}_l
q_0		✓	✓	✓
q_1	q_a	✓	✓	×
	q_l	✓	×	✓
q_2		✓	×	×

B. Finite Populations

1) *Major Agent*: The dynamics of the major trader in the market can be modeled as

$$\begin{aligned} d\nu_0(t) &= u_0(t)dt, \\ dQ_0(t) &= \nu_0(t)dt + \sigma_0^Q dw_0^Q, \\ dS_0(t) &= (\lambda_0 \nu_0(t) + \lambda \nu^{N_t}(t))dt + a_0 u_0(t)dt + \sigma dw_0^F(t). \end{aligned}$$

Let the major agent's state be denoted by $x_0 = [\nu_0, Q_0, S_0]^T$, then its dynamics can be expressed as

$$dx_0 = A_0 x_0 dt + B_0 u_0 dt + E_0 x^{N_t} dt + D_0 dw_0 \quad (8)$$

with the matrices

$$\begin{aligned} A_0 &= \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ \lambda_0 & 0 & 0 \end{bmatrix}, \quad B_0 = \begin{bmatrix} 1 \\ 0 \\ a_0 \end{bmatrix}, \quad w_0 = \begin{bmatrix} w_0^Q \\ w_0^F \end{bmatrix} \\ E_0 &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \lambda & 0 & 0 \end{bmatrix}, \quad D_0 = \begin{bmatrix} 0 & 0 \\ \sigma_0^Q & 0 \\ 0 & \sigma \end{bmatrix}. \end{aligned}$$

Note that in (8), N_t takes the following values.

$$N_t = \begin{cases} N_a + N_l & \text{for } q_0, \\ N_a & \text{for } q_a, \\ N_l & \text{for } q_l, \\ 0 & \text{for } q_2. \end{cases} \quad (9)$$

The major trader's cost function (5) can also be described in terms of its states with replacing the final cash process by $\mathbb{E}[Z_0(T)] = -\mathbb{E}[\int_0^T S_0(s)\nu_0(s)ds]$, and the fundamental asset price $F_0(T)$ using (1). The equation (8) together with the cost function (5) form the stochastic LQG problem for the major trader. Note that the major trader is involved with the market's average trading rate in its dynamics while involved with the market's average selling rate in its cost function.

2) *Minor Liquidator*: Similarly, the stochastic optimal control problem for a minor liquidator $\mathcal{A}_i \in \mathcal{A}_l$, is given by the set of dynamical equations

$$\begin{aligned} d\nu_i(t) &= u_i(t)dt, \\ dQ_i(t) &= \nu_i(t)dt + \sigma_i^Q dw_i^Q, \\ dS_i(t) &= (\lambda_0 \nu_0(t) + \lambda \nu^{N_t}(t))dt + a u_i(t)dt + \sigma dw_i^F. \end{aligned}$$

Similar to the major trader, we define a generic minor trader's state vector as $x_i = [\nu_i, Q_i, S_i]^T$, and its dynamics can be written as

$$dx_i = A_l x_i dt + B_l u_i dt + E_l x^{N_t} dt + D_l dw_i \quad (10)$$

with

$$\begin{aligned} A_l &= \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad E_l = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \lambda & 0 & 0 \end{bmatrix}, \quad B_l = \begin{bmatrix} 1 \\ 0 \\ a \end{bmatrix} \\ G_l &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \lambda_0 & 0 & 0 \end{bmatrix}, \quad D_l = \begin{bmatrix} 0 & 0 \\ \sigma_i^Q & 0 \\ 0 & \sigma \end{bmatrix}, \quad w_{li} = \begin{bmatrix} w_{li}^Q \\ w_{li}^F \end{bmatrix}. \end{aligned}$$

The quadratic cost function (6) can also be expressed in terms of the minor agent's state when the final cash process in (6) is replaced by $\mathbb{E}[Z_i(t_s^l)] = -\mathbb{E}[\int_0^{t_s^l} S_i(s)\nu_i(s)ds]$ using (4), and the fundamental asset price $F_i(t_s^l)$ is replaced using (2). The equations (10) and (6) form the stochastic LQG problem for a generic minor liquidator. Additionally, they show that a minor liquidator is coupled with the major agent's trading rate and the market's average trading rate in its dynamics while coupled with the market's average selling rate in its cost function.

3) *Minor Acquirer Agent*: The stochastic optimal control problem for a minor acquirer $\mathcal{A}_i \in \mathcal{A}_a$, is given by the set of dynamical equations

$$\begin{aligned} d\nu_i(t) &= u_i(t)dt, \\ dY_i(t) &= -\nu_i(t)dt + \sigma_i^Q dw_i^Q, \\ dS_i(t) &= (\lambda_0 \nu_0(t) + \lambda \nu^{N_t}(t))dt + a u_i(t)dt + \sigma dw_i^F, \end{aligned}$$

where $Y_i(t) = N_a - Q_i(t)$ is the remaining shares at t to be acquired until the end of trading horizon. We define a generic minor acquirer's state vector as $x_i = [\nu_i, Y_i, S_i]$, hence its dynamics in compact form would be

$$dx_i = A_a x_i dt + B_a u_i dt + E_a x^{N_t} dt + D_a dw_{ai}, \quad (11)$$

where

$$\begin{aligned} A_a &= \begin{bmatrix} 0 & 0 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad E_a = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \lambda & 0 & 0 \end{bmatrix}, \quad B_a = \begin{bmatrix} 1 \\ 0 \\ a \end{bmatrix} \\ G_a &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \lambda_0 & 0 & 0 \end{bmatrix}, \quad D_a = \begin{bmatrix} 0 & 0 \\ \sigma_i^Q & 0 \\ 0 & \sigma \end{bmatrix}, \quad w_{ai} = \begin{bmatrix} w_{ai}^Q \\ w_{ai}^F \end{bmatrix}. \end{aligned}$$

Note that N_t in (11) again takes values as in (9) over the trading horizon. Accordingly, the cost function for acquisition is given by

$$\begin{aligned} J_i(u_i, u_{-i}) &= \mathbb{E} \left[p_a Y_i(t_s^a) (S_i(t_s^a) - a \nu_i(t_s^a) + \psi_a Y_i(t_s^a)) \right. \\ &\quad + \xi_a S_i^2(t_s^a) + \mu_a (\nu_i(t_s^a) - \rho_a \bar{\nu}(t_s^a))^2 + \\ &\quad \int_0^{t_s^a} (\kappa_a Y_i(s)^2 + \gamma_a S_i^2(s) + \varrho_a \nu_i^2(s) \\ &\quad \left. - r_a S_i(s) \nu_i(s) + R_a u_i^2(s)) ds \right], \text{ for } \mathcal{A}_i \in \mathcal{A}_a. \quad (12) \end{aligned}$$

The set of equations (11)-(12) constitute the standard stochastic LQG problem for a minor acquirer. It can be seen that a generic minor acquirer interacts with the major agent's trading rate as well as the market's average trading rate through its dynamics, and with the market's average buying rate through its cost function.

C. Mean Field Evolution

Following the LQG MFG methodology [7], the mean field, \bar{x} , is defined as the L^2 limit, when it exists, of the average of minor agents' states when population size goes to infinity

$$\bar{x}(t) = \lim_{N_t \rightarrow \infty} x^{N_t}(t) = \lim_{N \rightarrow \infty} \frac{1}{N_t} \sum_{i=1}^{N_t} x_i(t), \text{ a.s.}$$

Now, if the control strategy for each minor agent is considered to have the general feedback form

$$u_i = L_1 x_i + L_2 x_0 + \sum_{j \neq i, j=1}^{N_t} L_4 x_j + L_3, \quad 1 \leq i \leq N_t, \quad (13)$$

then the mean field dynamics can be obtained by substituting (13) in the minor liquidator (respectively, acquirer) agents' dynamics (10) (respectively, (11)), and taking the average and then its L^2 limit as $N \rightarrow \infty$.

The set of mean field equations for the optimal execution problem can be written as

$$d\bar{x} = \bar{A}\bar{x}dt + \bar{G}\bar{x}_0dt + \bar{m}dt. \quad (14)$$

For q_0 , $\bar{x} = [\bar{x}_a^T, \bar{x}_l^T]^T$ consists of the mean field \bar{x}_l of the liquidator population, and the mean field \bar{x}_a of the acquirer population. The matrices in (14) are defined as

$$\bar{A} = \begin{bmatrix} \bar{A}_a & \bar{A}_{al} \\ \bar{A}_{la} & \bar{A}_l \end{bmatrix}, \bar{G} = \begin{bmatrix} \bar{G}_a \\ \bar{G}_l \end{bmatrix}, \bar{m} = \begin{bmatrix} \bar{m}_a \\ \bar{m}_l \end{bmatrix}, \quad (15)$$

which shall be determined from consistency equations discussed in section III-E.

For q_a , $\bar{x} = \bar{x}_a$, and the matrices in (14) are given as

$$\bar{A} = \bar{A}_a, \quad \bar{G} = \bar{G}_a, \quad \bar{m} = \bar{m}_a. \quad (16)$$

For q_l , $\bar{x} = \bar{x}_l$, and the matrices in (14) are given by

$$\bar{A} = \bar{A}_l, \quad \bar{G} = \bar{G}_l, \quad \bar{m} = \bar{m}_l. \quad (17)$$

Finally, for q_2 , $\bar{x} = 0$.

D. Infinite Populations

Following the mean field game methodology with a major agent [40], the hybrid optimal execution problem is first solved in the infinite population case where the average term in the finite population dynamics and cost function of each agent is replaced by its infinite population limit, i.e. the mean field. Then specializing to linear systems [7], the major agent's state is extended with the mean field, while the minor agent's state is extended with the mean field and the major agent's state; this yields LQG problems for each trader linked only through the mean field and the major agent's state. Then the main results of [7], [40] are (i) the existence of infinite population best response strategies which yield the Nash equilibria, and (ii) the infinite population best response strategies applied to the finite population system yield an ϵ -Nash equilibria (see Theorem 3.1).

1) *Major Liquidator Agent:* The extended dynamics of the major agent in the infinite population, i.e. the dynamic for the x_0^{ex, q_l} is given by

$$dx_0^{ex, q_j} = (\mathbb{A}_0^{q_j} x_0^{ex, q_j} + \mathbb{M}_0^{q_j} + \mathbb{B}_0^{q_j} u_0^{q_j})dt + \mathbb{D}_0^{q_j} dW_0, \quad (18)$$

$0 \leq j \leq 2$, and the cost function for the extended major agent's system would be

$$J_0(u_0, u_{-0}) = \mathbb{E} \left[\|x_0^{ex, q_2}(T)\|_{\mathbb{P}^{q_2}}^2 + \sum_{j=0}^2 \int_{t_j}^{t_{j+1}} (\|x_0^{ex, q_j}(s)\|_{\mathbb{P}^{q_j}}^2 + \|u_0^{q_j}(s)\|_{R_0^{q_j}}^2) ds \right], \quad (19)$$

where $t_0 = 0, t_3 = T$. Over the interval $[t_0, t_1)$, and in the discrete state q_0 , the dynamics of the continuous state $x_0^{ex, q_0} = [x_0^T, \bar{x}_a^T, \bar{x}_l^T]^T$ is determined from (18) with

$$\mathbb{A}_0^{q_0} = \begin{bmatrix} A_0 & [E_0, E_0] \\ \bar{G} & \bar{A} \end{bmatrix}, \quad \mathbb{M}_0^{q_0} = \begin{bmatrix} 0_{3 \times 1} \\ \bar{m} \end{bmatrix}, \\ \mathbb{B}_0^{q_0} = \begin{bmatrix} B_0 \\ 0_{6 \times 1} \end{bmatrix}, \quad \mathbb{D}_0^{q_0} = \begin{bmatrix} D_0 & 0_{3 \times 6} \\ 0_{6 \times 3} & 0_{6 \times 6} \end{bmatrix}.$$

and $\mathbb{P}_0^{q_0}$ in (19) is given by

$$\mathbb{P}_0^{q_0} = [I_{3 \times 3}, 0_{3 \times 3}, 0_{3 \times 3}]^T P_0 [I_{3 \times 3}, 0_{3 \times 3}, 0_{3 \times 3}]$$

In case (i) where $q_1 = q_a$ over the interval $[t_1, t_2)$, the dynamics for $x_0^{ex, q_a} = [x_0^T, \bar{x}_a^T]^T$ is determined from (18) with

$$\mathbb{A}_0^{q_a} = \begin{bmatrix} A_0 & E_0 \\ \bar{G}_a & \bar{A}_a \end{bmatrix}, \quad \mathbb{M}_0^{q_a} = \begin{bmatrix} 0_{3 \times 1} \\ \bar{m}_a \end{bmatrix}, \\ \mathbb{B}_0^{q_a} = \begin{bmatrix} B_0 \\ 0_{3 \times 1} \end{bmatrix}, \quad \mathbb{D}_0^{q_a} = \begin{bmatrix} D_0 & 0_{3 \times 3} \\ 0_{3 \times 3} & 0_{3 \times 3} \end{bmatrix}.$$

and $\mathbb{P}_0^{q_a}$ is given by

$$\mathbb{P}_0^{q_a} = [I_{3 \times 3}, 0_{3 \times 3}]^T P_0 [I_{3 \times 3}, 0_{3 \times 3}].$$

In this case, the values of the continuous state before and after t_1 are related by the jump map

$$x_0^{ex, q_a}(t_1) = \Psi_{0,a} x_0^{ex, q_0}(t_1-) \quad (20)$$

where

$$\Psi_{0,a} = \begin{bmatrix} I_{3 \times 3} & 0_{3 \times 3} & 0_{3 \times 3} \\ 0_{3 \times 3} & I_{3 \times 3} & 0_{3 \times 3} \end{bmatrix}. \quad (21)$$

In case (ii) where $q_1 = q_l$ holds, $x_0^{ex, q_l} = [x_0^T, \bar{x}_l^T]^T$ and

$$\mathbb{A}_0^{q_l} = \begin{bmatrix} A_0 & E_0 \\ \bar{G}_l & \bar{A}_l \end{bmatrix}, \quad \mathbb{M}_0^{q_l} = \begin{bmatrix} 0_{3 \times 1} \\ \bar{m}_l \end{bmatrix}, \\ \mathbb{B}_0^{q_l} = \begin{bmatrix} B_0 \\ 0_{3 \times 1} \end{bmatrix}, \quad \mathbb{D}_0^{q_l} = \begin{bmatrix} D_0 & 0_{3 \times 3} \\ 0_{3 \times 3} & 0_{3 \times 3} \end{bmatrix}.$$

$$\mathbb{P}_0^{q_l} = [I_{3 \times 3}, 0_{3 \times 3}]^T P_0 [I_{3 \times 3}, 0_{3 \times 3}, 0_{3 \times 3}].$$

In this case, the values of the continuous state of the major trader before and after t_1 are related by the jump map

$$x_0^{ex, q_l}(t_1) = \Psi_{0,l} x_0^{ex, q_0}(t_1-) \quad (22)$$

where

$$\Psi_{0,l} = \begin{bmatrix} I_{3 \times 3} & 0_{3 \times 3} & 0_{3 \times 3} \\ 0_{3 \times 3} & 0_{3 \times 3} & I_{3 \times 3} \end{bmatrix}. \quad (23)$$

For the discrete state q_2 , the continuous state of the major trader is $x_0^{ex, q_2} \equiv x_0$, and

$$\mathbb{A}_0^{q_2} = A_0, \quad \mathbb{M}_0^{q_2} = 0_{3 \times 1}, \quad \mathbb{B}_0^{q_2} = B_0, \quad \mathbb{D}_0^{q_2} = D_0$$

$$\bar{\mathbb{P}}_0^{q_2} = \bar{P}_0, \quad \mathbb{P}_0^{q_2} = P_0$$

The values continuous state of the major trader before and after t_2 are related by the the jump map

$$x_0^{ex, q_2}(t_2) = \Psi_{0,2} x_0^{ex, q_1}(t_2-) \quad (24)$$

where

$$\Psi_{0,2} = \begin{bmatrix} I_{3 \times 3} & 0_{3 \times 3} \end{bmatrix}. \quad (25)$$

Best response hybrid control action:

By the definition of the terms \mathbb{D}_0^{qj} necessarily satisfy the condition A1 in [30], which in the LQG takes the following form

$$\mathbb{D}_0^{qj} = \Psi_{0,j} \mathbb{D}_0^{qj-1}, \quad j = 1, 2. \quad (26)$$

An application of the stochastic hybrid control theory of [30], specialized to the LQG case in [31], yield the infinite population best response hybrid control action as

$$u_0^{qj}(t) = -R_{0,qj}^{-1} \mathbb{B}_{0,qj}^T \Pi_0^{qj}(t) x_0^{ex,qj}(t), \quad (27)$$

where $\Pi_0^{qj}(t)$ is the solution of

$$-\dot{\Pi}_0^{qj} = \Pi_0^{qj} \mathbb{A}_0^{qj} + \mathbb{A}_{0,qj}^T \Pi_0^{qj} - \Pi_0^{qj} \mathbb{B}_0^{qj} R_{0,qj}^{-1} \mathbb{B}_{0,qj}^T \Pi_0^{qj} + \mathbb{P}_0, \quad (28)$$

subject to the terminal and boundary conditions

$$\Pi_0^{q2}(T) = \bar{\mathbb{P}}_0, \quad (29)$$

$$\Pi_0^{qj-1}(t_j) = \Psi_{0,j}^T \Pi_0^{qj}(t_j) \Psi_{0,j}, \quad (30)$$

$$\begin{aligned} & \mathbb{P}_0^{qj-1} + \Psi_{0,j}^T \Pi_0^{qj}(t_j) \Psi_{0,j} \mathbb{A}_0^{qj-1} + \mathbb{A}_{0,qj-1}^T \Psi_{0,j}^T \Pi_0^{qj}(t_j) \Psi_{0,j} \\ & - \Psi_{0,j}^T \Pi_0^{qj}(t_j) \Psi_{0,j} \mathbb{B}_0^{qj-1} R_{0,qj-1}^{-1} \mathbb{B}_{0,qj-1}^T \Psi_{0,j}^T \Pi_0^{qj}(t_j) \Psi_{0,j} \\ & = \Psi_{0,j}^T \left(\mathbb{P}_0^{qj} + \Pi_0^{qj}(t_j) \mathbb{A}_0^{qj} + \mathbb{A}_{0,qj}^T \Pi_0^{qj} \right. \\ & \left. - \Pi_0^{qj}(t_j) \mathbb{B}_0^{qj} R_{0,qj}^{-1} \mathbb{B}_{0,qj}^T \Pi_0^{qj}(t_j) \right) \Psi_{0,j}, \text{ for } j = 1, 2 \end{aligned} \quad (31)$$

We remark that $q_1 = q_a$ if $t_2^l < t_2^a$, where t_2^a denotes the time t_j , $j = 2$ calculated from (31) with the substitution of $q_1 = q_a$, and t_2^l denotes the time calculated with the substitution of $q_1 = q_l$. Otherwise, $q_1 = q_l$.

2) Minor Acquirer: A generic minor agent \mathcal{A}_i 's extended dynamics in the acquirer population with the extended state $x_i^{ex,qj}$ is

$$dx_i^{ex,qj} = (\mathbb{A}_a^{qj} x_i^{ex,qj} + \mathbb{M}_a^{qj} + \mathbb{B}_0^{qj} u_0^{qj} + \mathbb{B}_a^{qj} u_i^{qj}) dt + \mathbb{D}_a^{qj} dW_i, \quad (32)$$

where for q_0 , $x_i^{ex,q_0} = [x_i^T, x_0^T, \bar{x}_a^T, \bar{x}_l^T]^T$, and

$$\begin{aligned} \mathbb{A}_a^{q_0} &= \begin{bmatrix} A_a & [G_a, E_a, E_a] \\ 0_{9 \times 3} & \mathbb{A}_0^{q_0} - \mathbb{B}_0^{q_0} R_{0,q_0}^{-1} \mathbb{B}_{0,q_0}^T \Pi_0^{q_0} \end{bmatrix}, \\ \mathbb{M}_a^{q_0} &= \begin{bmatrix} 0_{3 \times 1} \\ \mathbb{M}_0 \end{bmatrix}, \quad \mathbb{B}_a^{q_0} = \begin{bmatrix} B_a \\ 0_{9 \times 1} \end{bmatrix}, \quad \mathbb{D}_a^{q_0} = \begin{bmatrix} D_a & 0_{3 \times 9} \\ 0_{9 \times 3} & \mathbb{D}_0^{q_0} \end{bmatrix}, \end{aligned}$$

and for q_a , $x_i^{ex,q_a} = [x_i^T, x_0^T, \bar{x}_a^T]^T$, and

$$\begin{aligned} \mathbb{A}_a^{q_a} &= \begin{bmatrix} A_a & [G_a, E_a] \\ 0_{6 \times 3} & \mathbb{A}_0^{q_a} - \mathbb{B}_0^{q_a} R_{0,q_a}^{-1} \mathbb{B}_{0,q_a}^T \Pi_0^{q_a} \end{bmatrix}, \\ \mathbb{M}_a^{q_a} &= \begin{bmatrix} 0_{3 \times 1} \\ \mathbb{M}_0 \end{bmatrix}, \quad \mathbb{B}_a^{q_a} = \begin{bmatrix} B_a \\ 0_{6 \times 1} \end{bmatrix}, \quad \mathbb{D}_a^{q_a} = \begin{bmatrix} D_a & 0_{3 \times 6} \\ 0_{6 \times 3} & \mathbb{D}_0^{q_a} \end{bmatrix}. \end{aligned}$$

In case (i) where the acquirer population is trading over $[t_1, t_2)$, i.e. $q_1 = q_a$, the total hybrid cost for a minor acquirer is given by

$$\begin{aligned} J_i^a(u_i, u_{-i}) &= \mathbb{E} \left[\|x_i^{ex,q_a}(t_2)\|_{\bar{\mathbb{P}}^{q_a}}^2 \right. \\ & \left. + \sum_{j=0}^1 \int_{t_j}^{t_{j+1}} (\|x_i^{ex,q_j}(s)\|_{\mathbb{P}_a^{q_j}}^2 + \|u_i^{q_j}(s)\|_{R_a^{q_j}}^2) ds \right], \end{aligned} \quad (33)$$

where

$$\bar{\mathbb{P}}_a^{q_a} = [I_{3 \times 3}, 0_{3 \times 6}]^T \bar{P} [I_{3 \times 3}, 0_{3 \times 6}] \quad (34)$$

$$\mathbb{P}_a^{q_a} = [I_{3 \times 3}, 0_{3 \times 6}]^T P [I_{3 \times 3}, 0_{3 \times 6}] \quad (35)$$

$$\mathbb{P}_a^{q_0} = [I_{3 \times 3}, 0_{3 \times 9}]^T P [I_{3 \times 3}, 0_{3 \times 9}]. \quad (36)$$

In this case, the extended state for a generic minor agent in the acquirer population at t_1 satisfies the jump transition map

$$x_i^{ex,q_a}(t_1) = \Psi_{i,a} x_i^{ex,q_0}(t_1 -)$$

with

$$\Psi_{i,a} = \begin{bmatrix} I_{3 \times 3} & 0_{3 \times 3} & 0_{3 \times 3} & 0_{3 \times 3} \\ 0_{3 \times 3} & I_{3 \times 3} & 0_{3 \times 3} & 0_{3 \times 3} \\ 0_{3 \times 3} & 0_{3 \times 3} & I_{3 \times 3} & 0_{3 \times 3} \end{bmatrix}.$$

In case (ii) where $q_1 = q_l$ holds over the interval $[t_1, t_2)$, the cost for the minor acquirer agent \mathcal{A}_i is given by

$$\begin{aligned} J_i^a(u_i, u_{-i}) &= \mathbb{E} \left[\|x_i^{ex,q_0}(t_1)\|_{\bar{\mathbb{P}}^{q_0}}^2 \right. \\ & \left. + \int_{t_0}^{t_1} (\|x_i^{ex,q_0}(s)\|_{\mathbb{P}_a^{q_0}}^2 + \|u_i^{q_0}(s)\|_{R_a^{q_0}}^2) ds \right], \end{aligned} \quad (37)$$

with $\bar{\mathbb{P}}^{q_0} = [I_{3 \times 3}, 0_{3 \times 9}]^T \bar{P} [I_{3 \times 3}, 0_{3 \times 9}]$.

Best response hybrid control action:

The optimal stopping problem for a minor acquirer is equivalent to a hybrid optimal control problem in which the dynamics and costs become zero after stopping. By the definition of the terms \mathbb{D}_a^{qj} necessarily satisfy the condition A1 in [30]. To be specific, for the case (i) the diffusion coefficients in (32) satisfy

$$\mathbb{D}_a^{q_a} = \Psi_{i,a} \mathbb{D}_a^{q_0}, \quad (38)$$

$$\mathbb{D}_a^{q_2} = \Psi_{i,\sigma_{q_a,q_{\text{stop}}}} \mathbb{D}_a^{q_a} \equiv 0, \quad (39)$$

where $\sigma_{q_j,q_{\text{stop}}}$ denotes the stopping event in the discrete state q_j . Both conditions in (39) are satisfied since $\mathbb{D}_a^{q_2} = 0$ due to the zero dynamics after stopping and $\Psi_{i,\sigma_{q_a,q_{\text{stop}}}} = 0$ due to removal of the minor acquirer trader's state from the market dynamics. For the case (ii) we also have

$$\mathbb{D}_a^{q_1} = \Psi_{i,\sigma_{q_0,q_{\text{stop}}}} \mathbb{D}_a^{q_0} \equiv 0, \quad (40)$$

which holds due to the stopping decision at t_1 . The results of [30], [31] yield

$$u_i^{qj}(t) = -R_{qj}^{-1} \mathbb{B}_{a,qj}^T \Pi_a^{qj}(t) x_i^{ex,qj}(t), \quad (41)$$

with

$$-\dot{\Pi}_a^{qj} = \Pi_a^{qj} \mathbb{A}_a^{qj} + \mathbb{A}_{a,qj}^T \Pi_a^{qj} - \Pi_a^{qj} \mathbb{B}_a^{qj} R_{a,qj}^{-1} \mathbb{B}_{a,qj}^T \Pi_a^{qj} + \mathbb{P}_a, \quad (42)$$

where for the case (i), in which $q_1 = q_a$, $\Pi_a^{q_j}(t)$ is the solution of (42) subject to the terminal conditions

$$\begin{aligned} \Pi_a^{q_a}(t_2) &= \bar{\mathbb{P}}_a^{q_a}, \\ \left(\mathbb{P}_a^{q_a} + \bar{\mathbb{P}}_a^{q_a} \mathbb{A}_a^{q_a} + \mathbb{A}_{a,q_a}^T \bar{\mathbb{P}}_a^{q_a} - \bar{\mathbb{P}}_a^{q_a} \mathbb{B}_{a,q_a}^{-1} \mathbb{B}_{a,q_a}^T \bar{\mathbb{P}}_a^{q_a} \right)_{t=t_2} &= 0 \end{aligned}$$

and the boundary conditions

$$\Pi_a^{q_0}(t_1) = \Psi_{i,a}^T \Pi_a^{q_a}(t_1) \Psi_{i,a}, \quad (43)$$

$$\begin{aligned} &\mathbb{P}_a^{q_0} + \Psi_{i,a}^T \Pi_a^{q_a}(t_1) \Psi_{i,a} \mathbb{A}_{a,q_0}^T + \mathbb{A}_{a,q_0}^T \Psi_{i,a}^T \Pi_a^{q_a}(t_1) \Psi_{i,a} \\ &- \Psi_{i,a}^T \Pi_a^{q_a}(t_1) \Psi_{i,a} \mathbb{B}_{a,q_0}^{-1} \mathbb{B}_{a,q_0}^T \Psi_{i,a}^T \Pi_a^{q_a}(t_1) \Psi_{i,a} \\ &= \Psi_{i,a}^T \left(\mathbb{P}_a^{q_a} + \Pi_a^{q_a}(t_1) \mathbb{A}_a^{q_a} + \mathbb{A}_{a,q_a}^T \Pi_a^{q_a}(t_1) \right. \\ &\quad \left. - \Pi_a^{q_a}(t_1) \mathbb{B}_{a,q_a}^{-1} \mathbb{B}_{a,q_a}^T \Pi_a^{q_a}(t_1) \right) \Psi_{i,a}, \quad (44) \end{aligned}$$

and in case (ii) where $q_1 = q_l$ holds, $\Pi_a^{q_0}(t)$ is the solution of (42) subject to the terminal conditions

$$\begin{aligned} \Pi_a^{q_0}(t_1) &= \bar{\mathbb{P}}_a^{q_0} \\ \left(\mathbb{P}_a^{q_0} + \bar{\mathbb{P}}_a^{q_0} \mathbb{A}_a^{q_0} + \mathbb{A}_{a,q_0}^T \bar{\mathbb{P}}_a^{q_0} - \bar{\mathbb{P}}_a^{q_0} \mathbb{B}_{a,q_0}^{-1} \mathbb{B}_{a,q_0}^T \bar{\mathbb{P}}_a^{q_0} \right)_{t=t_1} &= 0. \quad (46) \end{aligned}$$

We remark that the the solution of (44) (respectively (46)) result in the same stopping times t_2^a (respectively t_1) determined from (31).

3) *Minor Liquidator*: The hybrid dynamics, jump maps and performance measures for a minor liquidator are presented in a similar form as the minor acquirer, and therefore, due to space limitations, are not presented here. The infinite population best response hybrid control action as

$$u_i^{q_j}(t) = -R_{q_j}^{-1} \mathbb{B}_{l,q_j}^T \Pi_l^{q_j}(t) x_i^{e_x, q_j}(t), \quad (47)$$

with

$$-\dot{\Pi}_l^{q_j} = \Pi_l^{q_j} \mathbb{A}_l^{q_j} + \mathbb{A}_{l,q_j}^T \Pi_l^{q_j} - \Pi_l^{q_j} \mathbb{B}_{l,q_j}^{-1} \mathbb{B}_{l,q_j}^T \Pi_l^{q_j} + \mathbb{P}_l, \quad (48)$$

where for the case (i), in which $q_1 = q_a$, $\Pi_l^{q_j}(t)$ is the solution of (48) subject to the terminal conditions

$$\begin{aligned} \Pi_l^{q_0}(t_1) &= \bar{\mathbb{P}}_l^{q_0} \\ \left(\mathbb{P}_l^{q_0} + \bar{\mathbb{P}}_l^{q_0} \mathbb{A}_l^{q_0} + \mathbb{A}_{l,q_0}^T \bar{\mathbb{P}}_l^{q_0} - \bar{\mathbb{P}}_l^{q_0} \mathbb{B}_{l,q_0}^{-1} \mathbb{B}_{l,q_0}^T \bar{\mathbb{P}}_l^{q_0} \right)_{t=t_1} &= 0. \end{aligned}$$

and in case (ii) where $q_1 = q_l$ holds, $\Pi_l^{q_0}(t)$ is the solution of (42) subject to the terminal conditions

$$\begin{aligned} \Pi_l^{q_l}(t_2) &= \bar{\mathbb{P}}_l^{q_l}, \\ \left(\mathbb{P}_l^{q_l} + \bar{\mathbb{P}}_l^{q_l} \mathbb{A}_l^{q_l} + \mathbb{A}_{l,q_l}^T \bar{\mathbb{P}}_l^{q_l} - \bar{\mathbb{P}}_l^{q_l} \mathbb{B}_{l,q_l}^{-1} \mathbb{B}_{l,q_l}^T \bar{\mathbb{P}}_l^{q_l} \right)_{t=t_2} &= 0 \end{aligned}$$

and the boundary conditions

$$\Pi_l^{q_0}(t_1) = \Psi_{i,l}^T \Pi_l^{q_l}(t_1) \Psi_{i,l}, \quad (49)$$

$$\begin{aligned} &\mathbb{P}_l^{q_0} + \Psi_{i,l}^T \Pi_l^{q_l}(t_1) \Psi_{i,l} \mathbb{A}_{l,q_0}^T + \mathbb{A}_{l,q_0}^T \Psi_{i,l}^T \Pi_l^{q_l}(t_1) \Psi_{i,l} \\ &- \Psi_{i,l}^T \Pi_l^{q_l}(t_1) \Psi_{i,l} \mathbb{B}_{l,q_0}^{-1} \mathbb{B}_{l,q_0}^T \Psi_{i,l}^T \Pi_l^{q_l}(t_1) \Psi_{i,l} \\ &= \Psi_{i,l}^T \left(\mathbb{P}_l^{q_l} + \Pi_l^{q_l}(t_1) \mathbb{A}_l^{q_l} + \mathbb{A}_{l,q_l}^T \Pi_l^{q_l}(t_1) \right. \\ &\quad \left. - \Pi_l^{q_l}(t_1) \mathbb{B}_{l,q_l}^{-1} \mathbb{B}_{l,q_l}^T \Pi_l^{q_l}(t_1) \right) \Psi_{i,l}, \quad (50) \end{aligned}$$

The infinite population equilibria is linked to the finite population equilibria by the following theorem.

Theorem 3.1 (ϵ -Nash Equilibria for CO MM-MF Systems):

[7] Subject to reasonable technical conditions (see [7]), the mean field equations generate the set of control laws $\mathcal{U}_{MF}^N \triangleq \{u_i^0; 0 \leq i \leq N\}$, $1 \leq N < \infty$, given by (27), (41), and (47) such that

- (i) All agent systems \mathcal{A}_i , $0 \leq i \leq N$, are second order stable.
- (ii) $\{\mathcal{U}_{MF}^N; 1 \leq N < \infty\}$ yields an ϵ -Nash equilibrium for all ϵ , i.e. for all $\epsilon > 0$, there exists $N(\epsilon)$ such that for all $N \geq N(\epsilon)$;

$$J_i^{s,N}(u_i^0, u_{-i}^0) - \epsilon \leq \inf_{u_i \in \mathcal{U}_{i,y}^N} J_i^{s,N}(u_i, u_{-i}^0) \leq J_i^{s,N}(u_i^0, u_{-i}^0). \quad \square$$

E. Consistency Condition

The closed loop trading dynamics of a minor acquirer $\mathcal{A}_i \in \mathcal{A}_a$ applying (41), or correspondingly a minor liquidator $\mathcal{A}_i \in \mathcal{A}_l$ applying (47) is consequently

$$d\nu_i = -R_{a/l}^{-1} \mathbb{B}_{a/l}^T \Pi_{a/l}(x_i^T, x_0^T, \bar{x}^T)^T dt - R_{a/l}^{-1} \mathbb{B}_{a/l}^T s_{a/l}(t) dt,$$

then the average of closed loop trading dynamics over acquirer or liquidator population is obtained as

$$\begin{aligned} \frac{1}{N_{a/l}} \sum_{i=1}^{N_{a/l}} d\nu_i &= -\frac{1}{N_{a/l}} \sum_{i=1}^{N_{a/l}} R_{a/l}^{-1} \mathbb{B}_{a/l}^T \Pi_{a/l}(x_i^T, x_0^T, \bar{x}^T)^T dt \\ &\quad - \frac{1}{N_{a/l}} \sum_{i=1}^{N_{a/l}} R_{a/l}^{-1} \mathbb{B}_{a/l}^T s_{a/l}(t) dt, \quad (51) \end{aligned}$$

where $\bar{x} = [\bar{x}_a^T, \bar{x}_l^T]^T$. Then taking the L^2 limit of (51) as the population size $N_{a/l}$ goes to infinity yields the trading rate mean field dynamics

$$\begin{aligned} d\bar{\nu}_{a/l} &= \lim_{N_{a/l} \rightarrow \infty} d\nu^{N_{a/l}} = -R_{a/l}^{-1} \mathbb{B}_{a/l}^T \Pi_{a/l} \\ &\times \lim_{N_{a/l} \rightarrow \infty} ((x^{N_{a/l}})^T, x_0^T, \bar{x}_a^T, \bar{x}_l^T)^T dt - R_{a/l}^{-1} \mathbb{B}_{a/l}^T s_{a/l} dt, \end{aligned}$$

and hence the consistency equations become

$$\begin{aligned} \bar{A}_{a,11} &= -R_a^{-1}(\Pi_{a,11} + \Pi_{a,17}) - aR_a^{-1}(\Pi_{a,31} + \Pi_{a,37}), \\ \bar{A}_{a,12} &= -R_a^{-1}(\Pi_{a,12} + \Pi_{a,18}) - aR_a^{-1}(\Pi_{a,32} + \Pi_{a,38}), \\ \bar{A}_{a,13} &= -R_a^{-1}(\Pi_{a,13} + \Pi_{a,19}) - aR_a^{-1}(\Pi_{a,33} + \Pi_{a,39}), \\ \bar{A}_{al,11} &= -R_a^{-1}(\Pi_{a,110} + a\Pi_{a,310}), \\ \bar{A}_{al,12} &= -R_a^{-1}(\Pi_{a,111} + a\Pi_{a,311}) \\ \bar{A}_{al,13} &= -R_a^{-1}(\Pi_{a,112} + a\Pi_{a,312}) \\ \bar{G}_{a/l,11} &= -R_{a/l}^{-1}(\Pi_{a/l,14} + a\Pi_{a/l,34}), \\ \bar{G}_{a/l,12} &= -R_{a/l}^{-1}(\Pi_{a/l,15} + a\Pi_{a/l,35}), \\ \bar{G}_{a/l,13} &= -R_{a/l}^{-1}(\Pi_{a/l,16} + a\Pi_{a/l,36}), \\ \bar{m}_{a/l,1} &= 0, \quad (52) \end{aligned}$$

where $\Pi_{a/l,ij} = \Pi_{a/l}(i, j)$ for $i = \{1, 3\}$, $j = \{1, 2, 3, \dots, 12\}$. Hence the matrices in (15) are given as

$$\bar{A}_{a/l} = \begin{bmatrix} \bar{A}_{a/l,11} & \bar{A}_{a/l,12} & \bar{A}_{a/l,13} \\ 1 & 0 & 0 \\ (\lambda + a\bar{A}_{a/l,11}) & a\bar{A}_{a/l,12} & a\bar{A}_{a/l,13} \end{bmatrix},$$

$$\bar{A}_{al} = \begin{bmatrix} \bar{A}_{al,11} & \bar{A}_{al,12} & \bar{A}_{al,13} \\ 0 & 0 & 0 \\ a\bar{A}_{al,11} & a\bar{A}_{al,12} & a\bar{A}_{al,13} \end{bmatrix}, \bar{m} = \begin{bmatrix} \bar{m}_{a/l,1} \\ 0 \\ a\bar{m}_{a/l,1} \end{bmatrix}$$

$$\bar{A}_{la} = \begin{bmatrix} \bar{A}_{la,11} & \bar{A}_{la,12} & \bar{A}_{la,13} \\ 0 & 0 & 0 \\ a\bar{A}_{la,11} & a\bar{A}_{la,12} & a\bar{A}_{la,13} \end{bmatrix},$$

$$\bar{G}_{a/l} = \begin{bmatrix} \bar{G}_{a/l,12} & \bar{G}_{a/l,22} & \bar{G}_{a/l,23} \\ 0 & 0 & 0 \\ (\lambda_0 + a\bar{G}_{a/l,21}) & a\bar{G}_{a/l,22} & a\bar{G}_{a/l,23} \end{bmatrix}.$$

where the equations for the entries of matrices \bar{A}_{la} , \bar{A}_l due to space limitations are not presented (see [41]).

REFERENCES

- [1] M. Huang, P. E. Caines, and R. P. Malhamé, "Individual and mass behaviour in large population stochastic wireless power control problems: centralized and Nash equilibrium solutions," in *Proceedings of the 42nd IEEE Conference on Decision and Control*, 2003, pp. 98–103.
- [2] M. Huang, R. P. Malhamé, and P. E. Caines, "Large population stochastic dynamic games: Closed loop mckean-vlasov systems and the Nash certainty equivalence principle," *Special issue in honour of the 65th birthday of Tyrone Duncan, Communications in Information and Systems*, vol. 6, p. 221?252, 2006.
- [3] M. Huang, P. E. Caines, and R. P. Malhamé, "Large population cost-coupled lqg problems with non-uniform agents: individual-mass behaviour and decentralized ϵ -Nash equilibria," *IEEE Transactions on Automatic Control*, vol. 52, no. 9, p. 1560?1571, 2007.
- [4] P. E. Caines, "Mean field games," *Encyclopedia of Systems and Control*, pp. 1–6, 2014. [Online]. Available: [Online]. Available: <http://dx.doi.org/10.1007/978-1-4471-5102-930-1>
- [5] J. M. Lasry and P. L. Lions, "Jeux a champ stationnaire," *C. R. Acad. Sci*, vol. 343, pp. 619–625, 2006.
- [6] —, "Jeux a champ moyen. ii - horizon fini et controle optimal," *C. R. Acad. Sci*, vol. 343, pp. 679–684, 2006.
- [7] M. Huang, "Large-population lqg games involving a major player: The Nash certainty equivalence principle," *SIAM Journal on Control and Optimization*, vol. 48, no. 5, pp. 3318–3353, 2010.
- [8] S. L. Nguyen and M. Huang, "Linear-quadratic-gaussian mixed games with continuum-parametrized minor players," *SIAM Journal on Control and Optimization*, vol. 50, no. 5, pp. 2907–2937, 2012.
- [9] P. E. Caines and A. C. Kizilkale, "Recursive estimation of common partially observed disturbances in mfg systems with application to large scale power markets," in *Proceedings of the 52nd IEEE Conference on Decision and Control*, 2013, pp. 2505–2512.
- [10] —, "Mean field estimation for partially observed lqg systems with major and minor agents," in *Proceedings of the 19th World Congress of the International Federation of Automatic Control (IFAC)*, 2014, pp. 8705–8709.
- [11] —, " ϵ -Nash equilibria for partially observed LQG mean field games with major player," *IEEE Transaction on Automatic Control*, 2016. [Online]. Available: <http://ieeexplore.ieee.org/stamp/stamp.jsp?arnumber=7778116>
- [12] N. Şen and P. E. Caines, "Mean field games with partially observed major player and stochastic mean field," in *Proceedings of the 53rd IEEE Conference on Decision and Control*, 2014, pp. 2709–2715.
- [13] —, " ϵ -Nash equilibria for a partially observed mean field game with major player," in *Proceedings of the 2015 American Control Conference (ACC)*, 2015, pp. 4791–4797.
- [14] —, "Mean field game theory with a partially observed major agent," *SIAM Journal on Control and Optimization*, vol. 54, no. 6, pp. 3174–3224, 2016. [Online]. Available: <http://dx.doi.org/10.1137/16M1063010>
- [15] D. Firoozi and P. E. Caines, " ϵ -Nash equilibria for partially observed lqg mean field games with major agent: Partial observations by all agents," in *Proceedings of the 54th IEEE Conference on Decision and Control (CDC)*, 2015, pp. 4430–4437.
- [16] F. H. Clarke and R. B. Vinter, "Optimal Multiprocesses," *SIAM Journal on Control and Optimization*, vol. 27, no. 5, pp. 1072–1091, 1989.
- [17] A. Bensoussan and J. L. Menaldi, "Hybrid Control and Dynamic Programming," *Dynamics of Continuous, Discrete and Impulsive Systems Series B: Application and Algorithm*, vol. 3, no. 4, pp. 395–442, 1997.
- [18] J. Lygeros, C. Tomlin, and S. Sastry, "Multiobjective Hybrid Controller Synthesis," in *Hybrid and Real-Time Systems, International Workshop*, 1997, pp. 109–123.
- [19] M. S. Branicky, V. S. Borkar, and S. K. Mitter, "A Unified Framework for Hybrid Control: Model and Optimal Control Theory," *IEEE Transactions on Automatic Control*, vol. 43, no. 1, pp. 31–45, 1998.
- [20] H. J. Sussmann, "A Nonsmooth Hybrid Maximum Principle," *Lecture Notes in Control and Information Sciences, Springer London, Volume 246*, pp. 325–354, 1999.
- [21] X. Xu and P. J. Antsaklis, "Optimal Control of Switched Systems based on Parameterization of the Switching Instants," *IEEE Transactions on Automatic Control*, vol. 49, no. 1, pp. 2–16, 2004.
- [22] S. Dharmatti and M. Ramaswamy, "Hybrid Control Systems and Viscosity Solutions," *SIAM Journal on Control and Optimization*, vol. 44, no. 4, pp. 1259–1288, 2005.
- [23] M. Garavello and B. Piccoli, "Hybrid Necessary Principle," *SIAM Journal on Control and Optimization*, vol. 43, no. 5, pp. 1867–1887, 2005.
- [24] M. S. Shaikh and P. E. Caines, "On the Hybrid Optimal Control Problem: Theory and Algorithms," *IEEE Transactions on Automatic Control*, vol. 52, no. 9, pp. 1587–1603, 2007, corrigendum: vol. 54, no. 6, pp. 1428, 2009.
- [25] F. Taringoo and P. E. Caines, "On the Optimal Control of Impulsive Hybrid Systems on Riemannian Manifolds," *SIAM Journal on Control and Optimization*, vol. 51, no. 4, pp. 3127–3153, 2013.
- [26] A. Pakniyat and P. E. Caines, "On the Relation between the Minimum Principle and Dynamic Programming for Classical and Hybrid Control Systems," *IEEE Transactions on Automatic Control*, 2017, doi: 10.1109/TAC.2017.2667043.
- [27] —, "Hybrid Optimal Control of an Electric Vehicle with a Dual-Planetary Transmission," *Nonlinear Analysis: Hybrid Systems*, 2016, DOI: 10.1016/j.nahs.2016.08.004.
- [28] A. Bensoussan and J. Menaldi, "Stochastic Hybrid Control," *Mathematical Analysis and Applications*, vol. 249, no. 1, pp. 261–288, 2000.
- [29] C. Aghayeva and Q. Abushov, "The Maximum Principle for the Nonlinear Stochastic Optimal Control Problem of Switching Systems," *Global Optimization*, vol. 56, no. 2, pp. 341–352, 2011.
- [30] A. Pakniyat and P. E. Caines, "On the Stochastic Minimum Principle for Hybrid Systems," in *Proceedings of the 55th IEEE Conference on Decision and Control, Las Vegas, NV, USA*, 2016, pp. 1139–1144.
- [31] —, "A Class of Linear Quadratic Gaussian Hybrid Optimal Control Problems with Realization-Independent Riccati Equations," in (to appear) *Proceedings of the 20th World Congress of the International Federation of Automatic Control, IFAC, Toulouse, France*, 2017.
- [32] Á. Cartea and S. Jaimungal and J. Penalva, *Algorithmic and High-Frequency Trading*. United Kingdom : Cambridge University Press, 2015.
- [33] S. Jaimungal and D. Kinzebulatov, "Optimal execution with a price limiter," *Journal of Risk*, 2014.
- [34] R. Almgren and N. Chriss, "Optimal execution of portfolio transactions," *Journal of Risk*, pp. 5–39, 2001.
- [35] A. Alfonsi, A. Fruth, and A. Schied, "Optimal execution strategies in limit order books with general shape functions," *Quantitative Finance*, vol. 10, no. 2, pp. 143–157, 2010.
- [36] E. Bayraktar and M. Ludkovski, "Optimal trade execution in illiquid markets," *Mathematical Finance*, vol. 21, no. 4, pp. 681–701, 2011.
- [37] S. Jaimungal and M. Nourian, "Mean-field game strategies for a major-minor agent optimal execution problem," *Available at SSRN*, 2015.
- [38] D. Firoozi and P. E. Caines, "Mean field game ϵ -Nash equilibria for partially observed optimal execution problems in finance," in *Proceedings of the 55th IEEE Conference on Decision and Control*, 2016.
- [39] —, "An optimal execution problem in finance with acquisition and liquidation objectives: an MFG formulation," in (to appear) *Proceedings of the 20th World Congress of the International Federation of Automatic Control (IFAC)*, 2017.
- [40] M. Nourian and P. E. Caines, " ϵ -Nash mean field game theory for nonlinear stochastic dynamical systems with major and minor agents," *SIAM Journal on Control and Optimization*, vol. 51, no. 4, pp. 3302–3331, 2013.
- [41] D. Firoozi and P. E. Caines, "Optimal execution problem in finance with mean field tracking," (in preparation) *research report, CIM, McGill University*, 2017.