

# Mathematical Statistics Recitation 2

Section 002 (Prof. Niles-Weed)

TA: Sonia Reilly

February 6, 2026

Reminder: Midterm 1 on Thursday Feb 12! Covers Lectures 1-6 (everything up through today).

## Lecture Review

### Lecture 5

#### Covariance

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}X)(Y - \mathbb{E}Y)] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2 \text{Cov}(X, Y)$$

Correlation coefficient:

$$\rho = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- $-1 \leq \rho \leq 1$
- If  $X, Y$  are independent, then  $\rho = 0$  (not the converse).
- $\rho = \pm 1$  indicates perfect (linear) correlation.

#### Conditional Expectation and Variance

Conditional Expectation:

$$\mathbb{E}[Y \mid X = x] = \int y f_{Y|X}(y \mid x) dy,$$

which is a function of  $x$ .  $\mathbb{E}[Y \mid X]$  is a random variable.

Law of Total Expectation:

$$\mathbb{E}[Y] = \mathbb{E}[\mathbb{E}[Y \mid X]]$$

(inner expectation in  $Y$ , outer expectation in  $X$ )

Law of Total Variance

$$\text{Var}(Y) = \text{Var}(\mathbb{E}[Y \mid X]) + \mathbb{E}[\text{Var}(Y \mid X)]$$

## Lecture 6

### Convergence of Random Variables

Convergence in probability:

$$X_n \xrightarrow{p} X \quad \text{if} \quad \mathbb{P}(|X_n - X| > \varepsilon) \rightarrow 0 \quad \forall \varepsilon$$

Convergence in distribution:

$$X_n \xrightarrow{d} X \quad \text{if} \quad \lim_{n \rightarrow \infty} F_n(x) = F(x),$$

where  $F_n, F$  are CDFs of  $X_n, X$ .

### Moment Generating Functions

$$M(t) = \mathbb{E}(e^{tX})$$

uniquely determines the distribution if it exists.

Moment generating because  $k$ -th derivative at 0 is  $k$ -th moment:

$$M^{(k)}(0) = \mathbb{E}(X^k)$$

$$M_n(t) \rightarrow M(t) \quad \text{in open interval around 0} \implies F_n(t) \rightarrow F(t)$$

(used to prove CLT).

### Law of Large Numbers (LLN)

$$\frac{1}{n} \sum_{i=1}^n X_i \xrightarrow{p} \mathbb{E}[X_i] \quad \text{as } n \rightarrow \infty$$

if the  $X_i$  are iid. (Proof: Uses Chebyshev's inequality.)

### Central Limit Theorem:

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1) \quad \text{as } n \rightarrow \infty$$

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

for  $X_i$  i.i.d. with mean  $\mu$ , variance  $\sigma^2$ , and MGF defined in neighborhood of 0.  
Can interpret as

$$\bar{X}_n \sim N\left(\mu, \frac{\sigma^2}{n}\right) \quad \text{approximately, for large } n$$

(rule of thumb:  $n \geq 30$ )

### Reminder from probability: Gamma Distribution

pdf with parameters  $\alpha, \lambda$ :

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \geq 0$$

$$\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} dt.$$

In particular,  $\Gamma(n) = (n-1)!$  for integers.

Note: only the constant scaling is complicated. At heart, the distribution is just a mix of powers of  $x$  and exponentials of  $x$ .




Figure 1: Gamma distributions for different values of  $\alpha$  and  $\lambda$  (here  $\theta = 1/\lambda$ ). Credit: Wikipedia.

- $\alpha$  — shape parameter
- $\lambda$  — scale parameter (units of  $x$ , does not affect shape)

Family of probability distributions that includes special cases:

- exponential (for  $\alpha = 1$ )
- $\chi_n^2$  (for  $\alpha = \frac{n}{2}$ ,  $\lambda = \frac{1}{2}$ ) — we'll learn more about this later
- approximates normal as  $\alpha \rightarrow \infty$

## Problems

1. [4.74] The number of offspring of an organism is a discrete random variable with mean  $\mu$  and variance  $\sigma^2$ . Each of its offspring reproduces in the same manner. Find the expectation and variance of the number of offspring in the third generation.

**Solution:**

Let  $X_1 = 1$  be the number of organisms in the first generation,  $X_2$  be the number of offspring it has, and  $X_3$  be the total number of their offspring (the number in the third generation). Thus we know that

$$\mathbb{E}[X_2] = \mu, \quad \text{Var}(X_2) = \sigma^2.$$

We also have the same expectation and variance for the offspring of each second-generation organism. So

$$\mathbb{E}[X_3 | X_2] = X_2\mu, \quad \text{Var}(X_3 | X_2) = X_2\sigma^2$$

(because  $\text{Var}(\sum X_i) = \sum \text{Var}(X_i)$  for independent  $X_i$ 's). By the law of total expectation,

$$\mathbb{E}[X_3] = \mathbb{E}[\mathbb{E}[X_3 | X_2]] = \mathbb{E}[X_2\mu] = \boxed{\mu^2}.$$

By the law of total variance,

$$\begin{aligned}\text{Var}(X_3) &= \text{Var}(\mathbb{E}[X_3 \mid X_2]) + \mathbb{E}[\text{Var}(X_3 \mid X_2)] \\ &= \text{Var}(X_2\mu) + \mathbb{E}[X_2\sigma^2] \\ &= \sigma^2\mu^2 + \mu\sigma^2 \\ &= \boxed{\sigma^2\mu(\mu + 1)}.\end{aligned}$$

2. [5.18] Suppose that a company ships packages that are variable in weight, with an average weight of 15 lb and a standard deviation of 10 lb. Assuming that the packages come from a large number of different customers so that it is reasonable to model their weights as independent random variables, find the probability that 100 packages will have a total weight exceeding 1700 lb.

**Solution:**

Use the CLT

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$$

with

$$\mu = 15, \quad \sigma = 10.$$

Letting  $\Phi$  denote the CDF of  $N(0, 1)$ ,

$$\begin{aligned}\mathbb{P}\left(\sum_{i=1}^{100} X_i > 1700\right) &= \mathbb{P}(\bar{X}_{100} > 17) \\ &= \mathbb{P}\left(\frac{\bar{X}_{100} - 15}{10/\sqrt{100}} > 2\right) \\ &\approx 1 - \Phi(2) \\ &= 1 - 0.9772 = \boxed{0.0228}\end{aligned}$$

3. Find the mean and variance of a gamma-distributed random variable

$$X \sim \text{Gamma}(\alpha, \lambda).$$

- (a) Find the moment generating function of the gamma distribution.  
(b) Find the mean and variance of  $X$ .

**Solution:**

(a)

$$\begin{aligned}M(t) &= \mathbb{E}[e^{tX}] = \int_0^\infty e^{tx} \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} dx \\ &= \frac{\lambda^\alpha}{\Gamma(\alpha)} \int_0^\infty x^{\alpha-1} e^{-x(\lambda-t)} dx\end{aligned}$$

We know from the pdf of the gamma distribution that

$$\int_0^\infty \frac{L^A}{\Gamma(A)} x^{A-1} e^{-Lx} dx = 1 \implies \int_0^\infty x^{A-1} e^{-Lx} dx = \frac{\Gamma(A)}{L^A}.$$

So letting  $A = \alpha$ ,  $L = \lambda - t$ , we get

$$\int_0^\infty x^{\alpha-1} e^{-x(\lambda-t)} dx = \frac{\Gamma(\alpha)}{(\lambda-t)^\alpha}.$$

So

$$M(t) = \frac{\lambda^\alpha}{\Gamma(\alpha)} \frac{\Gamma(\alpha)}{(\lambda-t)^\alpha} = \left( \frac{\lambda}{\lambda-t} \right)^\alpha.$$

(b) To find the moments, we differentiate  $M(t)$ :

$$M'(t) = \alpha \left( \frac{\lambda}{\lambda-t} \right)^{\alpha-1} \left( \frac{\lambda}{(\lambda-t)^2} \right) = \frac{\alpha}{\lambda} \left( \frac{\lambda}{\lambda-t} \right)^{\alpha+1}$$

$$M''(t) = \frac{\alpha(\alpha+1)}{\lambda} \left( \frac{\lambda}{\lambda-t} \right)^\alpha \left( \frac{\lambda}{(\lambda-t)^2} \right) = \frac{\alpha(\alpha+1)}{\lambda^2} \left( \frac{\lambda}{\lambda-t} \right)^{\alpha+2}$$

So

$$M'(0) = \boxed{\mathbb{E}X = \frac{\alpha}{\lambda}}$$

$$M''(0) = \mathbb{E}[X^2] = \frac{\alpha(\alpha+1)}{\lambda^2}$$

$$\Rightarrow \text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}X)^2 = \frac{\alpha(\alpha+1)}{\lambda^2} - \frac{\alpha^2}{\lambda^2} = \boxed{\frac{\alpha}{\lambda^2}}.$$

4. Show that if  $X_i \stackrel{\text{iid}}{\sim} \text{Exp}(\lambda)$ , then

$$Y = \sum_{i=1}^n X_i \sim \text{Gamma}(n, \lambda).$$

**Solution:**

Recall that  $\text{Exp}(\lambda) = \text{Gamma}(1, \lambda)$ . We already found the mgf of  $\text{Gamma}(\alpha, \lambda)$ ,

$$\left( \frac{\lambda}{\lambda-t} \right)^\alpha,$$

so the mgf of  $\text{Exp}(\lambda)$  is

$$\frac{\lambda}{\lambda-t}.$$

Since the  $X_i$ 's are independent,

$$\begin{aligned} M_Y(t) &= \mathbb{E} [e^{tY}] = \mathbb{E} [e^{t \sum_{i=1}^n X_i}] = \mathbb{E} \left[ \prod_{i=1}^n e^{tX_i} \right] \\ &= \prod_{i=1}^n \mathbb{E} [e^{tX_i}] = \prod_{i=1}^n \left( \frac{\lambda}{\lambda-t} \right) = \left( \frac{\lambda}{\lambda-t} \right)^n. \end{aligned}$$

This is the same as the mgf of  $\text{Gamma}(n, \lambda)$ , and since we know that the mgf uniquely describes distributions, we know that

$$Y \sim \text{Gamma}(n, \lambda).$$