

Determination of Functional Network Structure from Local Parameter Dependence Data

Boris N. Kholodenko

Department of Pathology, Anatomy and Cell Biology

Thomas Jefferson University, PA

Boris.Kholodenko@mail.tju.edu

Eduardo D. Sontag

Dept. of Mathematics

Rutgers University, NJ, USA

sontag@hilbert.rutgers.edu

Abstract

In many applications, such as those arising from the field of cellular networks, it is often desired to determine the interaction (graph) structure of a set of differential equations, using as data measured sensitivities. This note proposes an approach to this problem.

1 Introduction

Suppose given a system of differential equations

$$\begin{aligned}\dot{x}_1 &= f_1(x_1, \dots, x_n, p_1, \dots, p_m) \\ \dot{x}_2 &= f_2(x_1, \dots, x_n, p_1, \dots, p_m) \\ &\vdots \\ \dot{x}_n &= f_n(x_1, \dots, x_n, p_1, \dots, p_m),\end{aligned}\tag{1}$$

where the vector of state variables $\mathbf{x}(t) = (x_1(t), \dots, x_n(t))$ evolves in some open subset $X \subseteq \mathbb{R}^n$ and the vector of parameters $p = (p_1, \dots, p_m)$ can be chosen from an open subset $\mathcal{P} \subseteq \mathbb{R}^m$, for some n and m . In cellular networks, for instance, the state variables x_i might represent the concentrations of certain proteins, mRNA, etc., at different time instants, and the parameters p_i might represent the concentration levels of certain enzymes which are maintained at a constant value during a particular experiment (see, e.g., [1, 2]).

In many fields of application, it is often the case that the equations defining the system (that is, the form of the functions f_i describing the vector field) are unknown, even in general form, but one wishes nonetheless to determine the interaction graph of a system (1), that is to say, to know which variables directly influence which variables, as well as the relative strengths of these interactions ([3]). A more limited goal might be to determine if a certain feedback loop is present in the system (cf. [4]).

To help in this task, experimental data is usually available, measuring solutions of system (1) for various initial states and parameter settings. A special case, the one treated in this note, is that in which experimental data provides us with the location of steady states associated to parameter values near a given set of parameters $\bar{p}$. We show in this note how, starting from such data, and under an assumption that seems natural, it is indeed possible to determine this interaction graph and relative strengths of interactions. We then extend the method to non-steady state measurements. A more global analysis of the problem is also possible, and will be presented in a follow-up report.

2 Problem Formulation

The steady states of system (1) associated to a given parameter vector $p = (p_1, \dots, p_m)$ are the solutions $\mathbf{x} = (x_1, \dots, x_n)$ of the set of n simultaneous algebraic equations

$$\begin{aligned} f_1(x_1, \dots, x_n, p_1, \dots, p_m) &= 0 \\ f_2(x_1, \dots, x_n, p_1, \dots, p_m) &= 0 \\ &\vdots \\ f_n(x_1, \dots, x_n, p_1, \dots, p_m) &= 0. \end{aligned} \tag{2}$$

We will assume that there is a function $\xi : \mathcal{P} \rightarrow \mathcal{X}$ which assigns, to each parameter vector $p \in \mathcal{P}$, a steady state $\xi(p)$ of (1), that is to say, one has that $f_i(\xi(p), p) = 0$ for all $i = 1, \dots, n$ and all $p \in \mathcal{P}$. We suppose that the functions f_i as well as ξ are continuously differentiable. A particular parameter vector $\bar{p}$ is also given, and the problem will be formulated in terms of the behavior of steady states near $\bar{\mathbf{x}} = \xi(\bar{p})$.

Experimental Data

It will be assumed that an $n \times m$ matrix $\Sigma = (\sigma_{kj})$ is given, representing the “sensitivities”

$$\sigma_{kj} = \frac{\partial \xi_k}{\partial p_j}(\bar{p}) \tag{3}$$

for each $k = 1, \dots, n$ and each $j = 1, \dots, m$. This matrix of partial derivatives may be estimated numerically from the values of $\xi(p)$ on a neighborhood of the chosen parameter $\bar{p}$.

Desired Information

Consider the $n \times n$ matrix $A = (a_{ij})$ defined by:

$$a_{ij} = \frac{\partial f_i}{\partial x_j}(\bar{\mathbf{x}}, \bar{p})$$

for every $i, j = 1, \dots, n$.

Ideally, one would want to find the matrix A , since this matrix completely describes the influence of each variable x_j upon the rate of change of each other variable x_i . Unfortunately, such an objective is impossible to achieve from the local steady-state data Σ , or even from the knowledge of the complete (global) mapping ξ . This is because the same mapping ξ also solves the set of equations

$$\begin{aligned} \lambda_1 f_1(x_1, \dots, x_n, p_1, \dots, p_m) &= 0 \\ \lambda_2 f_2(x_1, \dots, x_n, p_1, \dots, p_m) &= 0 \\ &\vdots \\ \lambda_n f_n(x_1, \dots, x_n, p_1, \dots, p_m) &= 0, \end{aligned} \tag{4}$$

for any constants λ_i , but, on the other hand, multiplication of f_i by λ_i results in $a_{ij} = \lambda_i a_{ij}$. In other words, the best that one could hope is for the data Σ to determine the rows

$$A_i = (a_{i1}, \dots, a_{in}), \quad i = 1, \dots, n$$

of A only up to scalar multiples.

Thus, a more realistic objective is to attempt to identify the rows A_i up to a scalar multiple only. For example, if we assume that $a_{ii} \neq 0$ for each i (a realistic assumption when stable systems are being interconnected), this amounts to finding the ratios a_{ij}/a_{ii} for each $i \neq j$.

Assumptions

We will make two assumptions which will suffice for us to solve the problem of determining the rows A_i of A up to scalar multiples. The first assumption is a strong but reasonable structural one, while the second represents a weak algebraic nondegeneracy condition.

We will suppose known, for each $i \in \{1, \dots, n\}$, a subset S_i of the index set $\{1, \dots, m\}$ so that the following property holds:

$$(\forall j \in S_i) \quad \frac{\partial f_i}{\partial p_j}(\bar{\mathbf{x}}, \bar{p}) = 0 \quad (5)$$

which is in turn implied by the structural condition:

$$(\forall j \in S_i) \quad f_i \text{ does not depend upon } p_j.$$

This prior information about the system structure is far less restrictive than it might appear at first sight. Indeed, it is usually the case that “compartmental” information is available, for instance telling us that the concentration of a certain enzyme has no direct influence on an unrelated biochemical reaction, that an extracellular signaling molecule does not affect directly a cytoplasmatic reaction, and so forth.

The second assumption is as follows. For each $j \in \{1, \dots, m\}$, we introduce the vector

$$\Sigma_j = \text{col}(\sigma_{1j}, \dots, \sigma_{nj}) = \begin{pmatrix} \frac{\partial \xi_1}{\partial p_j}(\bar{p}) \\ \vdots \\ \frac{\partial \xi_n}{\partial p_j}(\bar{p}) \end{pmatrix} \quad (6)$$

representing the j th column of the matrix Σ , and we consider, for each $i \in \{1, \dots, n\}$ the linear subspace H_i of $\mathbb{R}^n$ spanned by the vectors

$$\{\Sigma_j \mid j \in S_i\}.$$

The assumption is:

$$\dim H_i \geq n - 1 \quad \forall i = 1, \dots, n.$$

Note that this amounts to saying that the dimension of H_i is either n or $n - 1$. (Generically, we may expect this dimension to be $n - 1$, because the orthogonality relation to be shown below is the only algebraic constraint.)

3 Solution

With the above assumptions, the problem that we posed can be solved as follows. We fix any index $i \in \{1, \dots, n\}$, then take partial derivatives in the equation $F_i(p) = f_i(\xi(p), p) = 0$ with respect to the variable p_j , for each index j in the set S_i , and finally evaluate at $\mathbf{x} = \bar{\mathbf{x}}$ and $p = \bar{p}$:

$$0 = \frac{\partial F_i}{\partial p_j}(\bar{p}) = \sum_{k=1}^n \frac{\partial f_i}{\partial x_k}(\bar{\mathbf{x}}, \bar{p}) \frac{\partial \xi_k}{\partial p_j}(\bar{\mathbf{x}}) + \frac{\partial f_i}{\partial p_j}(\bar{\mathbf{x}}, \bar{p}) = A_i \cdot \Sigma_j$$

where the second term vanishes by assumption (5).

Since this happens for every $j \in S_i$, we conclude that the vector A_i is orthogonal to H_i . As H_i has dimension n or $n - 1$, this determines A_i up to a scalar multiple, which is what we wanted to prove.

Of course, it is trivial to actually compute A_i from the data. If H_i has dimension n , then $A_i = 0$; this is a degenerate case. When the dimension is $n - 1$, one simply picks a basis $\{\Sigma_{j_1}, \dots, \Sigma_{j_{n-1}}\}$ of H_i , and any vector Σ_0 linearly independent from the elements of this basis (a randomly chosen vector has this property), and then solves (for example) the nonsingular set of equations $A_i \cdot \Sigma_0 = 1$, $A_i \cdot \Sigma_{j_\ell} = 0$, $\ell = 1, \dots, n - 1$, to find a nonzero A_i (all possible A_i are scalar multiples of this one). Alternatively, provided that one knows that $a_{ii} \neq 0$, one may simply normalize to $a_{ii} = -1$ and then determine the

remaining entries of A_i by solving a linear set of $n - 1$ equations. Observe also, that if it is known *a priori* that certain entries a_{ij} vanish, then one may redefine the space H_i to be spanned only by the vectors listing the appropriate components of the sensitivities $\partial \xi_i / \partial p_j$'s, and a potentially much smaller number of parameter perturbations may be required.

4 Modular Approach

It is also possible to apply our techniques in a “modular” context, in which only the derivatives $\partial f_i / \partial x_j$ with respect to communicating intermediaries are calculated ([4]). Let us briefly explain this.

We assume that the entire network consists of an interconnection of n subsystems or “modules”, each of which is described by a set of differential equations such as:

$$\begin{aligned}\dot{x}_j &= g_j(\mathbf{y}_j, x_1, \dots, x_n, p_1, \dots, p_m) \\ \dot{\mathbf{y}}_j &= G_j(\mathbf{y}_j, x_1, \dots, x_n, p_1, \dots, p_m),\end{aligned}\tag{7}$$

where the variables x_j represent “communicating” or “connecting” intermediaries of module j that transmit information to other modules, whereas the *vector variables* $\mathbf{y}_j$ represent chemical species that interact within module j . Each vector $\mathbf{y}_j$ has dimension ℓ_j . The integers ℓ_j , $j = 1, \dots, n$ are in general different for each of the n modules, and they represent one less than the number of chemical species in the j th module respectively. Observe that, for each j , g_j is function from an open subset of $\mathbb{R}^{\ell_j + n + m}$ into $\mathbb{R}$, and G_j is function from an open subset of $\mathbb{R}^{\ell_j + n + m}$ into $\mathbb{R}^{\ell_j}$.

We will assume that, for each fixed module, the Jacobian of G_j with respect to the vector variable $\mathbf{y}_j$ evaluated at the steady state corresponding to $\bar{p}$ (assumed to exist, as before) is nonsingular. The Implicit Mapping Theorem then implies that one may, in a neighborhood of this steady state, solve

$$G_j(\mathbf{y}_j, x_1, \dots, x_n, p_1, \dots, p_m) = 0$$

for the vector variable $\mathbf{y}_j$ as a function of $x_1, \dots, x_n, p_1, \dots, p_m$, the solution being given locally by a function

$$\mathbf{y}_j = \mathbf{M}_j(x_1, \dots, x_n, p_1, \dots, p_m).$$

One concludes that, around this steady state corresponding to $\bar{p}$, the functions x_j satisfy implicit equations of the form

$$0 = g_j(\mathbf{M}_j(x_1, \dots, x_n, p_1, \dots, p_m), x_1, \dots, x_n, p_1, \dots, p_m) = h_j(x_1, \dots, x_n, p_1, \dots, p_m)$$

so the analysis then proceeds as before, using the h_j 's instead of the f_j 's. The generalization to the case of more than one communicating intermediate in a module, namely a vector $(x_{j,1}, \dots, x_{j,k_j})$, is obvious.

5 Avoiding Derivatives

The technique that was described assumes that we know the sensitivity matrix Σ , which is obtained by evaluating the partial derivatives $\partial \xi_i / \partial p_j$ at the particular parameter value $\bar{p}$. Ordinarily, these derivatives would be estimated by finite differences. For instance, suppose that one measures $\bar{\mathbf{x}} = \xi(\bar{p})$ as well as $\xi(\bar{p} + d_j p)$, where

$$d_j p = \text{col}(0, \dots, 0, dp_j, 0, \dots, 0)$$

(entry in j th position) and we view dp_j as a “small” perturbation of the j th parameter. Denoting

$$d_j x_i := \xi_i(\bar{p} + d_j p) - \bar{x}_i\tag{8}$$

obviously one may estimate Σ using the following approximation:

$$\frac{\partial \xi_i}{\partial p_j}(\bar{p}) \approx \frac{d_j x_i}{dp_j}.$$

In order to calculate this ratio, both $d_j x_i$ and dp_j must be known.

However, in certain experimental situations it may well be impossible to estimate the values of dp_j . This might appear to be contradictory, since we are assuming that we perform experiments which change the values of p . But one can easily envision an experimental setup in which a certain external variable (in a cell biology situation, for instance, a growth factor) is known to influence a certain parameter p_j . Varying this external variable therefore produces a perturbation in p_j , and hence an appropriate $d_j x$ which is measured, but dp_j itself may be hard to measure.

It is rather surprising that we can still achieve our goal of estimating the rows of A (up to scalar multiples) even in the absence of information about the dp_j 's! To see intuitively why this is plausible, consider the following argument. Let us say that we have just a scalar parameter p and a scalar function $f(x_1, x_2)$ so that $f(\xi_1(p), \xi_2(p)) \equiv 0$, and that $\bar{\mathbf{x}} = \xi(\bar{p}) = (0, 0)$. In a neighborhood of $p = \bar{p}$, we may assume that f is linear, so we have a linear relation (with unknown coefficients)

$$a_1 \xi_1(p) + a_2 \xi_2(p) = 0.$$

The method discussed so far would take derivatives at $p = \bar{p}$:

$$a_1 \frac{d\xi_1}{dp}(\bar{p}) + a_2 \frac{d\xi_2}{dp}(\bar{p}) = 0$$

and thus (assuming that the derivative is not zero), we know that the row (a_1, a_2) must be a multiple of $(-(d\xi_2/dp)(p), (d\xi_1/dp)(p))$. A completely different argument (analogous to using a two-point as opposed to a slope-point formula in order to find the equation of a line) would simply take the original equation $a_1 \xi_1(p) + a_2 \xi_2(p) = 0$ (valid only $p \approx \bar{p}$, since this was an approximation of f) and say that the row (a_1, a_2) must be a multiple of $(-\xi_2(p), \xi_1(p))$, for any fixed $p \approx \bar{p}$. There is no inconsistency between the two estimates, since they only differ (approximately) by multiplication by the scalar dp :

$$(-\xi_2(p), \xi_1(p)) \approx (-(d\xi_2/dp)(p), (d\xi_1/dp)(p)) dp$$

and we only care about scalar multiples. Let us now say this in general.

Since $f_i(\xi(\bar{p} + d_j p), \bar{p} + d_j p) - f(\bar{\mathbf{x}}, \bar{p}) = 0 - 0 = 0$, we have, taking a Taylor expansion, that

$$\left. \frac{d}{dp_j} f_i(\xi(p), p) \right|_{p=\bar{p}} dp_j + o(dp_j) = \sum_{k=0}^n \frac{\partial f_i}{\partial x_k}(\bar{\mathbf{x}}, \bar{p}) \frac{\partial \xi_k}{\partial p_j}(\bar{p}) dp_j + o(dp_j) = 0. \quad (9)$$

whenever $j \in S_i$. Substituting

$$d_j x_k = \xi_k(\bar{p} + d_j p) - \bar{x}_k = \frac{\partial \xi_k}{\partial p_j}(\bar{p}) dp_j + o(dp_j) \quad (10)$$

into (9), we conclude that

$$\sum_{k=0}^n \frac{\partial f_i}{\partial x_k}(\bar{\mathbf{x}}, \bar{p}) d_j x_k = o(dp_j) \text{ provided that } j \in S_i.$$

Since $dp_j \approx 0$ and $d_j x_k = O(dp_j)$, this is an approximate orthogonality relation, and we now make the approximation:

$$\sum_{k=0}^n \frac{\partial f_i}{\partial x_k}(\bar{\mathbf{x}}, \bar{p}) d_j x_k = 0 \text{ provided that } j \in S_i. \quad (11)$$

In conclusion, and introducing the matrix $\Gamma = (\gamma_{kj}) = (d_j x_k)$ instead of Σ , we have that $A_i \cdot \Gamma_j = 0$ for all $j \in S_i$, where $\Gamma_j = \text{col}(\gamma_{1j}, \dots, \gamma_{nj})$. Now we consider, for each $i \in \{1, \dots, n\}$, the linear subspace K_i of $\mathbb{R}^n$ spanned by the vectors $\{\Gamma_j \mid j \in S_i\}$ and assume that $\dim K_i \geq n - 1$ for all i . We conclude that the vector A_i is orthogonal to K_i , and this once again determines A_i up to a scalar multiple.

6 Non-Steady State Analysis

Let us sketch here how one might extend our methodology to use non-steady state data. In general, we denote by $\xi(t, x^0, p)$ the solution of (1) with initial condition x^0 , at time t and using parameters p . Let us suppose that we can measure the sensitivities (3) at some specific point in time, and for some specific solution $\xi(\bar{t}, \bar{x}^0, \bar{p})$:

$$\sigma_{kj} = \frac{\partial \xi_k}{\partial p_j}(\bar{t}, \bar{x}^0, \bar{p}) \quad (12)$$

for each $k = 1, \dots, n$ and each $j = 1, \dots, m$, and we let $\Sigma = (\sigma_{kj})$. We also need now the mixed second derivatives:

$$\eta_{ij} = \frac{\partial^2 \xi_i}{\partial p_j \partial t}(\bar{t}, \bar{x}^0, \bar{p})$$

and instead of $\Sigma_j = \text{col}(\sigma_{1j}, \dots, \sigma_{nj})$ as in (6), we consider for each $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, m\}$ the vector

$$\Sigma_{ij} = \text{col}(\eta_{ij}, \sigma_{1j}, \dots, \sigma_{nj}). \quad (13)$$

We define, for each $i \in \{1, \dots, n\}$, H_i as the linear subspace of $\mathbb{R}^{n+1}$ spanned by the vectors $\{\Sigma_{ij} \mid j \in S_i\}$. We let now $a_{ij} = \frac{\partial f_i}{\partial x_j}(\bar{t}, \bar{x}^0, \bar{p})$. Fixing any index $i \in \{1, \dots, n\}$, we take partial derivatives on both sides of the differential equation

$$\frac{\partial}{\partial t} \xi_i(t, x^0, p) = f_i(\xi(t, x^0, p), p)$$

with respect to the variable p_j , for each index j in the set S_i , and evaluate at $\mathbf{x} = \bar{\mathbf{x}}$, $t = \bar{t}$, and $p = \bar{p}$, to obtain:

$$\eta_{ij} = \frac{\partial}{\partial p_j} f_i(\bar{\mathbf{x}}, \bar{p}) = \sum_{k=1}^n a_{ik} \sigma_{kj}$$

from which we conclude that $[-1, A_i] \cdot \Sigma_j = 0$ whenever $j \in S_i$, and hence that $[-1, A_i]$ is orthogonal to H_i . With appropriate genericity conditions, this orthogonality, perhaps in conjunction with conditions at other times t or points p , will restrict the possible vectors A_i and more generally the interaction graph. (For example, if $\dim H_i = n + 1$, then we have a unique solution.) Derivatives with respect to parameter values can be replaced by differences, just as in the steady state case. We will discuss this further in a future contribution.

References

- [1] Kholodenko, B.N., Demin, O.V., Moehren, G., and Hoek, J.B. (1999) "Quantification of short term signaling by the epidermal growth factor receptor," *J. Biol. Chem.* 274, 30169-30181.
- [2] Moehren, G., Markevich, N., Demin, O., Kiyatkin, A., Goryanin, I., Hoek, J.B., and Kholodenko, B.N. (2002) "Temperature dependence of the epidermal growth factor receptor signaling network can be accounted for by a kinetic model," *Biochemistry* 41, 306-320.
- [3] Chevalier, T., Schreiber, T., and Ross, J. (1993) "Toward a systematic determination of complex reaction mechanisms," *J. Phys. Chem.* 97, 6776-6787.
- [4] Kholodenko, B.N., Kiyatkin A., Bruggeman F., Sontag E.D., Westerhoff H., Hoek J. (2002) "Untangling the wires: a novel strategy to trace functional interactions in signaling and gene networks," submitted for publication.