

## MATH 104: WORKSHEET 7

### 1. Concepts

(1) Differentiation Rules

### 2. Discussions

*Question 1.* Consider the following three derivatives of functions

$$[Df]_0 = \begin{bmatrix} 1 & -3 & 0 \\ 2 & 7 & 1 \\ 3 & 1 & 0 \\ -2 & 1 & -1 \end{bmatrix},$$

$$[Dg]_0 = \begin{bmatrix} 4 & 1 \\ -2 & 0 \\ -1 & 1 \end{bmatrix},$$

$$[Dh]_0 = \begin{bmatrix} 3 & -1 & 2 & 0 \\ -2 & -1 & 1 & 4 \\ 5 & 0 & 4 & 2 \end{bmatrix}.$$

Which derivatives can you compute at the origin among:

$$f \circ g, \quad g \circ f, \quad g \circ h, \quad h \circ g, \quad f \circ h, \quad h \circ f?$$

*Question 2.* Consider the functions

$$f(x, y, z) = \begin{pmatrix} x^2 + yz^2 \\ 2x + y^3 - z \end{pmatrix}$$

and

$$g(u, v) = \begin{pmatrix} u^2 - v^2 \\ uv \\ 3u - 2v \end{pmatrix}$$

- (1) Compute  $[Df]$  and  $[Dg]$
- (2) Compute  $[D(f \circ g)]_a$  where  $a = (1, 1, 1)$ .
- (3) If all inputs of  $f \circ g$  equal 1 what is the most sensitive output of  $f \circ g$ ?

*Question 3.* Consider the function  $f(x) = u(x)^{v(x)}$ , where  $f, u, v : \mathbb{R} \rightarrow \mathbb{R}$  are all single variable functions.

---

*Date:* 01/31/2024.

- (1) Compute  $f'(x)$  using your technique from Calculus.
- (2) Find  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $h : \mathbb{R} \rightarrow \mathbb{R}^2$  so that you can rephrase  $f = g \circ h$ . Compute  $[Df]_x$  using the multivariable chain rule and compare with the previous result.

*Question 4.* What is the derivative of the function  $f(x) = (Q(x))^2$  where  $Q(x) = x \cdot Ax$ ?

Here,  $A$  is a square  $n \times n$  matrix and  $x \in \mathbb{R}^n$ .

*Question 5.* Let  $f, h : \mathbb{R} \rightarrow \mathbb{R}$  and define

$$F(x) = \int_0^{h(x)} f(t) dt.$$

Determine  $[DF]$  using what we learn in this class. Check your answer with what you learned in Calculus.