MATH 104: Multivariable Calculus (brief notes)

Truong-Son Van

Contents

1 Vectors **7**
1.1 Basics 7
1.2 Products 8
1.2.1 Dot product 8
1.2.2 3D special: Cross product 9

Spring 2024

Chapter 1

Vectors

1.1 Basics

Reading: Stewart Chapter 12, Thomas Calculus Chapter 12, Active Calculus Chapter 9

You should be able to answer the following questions after reading this section:

- What is a vector?
- What does it mean for two vectors to be equal?
- How do we add two vectors together and multiply a vector by a scalar?
- How do we determine the magnitude of a vector?
- What is a unit vector
- How do we find a unit vector in the direction of a given vector?

Typically, we talk about 3-dimensional vectors (as discussed in Stewart and Thomas). However, since talking about n-dimensional vectors doesn't require much more effort, we will talk about n-dimensional vectors instead.

Definition 1.1. An n-dimensional Euclidean space $\mathbb{R}^n$ is the Cartesian product of n Euclidean spaces $\mathbb{R}$.

Definition 1.2. An n-dimensional vector $\mathbf{v} \in \mathbb{R}^n$ is a tuple

$$\mathbf{v} = \langle v_1, \dots, v_n \rangle , \tag{1.1}$$

where $v_i \in \mathbb{R}$.

In dimensions less than or equal to 3, we represent a vector geometrically by an arrow, whose length represents the magnitude.

Remark. A point in $\mathbb{R}^n$ is also represented by an n-tuple but with round brackets. A vector connecting two points $A = (a_1, \dots, a_n)$ and $B = (b_1, \dots, b_n)$ can be constructed as

$$\mathbf{x} = \langle b_1 - a_1, \dots, b_n - a_n \rangle .$$

We denote the above vector as $\vec{AB}$ where A is the tail (initial point) and B is the tip/head (terminal point). We denote $\mathbf{0}$ to be the zero vector, i.e.,

$$\mathbf{0} = \langle 0, \dots, 0 \rangle .$$

Definition 1.3. The length of a vector $\mathbf{v}$ (denoted by $|\mathbf{v}|$) is defined to be

$$|\mathbf{v}| = \sqrt{v_1^2 + \cdots + v_n^2} . \tag{1.2}$$

Definition 1.4. A unit vector is a vector that has magnitude 1.

Exercise 1.1. Turn a vector $\mathbf{v} \in \mathbb{R}^n$ into a unit vector with the same direction.

Rules to manipulate vectors

Let $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ and $c, d \in \mathbb{R}$. Then,

$$c(\mathbf{a}+\mathbf{b}) = \langle ca_1 + cb_1, \dots, ca_n + cb_n \rangle = c\mathbf{a} + c\mathbf{b}\,,$$

and

$$(c+d)\mathbf{a} = c\mathbf{a} + d\mathbf{a}\,.$$

These formulas are deceptively simple. Make sure you understand all the implications.

Because of this rule, sometimes it is good to write vectors in terms of elementary vectors:

$$\mathbf{u} = u_1\mathbf{e_1} + \cdots + u_n\mathbf{e_n}\,,$$

where $e_i = \langle 0, \dots, 1, \dots, 0 \rangle$ is the vector which has zero at all entries except that the i^{th} entry is 1.

In 3D,

$$\mathbf{e_1} = \mathbf{i}\,, \qquad \mathbf{e_2} = \mathbf{j}\,, \qquad \mathbf{e_3} = \mathbf{k}\,.$$

Properties of vector operations

Read the book

(Make sure you understand the geometric intepretation)

1.2 Products

1.2.1 Dot product

- How is the dot product of two vectors defined and what geometric information does it tell us?
- How can we tell if two vectors in $\mathbb{R}^n$ are perpendicular?
- How do we find the projection of one vector onto another?

Definition 1.5. The dot product of vectors $\mathbf{u} = \langle u_1, \dots, u_n \rangle$ and $\mathbf{v} = \langle v_1, \dots, v_n \rangle$ in $\mathbb{R}^n$ is the scalar

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + \cdots + u_nv_n\,.$$

Properties of dot product

Let $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$. Then,

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$,
2. $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = (\mathbf{u} \cdot \mathbf{w}) + (\mathbf{v} \cdot \mathbf{w})$,
3. If c is a scalar, then $(c\mathbf{u}) \cdot \mathbf{w} = c(\mathbf{u} \cdot \mathbf{w})$.

Theorem 1.1 (Law of cosine)**.** *If θ is the angle between the vectors $\boldsymbol{u}$ and $\boldsymbol{v}$, then*

$$\boldsymbol{u} \cdot \boldsymbol{v} = |\boldsymbol{u}||\boldsymbol{v}| \cos\theta\,.$$

Corollary 1.1. *Two vectors $\boldsymbol{u}$ and $\boldsymbol{v}$ are orthogonal to each other if $\boldsymbol{u} \cdot \boldsymbol{v} = 0$.*

Projection

Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$. The component of $\mathbf{u}$ in the direction of $\mathbf{v}$ is the scalar

$$\text{comp}_{\mathbf{v}}\mathbf{u} = \frac{\mathbf{u}\cdot\mathbf{v}}{|\mathbf{v}|}\,,$$

and the projection of $\mathbf{u}$ onto $\mathbf{v}$ is the vector

$$\text{proj}_{\mathbf{v}}\mathbf{u} = \left(\mathbf{u}\cdot\frac{\mathbf{v}}{|\mathbf{v}|}\right)\frac{\mathbf{v}}{|\mathbf{v}|} = \frac{\mathbf{u}\cdot\mathbf{v}}{\mathbf{v}\cdot\mathbf{v}}\mathbf{v}\,.$$

Read the book for more details. Make sure you understand the geometric meaning.

1.2.2 3D special: Cross product

This concept is very specific to $\mathbb{R}^3$. It will not make sense in other dimensions.

Definition 1.6. Let $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$. The cross product of $\mathbf{a}$ and $\mathbf{b}$ is defined to be

$$\mathbf{a}\times\mathbf{b} = \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1\rangle\,.$$

Theorem 1.2. *Let θ be the angle between $\mathbf{a}$ and $\mathbf{b}$. Then,*

$$|\mathbf{a}\times\mathbf{b}| = |\mathbf{a}||\mathbf{b}|\sin\theta\,.$$

Theorem 1.3. *The vector $\mathbf{a}\times\mathbf{b}$ is orthogonal to both $\mathbf{a}$ and $\mathbf{b}$.*