

$$P(E, S, B, W, G) = P(E)P(B)P(S|E, B)P(W|S)P(G|S)$$

P(E)	е	-e
	1/10	9/10
(SIF R)	c	_c

P(B)	b	-b
	1/10	9/10

P(S E,B)	S	-S
e ∧ b	9/10	1/10
e ∧ -b	2/10	8/10
-e ∧ b	8/10	2/10
-e ∧ -b	0	1

/	- /	
P(W S)	W	-W
S	8/10	2/10
-S	2/10	8/10

P(G S)	g	-g
S	1/2	1/2
-S	0	1

► Given the alarm went off (s) what is the probability that Mrs. Gibbons phones you (g)?

Given the alarm went off (s) what is the probability that Mrs. Gibbons phones you (g)? probability that the alarm went off (s)?

$$P(g|s) = 1/2$$

► Given that Mrs. Gibbons phones you (g) what is the probability the alarm went off (s)?

- ► Given that Mrs. Gibbons phones you (g) what is the probability the alarm went off (s)?
- 1. Bayes Rule says: P(S|g) = P(g|S) * P(S)/P(g)
- 2. P(-s|g) = P(g|-s) * P(-s)/P(g) = 0 * P(-s)/P(g) = 0.
- 3. Therefore P(s|g) = 1 (P(s|g) + P(-s|g) must sum to 1.

$$P(s|g) = 1 P(-s|g) = 0$$

Alternatively: $-s \rightarrow -g$, so $g \rightarrow s$, so P(s|g) = 1.



▶ What is P(G|S)? (i.e., the four probability values) P(g|s), P(-g|s), P(g|-s), P(-g|-s).

What is P(G|S)? (i.e., the four probability values P(g|s), P(-g|s), P(g|-s), P(-g|-s). $P(g|s) = 1/2 \qquad P(-g|s) = 1/2 \qquad P(-g|-s) = 1$

▶ What is $P(G|S \wedge W)$? (i.e., the 8 probability values $P(g|s \wedge w)$, $P(g|s \wedge -w)$, ..., $P(-g|-s \wedge -w)$).

▶ What is $P(G|S \land W)$? (i.e., the 8 probability values $P(g|s \land w), P(g|s \land -w), \ldots, P(-g|-s \land -w)$).

$$P(g|s,-w) = P(g|s,w) = P(g|s) = 1/2$$

 $P(-g|s,-w) = P(-g|s,w) = P(-g|s) = 1/2$
 $P(g|-s,-w) = P(g|-s,w) = P(g|-s) = 0$
 $P(-g|-s,-w) = P(-g|-s,w) = P(g|-s) = 1$

► What do these values tell us about the relationship between *G*, *W* and *S*?

▶ What is P(G|W)? (i.e., the four probability values P(g|w), P(-g|w), P(g|-w), and P(-g|-w)).

▶ What is P(G|W)? (i.e., the four probability values P(g|w), P(-g|w), P(g|-w), and P(-g|-w)).

Must do variable elimination.

- ▶ What is P(G|W)? (i.e., the four probability values P(g|w), P(-g|w), P(g|-w), and P(-g|-w)).
- Query variable is G.
- ▶ First run of VE, evidence is W = w.
- ▶ Second run of VE, evidence is W = -w.
- ▶ Use same ordering for both runs of VE: E, B, S, G.
- ▶ With same ordering some factors can be reused between the two runs of VE.

▶ What is P(G|W)? (i.e., the four probability values P(g|w), P(-g|w), P(g|-w), and P(-g|-w)).

- 1. E: P(E), P(S|E,B)
- 2. B: P(B),
- 3. **S**: P(w|S), P(S|G)
- 4. **G**:

What is P(G|W)? (i.e., the four probability values P(g|w), P(-g|w), P(g|-w), and P(-g|-w)).
1. E: P(E), P(S|E,B)
2. B: P(B),
3. S: P(w|S), P(S|G)

$$F_1(S,B) = \sum_{E} P(E) \times P(S|E,B)$$

= $P(e) \times P(S|e,B) + P(-e) \times P(S|-e,B)$

$$F_1(-s,-b) = P(e)P(-s,e,-b) + P(-e)P(-s,-e,-b)$$

= $0.1 \times 0.8 + 0.9 \times 1 = 0.98$

$$F_1(-s,b) = P(e)P(-s,e,b) + P(-e)P(-s,-e,b)$$

= $0.1 \times 0.1 + 0.9 \times 0.2 = 0.19$

$$F_1(s,-b) = P(e)P(s,e,-b) + P(-e)P(s,-e,-b)$$

$$= 0.1 \times 0.2 + 0.9 \times 0 = 0.02$$

$$F_1(s,b) = P(e)P(s,e,b) + P(-e)P(s,-e,b)$$

= 0.1 × 0.9 + 0.9 × 0.8 = 0.81



```
1. E: P(E), P(S|E,B)
2. B: P(B), F_1(S, B)
3. S: P(w|S), P(S|G)
4. G:
    F_2(S) = \sum_B P(B) \times F_1(S, B)
           = P(b)F_1(S,b) + P(-b)F_1(S,-b)
  F_2(-s) = P(b)F_1(-s,b) + P(-b)F_1(-s,-b)
           = 0.1 \times 0.19 + 0.9 \times 0.98 = 0.901
  F_2(s) = P(b)F_1(s,b) + P(-b)F_1(s,-b)
           = 0.1 \times 0.81 + 0.9 \times 0.02 = 0.099
```

- 1. E: P(E), P(S|E,B)
- 2. *B*: P(B), $F_1(S, B)$
- 3. S: P(w|S), P(S|G), $F_2(S)$
- 4. *G*:

$$F_3(G) = \sum_{S} P(w|S) \times P(S|G) \times F_2(S) = P(w|s)P(s|G)F_2(s) + P(w|-s)P(-s|G)F_2(-s)$$

$$F_3(-g) = P(w|s)P(s|-g)F_2(s) + P(w|-s)P(-s|-g)F_2(-s)$$

$$= 0.8 \times 0.5 \times 0.099 + 0.2 \times 1 \times 0.901 = 0.2198$$

$$F_3(g) = P(w|s)P(s|g)F_2(s) + P(w|-s)P(-s|g)F_2(-s)$$

$$F_3(g) = P(w|s)P(s|g)F_2(s) + P(w|-s)P(-s|g)F_2(-s)$$

= 0.8 \times 0.5 \times 0.099 + 0.2 \times 0 \times 0.901 = \frac{0.0396}{0.0396}

- 1. E: P(E), P(S|E,B)
- 2. *B*: P(B), $F_1(S, B)$
- 3. S: P(w|S), P(S|G), $F_2(S)$
- 4. $G: F_3(G)$

Normalize $F_3(G)$:

$$P(-g|w) = \frac{0.2198}{0.2198 + 0.0396} = 0.8473$$

$$P(g|w) = \frac{0.0396}{0.2198 + 0.0396} = 0.1527$$

- Now P(G|-w)?
 - 1. E: P(E), P(S|E,B)
 - 2. B: P(B),
 - 3. *S*: P(-w|S), P(S|G)
 - 4. G:

Already computed as $F_1(S, B)$

- 1. E: P(E), P(S|E,B)
- 2. *B*: P(B), $F_1(S, B)$
- 3. *S*: P(-w|S), P(S|G)
- 4. G:

Already computed as $F_2(S)$

- 1. E: P(E), P(S|E,B)
- 2. *B*: P(B), $F_1(S, B)$
- 3. *S*: P(-w|S), P(S|G), $F_2(S)$
- 4. *G*:

$$F_3(G) = \sum_{S} P(-w|S) \times P(S|G) \times F_2(S) = P(-w|s)P(s|G)F_2(s) + P(-w|-s)P(-s|G)F_2(-s)$$

$$F_{3}(-g) = P(-w|s)P(s|-g)F_{2}(s) + P(-w|-s)P(-s|-g)F_{2}(-s)$$

$$= 0.2 \times 0.5 \times 0.099 + 0.8 \times 1 \times 0.901 = 0.7307$$

$$F_{3}(g) = P(-w|s)P(s|g)F_{2}(s) + P(-w|-s)P(-s|g)F_{2}(-s)$$

$$= 0.2 \times 0.5 \times 0.099 + 0.8 \times 0 \times 0.901 = 0.0099$$

- 1. E: P(E), P(S|E,B)
- 2. B: P(B), $F_1(S, B)$
- 3. *S*: P(-w|S), P(S|G), $F_2(S)$
- 4. $G: F_3(G)$

Normalize $F_3(G)$:

$$P(-g|w) = \frac{0.7307}{0.7307 + 0.0099} = 0.9866$$

$$P(g|w) = \frac{0.0099}{0.2198 + 0.00099} = 0.0134$$

What do these values tell us about the relationship between G and W, and why does this relationship differ when we know S?

What do these values tell us about the relationship between G and W, and why does this relationship differ when we know S?

 ${\cal G}$ and ${\cal W}$ are not independent of each other. But when ${\cal S}$ is known they become independent.