Protocol Synopsis: BREATHE-2 (FICTIONAL, for demonstration only)

A Phase 3, Randomized, Double-Blind, Placebo- and Active-Controlled Study of SF-202 in Adults with Moderate Persistent Asthma

1. Study Design

Participants will be randomized 2:2:1 to one of three treatment groups for 24 weeks, followed by a 4-week safety follow-up.

2. Treatment Arms

Arm A: SF-202 20 mg once daily (N=200)

Arm B: Active comparator fluticasone 250 mcg twice daily (N=200)

Arm C: Matching placebo once daily (N=100)

3. Objectives and Endpoints

Primary objective: Evaluate the efficacy of SF-202 compared with placebo

Endpoint: Change from baseline in pre-bronchodilator FEV1 at Week 24

Secondary objective: Compare SF-202 with the active comparator on asthma control

Endpoint: Change from baseline in ACQ-5 score at Week 24

Secondary objective: Evaluate the safety and tolerability of SF-202

Endpoint: Incidence of treatment-emergent adverse events through Week 28

Exploratory objective: Explore the effect of SF-202 on airway inflammation

Endpoint: Change from baseline in FeNO at Weeks 12 and 24

4. Key Eligibility Criteria

Adults aged 18 to 75 with a physician diagnosis of asthma for at least 12 months, pre-bronchodilator FEV1 between 50% and 85% of predicted, and an ACQ-5 score of 1.5 or higher at screening. Current smokers are excluded.