

ASSIGNED FORUM PROBLEM 4

Soobin Rho

July 8, 2022

Chapter 8.8: *Improper Integrals*

$$\mathbf{33.} \quad \int_{-1}^{\infty} \frac{d\theta}{\theta^2 + 5\theta + 6} \tag{1}$$

SOLUTION

The first thing that comes to my mind when I look at this problem is that the numerator can be factored into $(\theta + 3)(\theta + 2)$ which means our problem becomes

$$\mathbf{33.} \quad \int_{-1}^{\infty} \frac{d\theta}{\theta^2 + 5\theta + 6} = \int_{-1}^{\infty} \frac{d\theta}{(\theta + 3)(\theta + 2)} \tag{2}$$

After that, we can use partial fraction decomposition to make the integrand easier to integrate.

$$\int_{-1}^{\infty} \frac{d\theta}{(\theta + 3)(\theta + 2)} = \int_{-1}^{\infty} \frac{A}{\theta + 3} + \frac{B}{\theta + 2} d\theta \tag{3}$$

We can then multiply both sides of the equation with $(\theta + 3)(\theta + 2)$. For the sake of readability, let's skip the integral notation for now and look at only the coefficients now.

$$1 = A(\theta + 2) + B(\theta + 3) \tag{4}$$

$$(0)\theta + (1) = (A + B)\theta + (3A + 2B) \tag{5}$$

Solving for these two equations $A + B = 0$ and $3A + 2B = 1$ gives us

$$A = 1 \tag{6}$$

$$B = -1 \tag{7}$$

In other words, if we put everything back to the original equation together, we get something that's much more easier to integrate:

$$\int_{-1}^{\infty} \frac{d\theta}{(\theta + 3)(\theta + 2)} = \int_{-1}^{\infty} \frac{1}{\theta + 3} + \frac{-1}{\theta + 2} d\theta$$

Well, why is it easier to integrate? It's because $\int \frac{1}{x}$ is elegantly $\ln|x| + C$.

$$\int_{-1}^{\infty} \frac{d\theta}{(\theta+3)(\theta+2)} = \left[\ln|\theta+2| - \ln|\theta+3| + C \right]_{-1}^{\infty} \quad (8)$$

$$= \left[0 \right] - \left[-\ln 2 \right] \quad (9)$$

$$= \ln 2 \quad (10)$$