

Dataset: BRITISH NATIONAL HEALTH DPOLL
Analyses: Cronbach's Alphas, PCA, Means Tests

Version 6: New Analyses Highlighted

Variable names in dataset:

1. "t1payhlth" & "t2payhlth"
2. "t1poora" & "t2poora"
3. "tloptions" & "t2options"
4. « t1hlthfundoptions » & « t2hlthfundoptions »
5. « t1ctexptrtmentb » & « t2ctexptrtmentb »
6. "t1prtreath" & "t2prtreath"
7. « t1wl » & « t2wl »
8. "t1severity" & "t2severity"
9. "t1prevention" & "t2prevention"
10. "t1qoflife" & "t2qoflife"
11. « t1dispubgvt » & « t2dispubgvt »
12. « t1avgdisdod » & « t2avgdisdod »
13. "t1moresay" & "t2moresay"

Indices: DEPENDENT VARIABLES

1. PAYHLTH = Government Pay For Healthcare

Who should pay for healthcare? Which of these three statements comes closest to your view? Govt pay for everyone OR Govt pay for the poor OR People pay own w/ insurance?

*In Dataset: "t1payhlth" & "t2payhlth"

Coding: 0 = individual should pay, 1 = government should pay

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2payh~h	193	.8937824	.0147596	.2050475	.8646705	.9228942
t1payh~h	193	.8186528	.01884	.2617335	.7814929	.8558128
diff	193	.0751295	.0195484	.2715755	.0365723	.1136868

Ho: mean(t2payhlth - t1payhlth) = mean(diff) = 0

Ha: mean(diff) < 0	Ha: mean(diff) != 0	Ha: mean(diff) > 0
t = 3.8433	t = 3.8433	t = 3.8433
P < t = 0.9999	P > t = 0.0002	P > t = 0.0001

2. POORA = Priority to poor

Please say how much you agree or disagree with the following. If the NHS had to choose, priority should be given to treating people who can't afford private medical treatment? (5-pt scale)

*In Dataset: "t1poora" & "t2poora"

Coding: 0 = strongly disagree, 1 = strongly agree

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2poora	213	.5481221	.0226026	.3298745	.5035674	.5926768
t1poora	213	.6126761	.0204697	.2987456	.5723258	.6530263

diff	213	-.064554	.0241907	.3530519	-.1122391	-.0168688
------	-----	----------	----------	----------	-----------	-----------

Ho: mean(t2poora - t1poora) = mean(diff) = 0

Ha: mean(diff) < 0	Ha: mean(diff) != 0	Ha: mean(diff) > 0
t = -2.6685	t = -2.6685	t = -2.6685
P < t = 0.0041	P > t = 0.0082	P > t = 0.9959

3. OPTIONS = Increased Spending

Government spending If the government had to choose between the three options below, which do you think it should choose? Spend Less on the NHS; Keep spending on the same level; Spend more on the NHS.

*In Dataset: "tloptions" & "t2options"

Coding: 0 = spend less, 1 = spend more

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]
t2opti~s	212	.8915094	.0141882	.2065826	.8635407 .9194781
tlopti~s	212	.875	.0156369	.2276763	.8441755 .9058245
diff	212	.0165094	.0196045	.2854465	-.0221364 .0551553

Ho: mean(t2options - tloptions) = mean(diff) = 0

Ha: mean(diff) < 0	Ha: mean(diff) != 0	Ha: mean(diff) > 0
t = 0.8421	t = 0.8421	t = 0.8421
P < t = 0.7997	P > t = 0.4007	P > t = 0.2003

4. INCREASE CHARGES

Average of the following items:

CHGP = Charges for visiting your GP
One way of getting more money into the NHS is to charge people for certain things. How much would you be in favour of or against introducing charges for visiting your GP? (5-pt scale)
TTest (N=204): T1 = .225, T2 = .208; T2-T1 = -.017 (S.E. = 0.02); p = .349

CHGVIS = Charges for your GP visiting
One way of getting more money into the NHS is to charge people for certain things. How much would you be in favour of or against introducing charges for your GP visiting you at home? (5-pt scale)
TTest (N=205): T1 = .323, T2 = .299; T2-T1 = -.024 (S.E. = 0.02); p = .254

CHMEAL = Charges for hospital meals
One way of getting more money into the NHS is to charge people for certain things. How much would you be in favour of or against introducing charges for the cost of your hospital meals when you are an inpatient? (5-pt scale)
TTest (N=203): T1 = .379, T2 = .395; T2-T1 = .016 (S.E. = 0.02); p = .479

CHSTAY = Charges for accommodation in hospital
One way of getting more money into the NHS is to charge people for certain things. How much would you be in favour of or against introducing charges for the cost of accommodation while you are in hospital? (5-pt scale)
TTest (N=199): T1 = .245, T2 = .256; T2-T1 = .011 (S.E. = 0.02); p = .548

CHAMB = Charges for non-emergency ambulance

One way of getting more money into the NHS is to charge people for certain things. How much would you be in favour of or against introducing charges for using a non-emergency ambulance? (5-pt scale)

TTest (N=200): T1 = .503, T2 = .494; T2-T1 = -.009 (S.E. = 0.03); p = .731

*In Dataset : « t1hlthfundoptions » & « t2hlthfundoptions »

Coding: 0 = against, 1 = in favour

Cronbach's Alpha at T1: 0.802

Cronbach's Alpha at T2: 0.756

Factor Analysis

(iterated principal factors; 1 factor retained)

Factor	Eigenvalue	Difference	Proportion	Cumulative
1	2.28525	2.04693	1.0000	1.0000
2	0.23832	0.07094	0.1043	1.1043
3	0.16737	0.29705	0.0732	1.1775
4	-0.12968	0.14636	-0.0567	1.1208
5	-0.27604	.	-0.1208	1.0000

Factor Loadings		
Variable	1	Uniqueness
t2chgp	0.61784	0.61828
t2chvis	0.66711	0.55497
t2chmeal	0.76481	0.41507
t2chstay	0.82306	0.32257
t2chamb	0.44287	0.80387

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2hlth~s	217	.3455261	.0149433	.2201281	.3160728	.3749794
t1hlth~s	217	.3343318	.0147649	.2175004	.3052301	.3634335
diff	217	.0111943	.0159266	.2346131	-.0201971	.0425857

Ho: mean(t2hlthfundoptions - t1hlthfundoptions) = mean(diff) = 0

Ha: mean(diff) < 0

Ha: mean(diff) != 0

Ha: mean(diff) > 0

t = 0.7029

t = 0.7029

t = 0.7029

P < t = 0.7586

P > |t| = 0.4829

P > t = 0.2414

5. CUTTING EXPENSIVE TREATMENTS

Average of the following items:

CTHART = NHS cut down on heart transplants

Some people say the NHS should cut down on very expensive treatments or services and use the savings to provide less expensive treatments or services for more people. With this aim, how much would you be in favour of, or against, the NHS cutting down on heart transplants. (5-pt scale)

TTest (N=198): T1 = .740, T2 = .694; T2-T1 = -.045 (S.E. = 0.02); p = .014

CTNURS = NHS cut down on long term nursing care

Some people say the NHS should cut down on very expensive treatments or services and use the savings to provide less expensive treatments or services for more people. With this aim, how much would you be in favour of, or against, the NHS cutting down on long term nursing care of the elderly. (5-pt scale)

TTest (N=193): T1 = .699, T2 = .709; T2-T1 = .009 (S.E. = 0.02); p = .664

CTBABY = NHS cut down on intensive care for premature
Some people say the NHS should cut down on very expensive treatments or services and use the savings to provide less expensive treatments or services for more people. With this aim, how much would you be in favour of, or against, the NHS cutting down on intensive care for very premature babies whose survival is in doubt? (5-pt scale)

TTest (N=190): T1 = .709, T2 = .667; T2-T1 = -.042 (S.E. = .02); p = .078

Coding: 0 = make cuts, 1 = against cuts

and then average that with

TREATA = Lower priority to expensive treatments
Please say how much you agree or disagree with the following. If the NHS had to choose, it should give lower priority to expensive treatments so that more money is available to treat a larger number of sick people whose treatment costs less. (5-pt scale)

TTest (N=179): T1 = .557, T2 = .534; T2-T1 = -.024 (S.E. = 0.02); p = .264

In Dataset : « t1ctexptrtmentb » & « t2ctexptrtmentb »

Cronbach's Alpha at T1: 0.756

Cronbach's Alpha at T2: 0.713

Factor Analysis

(iterated principal factors; 1 factor retained)				
Factor	Eigenvalue	Difference	Proportion	Cumulative
1	1.62905	1.50636	1.0000	1.0000
2	0.12269	0.09749	0.0753	1.0753
3	0.02520	0.17310	0.0155	1.0908
4	-0.14790	.	-0.0908	1.0000

Factor Loadings		
Variable	1	Uniqueness
t2cthart	0.77225	0.40363
t2ctnurs	0.57396	0.67057
t2ctbaby	0.70775	0.49909
t2treata	0.44983	0.79765

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2ctex~b	217	.6044547	.0148076	.2181289	.5752689	.6336405
t1ctex~b	217	.6359447	.015329	.2258106	.6057311	.6661583
diff	217	-.03149	.0157562	.2321037	-.0625457	-.0004344

Ho: mean(t2ctexptrtmentb - t1ctexptrtmentb) = mean(diff) = 0

Ha: mean(diff) < 0

t = -1.9986

P < t = 0.0235

Ha: mean(diff) != 0

t = -1.9986

P > |t| = 0.0469

Ha: mean(diff) > 0

t = -1.9986

P > t = 0.9765

6. PRIVATIZING SOME TREATMENTS

Average of the following items:

CTFERT = NHS cut down on fertility treatment

Other people say the NHS should cut down on certain treatments or services that should be provided instead by private medicine or charities. With this aim, how much would you be in favour of, or against, the NHS cutting down on fertility treatment? (5-pt scale)

TTest (N=193): T1 = .475, T2 = .437; T2-T1 = -.039 (S.E. = 0.02); p = .090

CTCOSM

= NHS cut down on cosmetic surgery

Other people say the NHS should cut down on certain treatments or services that should be provided instead by private medicine or charities. With this aim, how much would you be in favour of, or against, the NHS cutting down on cosmetic surgery. (5-pt scale)

TTest (N=188): T1 = .392, T2 = .334; T2-T1 = -.059 (S.E. = 0.03); p = .035

Coding: 0 = favour cuts/privatization, 1 = against cuts/privatization

*In Dataset: "tlprtretrat" & "t2prtretrat"

Correlation at T1:

	tlctfert	tlctcosm
tlctfert	1.0000	
tlctcosm	0.3819	1.0000

Correlation at T2:

	t2ctfert	t2ctcosm
t2ctfert	1.0000	
t2ctcosm	0.5424	1.0000

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2prtretrat	207	.3846618	.0198264	.2852518	.3455732	.4237504
tlprtretrat	207	.4390097	.0169894	.2444344	.4055143	.472505
diff	207	-.0543478	.0223961	.3222237	-.0985028	-.0101929

Ho: mean(t2prtretrat - tlprtretrat) = mean(diff) = 0

Ha: mean(diff) < 0

t = -2.4267

P < t = 0.0080

Ha: mean(diff) != 0

t = -2.4267

P > |t| = 0.0161

Ha: mean(diff) > 0

t = -2.4267

P > t = 0.9920

Note: The following item is NOT included in this index. A footnote shall be made to indicate "cthosp" does not correlate well with the above two items.

CTHOSP

= NHS cut down on hospice care for terminally ill

Other people say the NHS should cut down on certain treatments or services that should be provided instead by private medicine or charities. With this aim, how much would you be in favour of, or against, the NHS cutting down on hospice care for the terminally ill. (5-pt scale)

TTest (N=191): T1 = .733, T2 = .709; T2-T1 = -.024 (S.E. = 0.02); p = .244

Correlation at T1:

	tlctfert	tlcthosp	tlctcosm
tlctfert	1.0000		
tlcthosp	0.0920	1.0000	
tlctcosm	0.3715	-0.0418	1.0000

Correlation at T2:

	t2ctfert	t2cthosp	t2ctcosm
t2ctfert	1.0000		
t2cthosp	0.1989	1.0000	
t2ctcosm	0.5331	0.0584	1.0000

Cronbach's Alpha at T1: 0.351

Cronbach's Alpha at T2: 0.514

Principal Components Analysis

(principal components; 3 components retained)

Component	Eigenvalue	Difference	Proportion	Cumulative
1	1.59001	0.62802	0.5300	0.5300
2	0.96199	0.51400	0.3207	0.8507
3	0.44799	.	0.1493	1.0000

Variable	1	2	3
t2ctfert	0.69327	-0.07796	-0.71645
t2cthosp	0.29862	0.93585	0.18713
t2ctcosm	0.65590	-0.34368	0.67208

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2priv~t	211	.485782	.0162667	.2362878	.453715	.517849
t1priv~t	211	.53594	.0141123	.2049933	.50812	.5637599
diff	211	-.050158	.0180293	.2618905	-.0856995	-.0146164

Ho: mean(t2privatetrt - t1privatetrt) = mean(diff) = 0

Ha: mean(diff) < 0

t = -2.7820

P < t = 0.0029

Ha: mean(diff) != 0

t = -2.7820

P > |t| = 0.0059

Ha: mean(diff) > 0

t = -2.7820

P > t = 0.9971

7. EQUAL PRIORITY REGARDLESS OF RISK

Coding: 1 = no difference, 0 = otherwise

PRSMOKB = Respondent's view on smoking and priority

Suppose two equally sick people in the same area go on a waiting list at the same time for the same heart operation. One is a heavy smoker, the other does not smoke. In your opinion which one should get the operation first?

TTest (N=199): T1 = .523, T2 = .598; T2-T1 = .075 (S.E. = 0.04); p = .035

AGEB = Respondent's age priority 1=30, .5=no 0= 0=70yr

Now, suppose two other equally sick people in the same area go on a waiting list for the same heart operation. In this case one is aged 70, the other 30. In your opinion which one should get the operation first?

TTest (N=196): T1 = .571, T2 = .679; T2-T1 = .107 (S.E. = 0.04); p = .006

*In Dataset : « t1wl » & « t2wl »

Correlation at T1:

	t1smki~w	tlageb
t1smkingview	1.0000	

```

t1ageb |    0.4006    1.0000
Correlation at T2:
      | t2smki~w    t2ageb
-----+-----
t2smkingview |    1.0000
t2ageb |    0.2172    1.0000

```

Paired t test

```

-----+-----
Variable |      Obs      Mean    Std. Err.    Std. Dev.    [95% Conf. Interval]
-----+-----
t2wl |      224    .6517857    .0254442    .3808139    .6016439    .7019276
tlwl |      224    .5446429    .0288422    .4316699    .4878048    .6014809
-----+-----
diff |      224    .1071429    .028972    .4336133    .0500489    .1642368
-----+-----

```

Ho: mean(t2wl - tlwl) = mean(diff) = 0

```

Ha: mean(diff) < 0      Ha: mean(diff) != 0      Ha: mean(diff) > 0
t = 3.6982              t = 3.6982              t = 3.6982
P < t = 0.9999          P > |t| = 0.0003          P > t = 0.0001

```

Note: We tried the following coding, but decided to go with the above coding scheme.

Coding: 1=non-smoker/30 yr old, .5= no difference, 0=smoker/70 yr old

Correlation at T1:

```

      | t1smkb~i t1ageb~i
-----+-----
t1smkbtri |    1.0000
t1agebtri |    0.3534    1.0000

```

Correlation at T2:

```

      | t2smkb~i t2ageb~i
-----+-----
t2smkbtri |    1.0000
t2agebtri |    0.2071    1.0000

```

Paired t test

```

-----+-----
Variable |      Obs      Mean    Std. Err.    Std. Dev.    [95% Conf. Interval]
-----+-----
t2lowr~k |      222    .6441441    .0137401    .2047224    .6170658    .6712225
t1lowr~k |      222    .6677928    .0165494    .2465804    .635178    .7004076
-----+-----
diff |      222   -.0236486    .0167033    .2488739   -.0565668    .0092695
-----+-----

```

Ho: mean(t2lowrisk - t1lowrisk) = mean(diff) = 0

```

Ha: mean(diff) < 0      Ha: mean(diff) != 0      Ha: mean(diff) > 0
t = -1.4158              t = -1.4158              t = -1.4158
P < t = 0.0791          P > |t| = 0.1582          P > t = 0.9209

```

8. PRIORITY TO SEVERITY VS PLACE IN QUEUE:

LISTA - SEVERA (Subtract SEVERA from LISTA):

LISTA = Priority to waiting list

Please say how much you agree or disagree with the following. If the NHS had to choose, a person's place in the waiting list for a particular treatment should be the only basis for choosing between patients. (5-pt scale)

TTest (N=206): T1 = .495, T2 = .455; T2-T1 = -.040 (S.E. = 0.02); p = .091

SEVERA

= Priority to severity

Please say how much you agree or disagree with the following. If the NHS had to choose, the seriousness of their condition should be the only basis for choosing between patients. (5-pt scale)

TTest (N=193): T1 = .655, T2 = .705; T2-T1 = .049 (S.E. = 0.03); p = .052

Coding: 1 =strongly agree

*In Dataset: "tlseverity" & "t2severity"

Correlation at T1:

	tl1lista	tlsevera
tl1lista	1.0000	
tlsevera	0.2833	1.0000

Correlation at T2:

	t21lista	t2severa
t21lista	1.0000	
t2severa	0.2672	1.0000

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2seve~y	183	-.2486339	.0268786	.3636065	-.3016675	-.1956002
tlseve~y	183	-.1653005	.0260993	.3530647	-.2167967	-.1138044
diff	183	-.0833333	.0333491	.4511382	-.1491339	-.0175328

Ho: mean(t2severity - tlseverity) = mean(diff) = 0

Ha: mean(diff) < 0

Ha: mean(diff) != 0

Ha: mean(diff) > 0

t = -2.4988

t = -2.4988

t = -2.4988

P < t = 0.0067

P > |t| = 0.0133

P > t = 0.9933

9. PREVA

= Priority to prevention

Please say how much you agree or disagree with the following. If the NHS had to choose, it should give higher priority to preventing illness than to treating the very ill. (5-pt scale)

Coding: 1 = strongly agree

*In Dataset: "t1prevention" & "t2prevention"

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2prev~n	201	.5858209	.0213547	.3027548	.5437117	.6279301
tlprev~n	201	.6268657	.0213883	.303232	.5846901	.6690412
diff	201	-.0410448	.0252962	.3586355	-.0909262	.0088367

Ho: mean(t2prevention - t1prevention) = mean(diff) = 0

Ha: mean(diff) < 0

Ha: mean(diff) != 0

Ha: mean(diff) > 0

t = -1.6226	t = -1.6226	t = -1.6226
P < t = 0.0531	P > t = 0.1063	P > t = 0.9469

10. LIFEA Priority To Quality Of Life Over Danger

= Quality of life

Please say how much you agree or disagree with the following. If the NHS had to choose, it should give higher priority to those whose quality of life will benefit most from treatment than to those whose life is most in danger. (5-pt scale)

Coding: 1 = strongly agree

*In Dataset: "tlqoflife" & "t2qoflife"

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2qoflife	188	.5292553	.0227617	.3120923	.4843527	.574158
tlqoflife	188	.474734	.022354	.3065025	.4306356	.5188325
diff	188	.0545213	.02523	.3459364	.0047493	.1042933

Ho: mean(t2qoflife - tlqoflife) = mean(diff) = 0

Ha: mean(diff) < 0	Ha: mean(diff) != 0	Ha: mean(diff) > 0
t = 2.1610	t = 2.1610	t = 2.1610
P < t = 0.9840	P > t = 0.0320	P > t = 0.0160

11. DISCRETION: PUBLIC & GOV'T

Average of the following two items:

Coding: 1 = most say

INPUBA = Amount of general public input

Suppose that the NHS did have to make these sorts of choices about who to treat, or the order in which they should be treated. (None of the say; Some of the say; All or most of the say). Please tick one box on each line below to show how much say you think each of these groups should have.
The general public?

TTest (N=168): T1 = .750, T2 = .664; T2-T1 = -.086 (S.E. = 0.02); p = .000

INGOVA = Amount of government input

The government?

TTest (N=175): T1 = .783, T2 = .731; T2-T1 = -.051 (S.E. = 0.02); p = .021

In Dataset : « tldispubgvt » & « t2dispubgvt »

Correlation at T1:

	tlinpuba	tlingova
tlinpuba	1.0000	
tlingova	0.2001	1.0000

Correlation at T2:

	t2inpuba	t2ingova
t2inpuba	1.0000	
t2ingova	0.2836	1.0000

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2avg2	191	.7028796	.0145673	.2013235	.6741453	.7316139

t1avg2		191	.7709424	.0146524	.2025006	.7420401	.7998447
-----+							
diff		191	-.0680628	.0171029	.2363671	-.1017988	-.0343268

Ho: mean(t2avg2 - t1avg2) = mean(diff) = 0

Ha: mean(diff) < 0	Ha: mean(diff) != 0	Ha: mean(diff) > 0
t = -3.9796	t = -3.9796	t = -3.9796
P < t = 0.0000	P > t = 0.0001	P > t = 1.0000

12. DISCRETION: DOCTORS

Average of the following two items:

Coding: 1 = most say

INGPA = Amount of GP input
Suppose that the NHS did have to make these sorts of choices about who to treat, or the order in which they should be treated. (None of the say; Some of the say; All or most of the say). Please tick one box on each line below to show how much say you think each of these groups should have. GPs?
TTest (N=194): T1 = .722, T2 = .740; T2-T1 = .018 (S.E. = 0.02); p = .294

INDOCA = Amount of doctor input
Hospital doctors?
TTest (N=197): T1 = .817, T2 = .812; T2-T1 = -.005 (S.E. = 0.02); p = .814

In Dataset : « t1avgdisdoc » & « t2avgdisdoc »

Correlation at T1 :

		t1ingpa	t1indoca
-----+			
t1ingpa		1.0000	
t1indoca		0.3146	1.0000

Correlation at T2 :

		t2ingpa	t2indoca
-----+			
t2ingpa		1.0000	
t2indoca		0.2991	1.0000

Paired t test

Variable		Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]
-----+						
t2avgd~c		206	.7791262	.0140407	.2015216	.7514436 .8068089
t1avgd~c		206	.7682039	.0139722	.2005387	.7406563 .7957515
-----+						
diff		206	.0109223	.0161671	.2320421	-.0209529 .0427975

Ho: mean(t2avgdisdoc - t1avgdisdoc) = mean(diff) = 0

Ha: mean(diff) < 0	Ha: mean(diff) != 0	Ha: mean(diff) > 0
t = 0.6756	t = 0.6756	t = 0.6756
P < t = 0.7500	P > t = 0.5001	P > t = 0.2500

13. SAY

= Patients have more say
How much do you agree or disagree with each of the following statements? Patients should have more of a say in the decisions affecting their medical treatment. (5-pt scale)

Coding: 1 = strongly agree

In Dataset: "t1moresay" & "t2moresay"

Paired t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
t2more~y	212	.7582547	.0111368	.1621545	.736301	.7802084
t1more~y	212	.7169811	.0134929	.19646	.6903829	.7435793
diff	212	.0412736	.0133325	.1941241	.0149916	.0675555

Ho: mean(t2moresay - t1moresay) = mean(diff) = 0

Ha: mean(diff) < 0

t = 3.0957

P < t = 0.9989

Ha: mean(diff) != 0

t = 3.0957

P > |t| = 0.0022

Ha: mean(diff) > 0

t = 3.0957

P > t = 0.0011