Linear programming and duality

2105623 Optimization of Chemical Processes

26 October 2026 · 180-minute class

Geometry, Simplex, optimality certificates, and resource values



Plan for the three-hour class

Minutes	Topic
0–23	LP geometry and corner solutions
23–56	Basic feasible solutions and Simplex
56–66	Break
66–103	Duality and complementary slackness
103–163	Sensitivity, solver output, capacity decisions
163–180	Exit ticket, solutions, and connected recap

What makes a model linear?

$\max c^T x$ subject to $Ax \leq b$, $x \geq 0$

Coefficients are fixed; decision variables are continuous.

$x + 2y$ is linear. xy and x^2 are not.

A binary restriction changes an LP into a MILP.

Running example: two production routes

Item	Route x	Route y	Daily limit
Resource 1	1 unit/t	1 unit/t	4 units/day
Resource 2	2 units/t	1 unit/t	5 units/day
Contribution	3 × $100/t	2 × $100/t	Maximize

x, y = tonnes/day; contribution = $100/day units.

Choose continuous production; no fixed startup cost.

The production LP

$$\max z = 3x + 2y$$

$$x + y \leq 4$$

$$2x + y \leq 5$$

$$x \geq 0,\ y \geq 0$$



Objective contours and the best feasible point

$3x + 2y = z$ gives parallel lines.

Move the line toward larger z.

The last contact is $(x^*, y^*) = (1, 3)$.

$z^* = 9$, or \$900/day.



Why compare extreme points?

Vertex (x, y)	z = 3x + 2y
(0, 0)	0
(2.5, 0)	7.5
(1, 3)	9
(0, 4)	8

On a nonempty bounded polytope, a linear objective has an optimal extreme point.

A convex combination cannot beat all vertices used in that combination.

An LP can have many optimal solutions

Change only the objective to max x + y.

Every point from (0,4) to (1,3) is optimal.

The entire optimal edge has value 4.



Infeasible and unbounded are different

Infeasible: $x \le 1$ and $x \ge 2$ cannot both hold.

Unbounded: max x with only $x \ge 0$ has no finite maximum.



Quick check: feasibility before profit

Original LP: max $3x+2y$; $x+y\le4$, $2x+y\le5$, $x,y\ge0$.

A: (2,2) gives $z=10$. B: (1,3) gives $z=9$.

Which point can be accepted? Explain the failed constraint.

Slack variables and standard form

$x + y + s_1 = 4$

$2x + y + s_2 = 5$

$x, y, s_1, s_2 \geq 0$

s_1 and s_2 measure unused resources.

Shown: $(x,y)=(2.5,0)$, so $(s_1,s_2)=(1.5,0)$.



A basis selects the variables we solve for

Basis	Nonbasic variables	Basic solution
$\{s_1, s_2\}$	$x=y=0$	$s_1=4,\ s_2=5$
$\{x, s_1\}$	$y=s_2=0$	$x=2.5,\ s_1=1.5$
$\{x, y\}$	$s_1=s_2=0$	$x=1,\ y=3$

Choose two independent columns as a basis for the two equations.

Set the other two variables to zero, then solve.

A basic solution is feasible only if every variable is nonnegative.

The Simplex decision at each step

1. Choose an improving nonbasic variable.

2. Increase it until a basic variable reaches zero.

3. Pivot: exchange the entering and leaving variables.

Positive reduced cost improves z if a positive feasible step exists.



Pivot 1: x enters, s_2 leaves

At the origin: $z = 3x + 2y$. Choose x.

With $y=0$: $s_1=4-x$; $s_2=5-2x$.

Ratio test: $4/1=4$; $5/2=2.5$.

Move to (2.5,0); $z=7.5$.



Dictionary after the first pivot

$x = 2.5 - 0.5y - 0.5s_2$

$s_1 = 1.5 - 0.5y + 0.5s_2$

$z = 7.5 + 0.5y - 1.5s_2$

Nonbasic: $y = s_2 = 0$. The +0.5 coefficient means y can improve z.

Pivot 2: y enters, s_1 leaves

Hold $s_2=0$ and increase y.

$x \geq 0$ gives $y \leq 2.5/0.5=5$.

$s_1 \geq 0$ gives $y \leq 1.5/0.5=3$.

Thus $y=3$, $x=1$, $s_1=s_2=0$; $z=9$.



The final dictionary certifies optimality

$$x = 1 + s_1 - s_2$$

$$y = 3 - 2s_1 + s_2$$

$$z = 9 - s_1 - s_2$$

Because $s_1, s_2 \geq 0$, every feasible point satisfies $z \leq 9$.

Degeneracy and starting feasibility

A degenerate BFS has at least one zero basic variable.

A pivot may change the basis without improving the point.

At b=(4,8), the basis {x,y} gives x=4, y=0.

If a feasible slack basis is unavailable, use Phase I or another initialization.



Practice: change the contribution

Keep $x+y\le 4$, $2x+y\le 5$, $x,y\ge 0$.

Change the objective to max $z=x+3y$.

Choose the first entering variable and apply the ratio test.

Predict the optimum and write a stopping argument.

Practice solution: one pivot is enough

y enters; min(4/1,5/1)=4, so s_1 leaves.

$y=4-x-s_1$; $s_2=1-x+s_1$.

$z=x+3y=12-2x-3s_1\leq 12$.

Optimal: (x,y)=(0,4), slack=(0,1), $z^*=12$.



Break

10 minutes

After the break: what is one more unit of a resource worth?

The dual prices the resources

Return to max z=3x+2y, b=(4,5).

u, v price resources 1 and 2 ($100/unit).

A tonne of x uses resources worth u+2v.

A tonne of y uses resources worth u+v.

Cover contributions: u+2v≥3, u+v≥2.



Primal and dual side by side

Primal: max $c^T x$, subject to $Ax \leq b$, $x \geq 0$.

Dual: min $b^T u$, subject to $A^T u \geq c$, $u \geq 0$.

Here: min $w = 4u + 5v$

subject to $u + 2v \geq 3$, $u + v \geq 2$, $u, v \geq 0$.

Signs depend on the formulation

For a maximization primal:

Primal feature	Dual counterpart
Constraint $a_i^T x \le b_i$	$u_i \ge 0$
Constraint $a_i^T x \ge b_i$	$u_i \le 0$
Constraint $a_i^T x = b_i$	u_i is free
Variable $x_j \ge 0$	$(A^T u)_j \ge c_j$
Variable $x_j \le 0$	$(A^T u)_j \le c_j$
Variable x_j is free	$(A^T u)_j = c_j$

Weak duality gives an upper bound

For any primal-feasible x and dual-feasible u:

$c^T x \le x^T A^T u = u^T A x \le u^T b$.

Example: (u,v)=(2,1) is feasible; w=13.

Therefore every feasible production plan has z≤13.

Dual geometry: tighten the upper bound

The dual feasible set lies above both price lines.

Minimize 4u+5v over that set.

Candidate corner prices: (3,0), (1,1), (0,2).

Their bounds are 12, 9, and 10.



Solve the dual and compare

$u+2v=3$ and $u+v=2 \rightarrow u=v=1$.

Dual value: $4(1)+5(1)=9$.

Primal point (1,3) is feasible and gives 9.

Feasible values meet: both solutions are optimal.



Strong duality for linear programs

If an LP is feasible with a finite optimum, its dual also has an optimum with the same value.

A primal unbounded above implies the dual is infeasible.

Primal infeasible does not by itself imply dual unbounded.

The duality gap separates two kinds of waste

$s = b - Ax \geq 0$; $r = A^T u - c \geq 0$.

$w - z = u^T s + x^T r \geq 0$.

Unused priced resources contribute $u^T s$.

Producing below the assigned resource cost contributes $x^T r$.

Complementary slackness

For primal-feasible x and dual-feasible u:

$u_i(b_i - a_i^T x) = 0$ for each resource.

$x_j((A^T u)_j - c_j) = 0$ for each product.

Unused resource implies zero price. Positive output implies a tight dual constraint.

A binding resource can have zero price.

LP duality is KKT in a special form

Rewrite $\max c^T x$ as $\min -c^T x$.

$L = -c^T x + u^T(Ax - b) - r^T x$.

Stationarity: $-c + A^T u - r = 0$.

Primal feasibility, $u, r \geq 0$, and complementarity complete the optimality conditions.

Practice: a dual certificate

For max x+3y with the same two resource constraints:

1. Write the dual.

2. Test primal (0,4) and dual (u,v)=(3,0).

3. Check both objectives and all complementarity products.

Practice solution: zero price for unused capacity

Dual: min $4u+5v$; $u+2v\geq 1$, $u+v\geq 3$; $u,v\geq 0$.

At $(3,0)$: $w=12=z(0,4)$.

Resource slack: $(0,1)$. Dual slack: $(2,0)$.

Products: $3\cdot 0=0$, $0\cdot 1=0$, $0\cdot 2=0$, $4\cdot 0=0$.

Shadow prices: value per extra unit

Return to c=(3,2), b=(4,5): u*=v*=1.

Hold A and c fixed; keep the same basis feasible and optimal.

Then $\Delta z^* = u^*\Delta b_1 + v^*\Delta b_2$ exactly.

Each additional resource unit is worth $100 within this region.

1. Resource 1 changes, resource 2 stays fixed

Maximize $z = 3x + 2y$.

$x + y \le b_1$, $2x + y \le 5$, $x,y \ge 0$.

At $b_1 = 4$: $(x,y) = (1,3)$, $z^* = 9$.

How far can we change b_1 and keep its shadow price equal to 1?



2. The intersection moves with b_1

Suppose both resources remain fully used.

$x + y = b_1$

$2x + y = 5$

Subtract the equations: $x = 5 - b_1$.

Substitute back: $y = 2b_1 - 5$.



3. Nonnegativity gives the allowable range

$x = 5 - b_1 \geq 0$ requires $b_1 \leq 5$.

$y = 2b_1 - 5 \geq 0$ requires $b_1 \geq 2.5$.

Together: $2.5 \leq b_1 \leq 5$.

At 2.5: (x,y) = (2.5,0).

At 5: (x,y) = (0,5).



4. Inside the range: $b_1 = 4.5$

$x = 5 - 4.5 = 0.5$.

$y = 2(4.5) - 5 = 4$.

$z^* = 3(0.5) + 2(4) = 9.5$.

From $b_1 = 4$: $\Delta b_1 = 0.5$, $\Delta z^* = 0.5$.

The extra contribution is \$50.



5. Duality confirms the shadow price

At the intersection: $z^* = 3(5-b_1) + 2(2b_1-5) = b_1 + 5$.

Dual prices (u,v) = (1,1) remain feasible.

Dual value: $w = b_1u + 5v = b_1 + 5$.

Primal value = dual value, so both are optimal.

Within the range: $\Delta z^* = 1 \times \Delta b_1$.



6. Below the range: $b_1 = 2$

The old intersection is $(3,-1)$, which is infeasible.

The optimum is $(x,y) = (2,0)$, $z^* = 6$.

Resource 2 has slack: $5 - 2(2) = 1$.

For $0 < b_1 < 2.5$: $z^* = 3b_1$.

Shadow price of resource 1: 3.



7. Above the range: $b_1 = 6$

The old intersection is $(-1,7)$, which is infeasible.

The optimum is $(x,y) = (0,5)$, $z^* = 10$.

Resource 1 has slack: $6 - 5 = 1$.

For $b_1 > 5$: $z^* = 10$.

Shadow price of resource 1: 0.



8. The value function changes slope

$0 \le b_1 \le 2.5$: $z^* = 3b_1$.

$2.5 \le b_1 \le 5$: $z^* = b_1 + 5$.

$b_1 \ge 5$: $z^* = 10$.

The middle formula includes both endpoints.

At 2.5 and 5, left and right slopes differ.



1. Resource 2 changes, resource 1 stays fixed

Maximize $z = 3x + 2y$.

$x + y \le 4$, $2x + y \le b_2$, $x,y \ge 0$.

At $b_2 = 5$: $(x,y) = (1,3)$, $z^* = 9$.

How far can b_2 change while its shadow price stays at 1?



2. The intersection as a function of b_2

Suppose both resources remain fully used.

$x + y = 4$

$2x + y = b_2$

Subtract the equations: $x = b_2 - 4$.

Substitute back: $y = 8 - b_2$.



3. Nonnegativity gives $4 \le b_2 \le 8$

$x = b_2 - 4 \ge 0$ requires $b_2 \ge 4$.

$y = 8 - b_2 \ge 0$ requires $b_2 \le 8$.

Together: $4 \le b_2 \le 8$.

At 4: (x,y) = (0,4). At 8: (x,y) = (4,0).

From $b_2 = 5$: decrease by 1 or increase by 3.



4. Inside the range: $b_2 = 6$

$x = 6 - 4 = 2.$

$y = 8 - 6 = 2.$

$z^* = 3(2) + 2(2) = 10.$

From $b_2 = 5$: $\Delta b_2 = 1$, $\Delta z^* = 1$.

The extra contribution is \$100.



5. Duality confirms the price of resource 2

At the intersection: $z^* = 3(b_2-4) + 2(8-b_2) = b_2 + 4$.

Dual prices $(u,v) = (1,1)$ remain feasible.

Dual value: $w = 4u + b_2v = b_2 + 4$.

Primal value = dual value, so both are optimal.

Within the range: $\Delta z^* = 1 \times \Delta b_2$.



6. Below the range: $b_2 = 3$

The intersection formula gives $(-1,5)$, which is infeasible.

The optimum is $(x,y) = (0,3)$, $z^* = 6$.

Resource 1 has slack: $4 - 3 = 1$.

For $0 < b_2 < 4$: $z^* = 2b_2$.

Shadow price of resource 2: 2.



7. Above the range: $b_2 = 9$

The intersection formula gives (5,−1), which is infeasible.

The optimum is (x,y) = (4,0), z* = 12.

Resource 2 has slack: 9 − 2(4) = 1.

For $b_2 > 8$: z* = 12.

Shadow price of resource 2: 0.



8. The value function for resource 2

$0 \le b_2 \le 4$: $z^* = 2b_2$.

$4 \le b_2 \le 8$: $z^* = b_2 + 4$.

$b_2 \ge 8$: $z^* = 12$.

At 4: left slope 2, right slope 1.

At 8: left slope 1, right slope 0.



9. Changing both resources needs a joint check

With both RHS values changing:

$x = b_2 - b_1$, $y = 2b_1 - b_2$.

The same basis requires $b_1 \le b_2 \le 2b_1$.

$(b_1,b_2) = (5,4)$ is outside this region.

Its true optimum is $(x,y)=(0,4)$, $z^*=8$.



1. Objective sensitivity keeps the feasible set fixed

Maximize $z = c_1x + 2y$.

$x + y \le 4$, $2x + y \le 5$, $x,y \ge 0$.

Baseline $c_1 = 3$: $(x,y) = (1,3)$, $z^* = 9$.

Only the contribution from product x changes.

Which c_1 values keep (1,3) optimal?



2. Complementary slackness determines the prices

At (1,3), both products have positive output.

Their dual constraints must be tight:

$u + 2v = c_1$, $u + v = c_2$.

Subtract: $v = c_1 - c_2$.

Then $u = 2c_2 - c_1$.



3. Nonnegative dual prices give the range

$u = 2c_2 - c_1 \geq 0$ requires $c_1 \leq 2c_2$.

$v = c_1 - c_2 \geq 0$ requires $c_1 \geq c_2$.

Together: $c_2 \leq c_1 \leq 2c_2$.

With $c_2 = 2$: $2 \leq c_1 \leq 4$.

The contour slope $-c_1/2$ lies between -2 and -1.



4. Below the range: $c_1 = 1$

Maximize $z = x + 2y$.

At (1,3): $z = 7$.

The optimum is (0,4), with $z^* = 8$.

The old price formula gives $v = -1$.

For $c_1 < 2$: (0,4) stays optimal.



5. At $c_1 = 2$, an entire edge is optimal

$z = 2x + 2y = 2(x+y) \le 8$.

Every feasible point on x+y=4 attains 8.

The optimal edge runs from (0,4) to (1,3).

(1,3) is still optimal, but not unique.

Dual prices: (u,v) = (2,0).



6. Inside the range: $c_1 = 3.5$

The optimum stays at (x,y) = (1,3).

$z^* = 3.5(1) + 2(3) = 9.5$.

$u = 4 - 3.5 = 0.5$, $v = 3.5 - 2 = 1.5$.

Dual value: $4u + 5v = 9.5$.

The production plan stays fixed, but resource prices change.



7. At $c_1 = 4$, the other edge is optimal

$z = 4x + 2y = 2(2x+y) \le 10.$

Every feasible point on 2x+y=5 attains 10.

The optimal edge runs from (1,3) to (2.5,0).

(1,3) remains one of the optimal solutions.

Dual prices: (u,v) = (0,2).



8. Above the range: $c_1 = 5$

Maximize $z = 5x + 2y$.

At (1,3): $z = 11$.

The optimum is (2.5,0), with $z^* = 12.5$.

The old price formula gives $u = -1$.

For $c_1 > 4$: (2.5,0) stays optimal.



9. Optimal value as the coefficient changes

$c_1 < 2$: $z^* = 8$, with unique $x^* = 0$.

$2 \le c_1 \le 4$: $z^* = c_1 + 6$.

$c_1 > 4$: $z^* = 2.5c_1$, with unique $x^* = 2.5$.

On open intervals, $dz^*/dc_1 = x^*$.

At 2: $x^* \in [0,1]$; at 4: $x^* \in [1,2.5]$.



1. Return to the original production model

Maximize $z = 3x + 2y$.

$x + y \le 4$; $2x + y \le 5$; $x, y \ge 0$.

Baseline: $(x,y) = (1,3)$, $z^* = 9$.

Resource prices: $u = 1$, $v = 1$.

Could a new product improve this plan?



2. Price the resources used by the new product

New product t: contribution $c_t = 4$.

Each unit uses resource 1: 1; resource 2: 2.

At the old prices, resource value = 1u + 2v = 3.

Contribution 4 exceeds resource value 3 by 1.



3. Derive reduced cost from a feasible exchange

Keep both resources fully used:

$x + y + t = 4$; $2x + y + 2t = 5$.

Subtract: $x + t = 1$, so $x = 1 - t$ and $y = 3$.

$z = 3(1-t) + 2(3) + 4t = 9 + t$.

Thus $\bar{c}_t = c_t - (u+2v) = 4 - 3 = 1$.



4. The feasible step ends when x reaches zero

$x = 1 - t \geq 0$ and $t \geq 0$ require $0 \leq t \leq 1$.

At $t = 1$: $(x,y,t) = (0,3,1)$, $z = 10$.

At $t = 1.5$, the same rule gives $x = -0.5$.

So $z = 9 + t$ cannot be extrapolated past $t = 1$.

The basis must change when x leaves.



5. The new product invalidates the old certificate

Max 3x + 2y + 4t; x,y,t ≥ 0.

x+y+t ≤ 4; 2x+y+2t ≤ 5.

Dual: u,v ≥ 0; u+2v ≥ 3; u+v ≥ 2; u+2v ≥ 4.

Old prices (1,1) fail: 3 < 4 for t.

So the old upper bound 9 is invalid.



6. Reoptimize and certify the new maximum

Choose new dual prices (u,v) = (0,2).

Resource values: x → 4 ≥ 3; y → 2; t → 4.

Every feasible plan has z ≤ 4u + 5v = 10.

Plan (0,3,1) attains 10, so it is optimal.

New reduced costs (x,y,t) = (−1,0,0).

7. More than one new production plan is optimal

With x = 0: y+t ≤ 4 and y+2t ≤ 5.

On y+2t = 5, profit 2y+4t = 10.

Feasibility gives 1 ≤ t ≤ 2.5 and y = 5−2t.

Examples: (0,3,1), (0,2,1.5), (0,0,2.5).

Beyond t=1, reallocate y; profit stays at 10.



8. Compare three possible product contributions

Keep the new column fixed at (1,2).

At old prices: $\bar{c}_t = c_t - 3$.

$c_t = 2$: $\bar{c}_t = -1$; optimal t = 0, z* = 9.

$c_t = 3$: $\bar{c}_t = 0$; feasible exchange keeps z* = 9.

$c_t = 4$: $\bar{c}_t = +1$; reoptimize to z* = 10.



9. A reduced-cost decision checklist

For max: $\bar{c}_j = c_j - a_j^T \pi$, where $\pi = (u,v)$.

1. Price the column at the current dual prices.

2. Check a feasible step and its limit.

3. Reoptimize; check a new dual certificate.

For nonbasic variables at lower bounds: max $\bar{c} \le 0$; min $\bar{c} \ge 0$ at an optimal basis.

From screening to a verified new optimum



1. Send a minimization model to SciPy

Return to the original LP; no new product t.

Max z = 3x+2y becomes min F = −3x−2y.

c = [−3, −2]; A = [[1,1], [2,1]].

b = [4,5]; default bounds: (0,None).

linprog(c, A_ub=A, b_ub=b, method="highs")

Same feasible set; opposite objective signs



2. Read the actual solver output

Output	Expected value	Meaning
success	True	Optimal termination
x	[1, 3]	Production
fun	−9	Minimized negative contribution
ineqlin.residual	[0, 0]	Unused capacity
ineqlin.marginals	[−1, −1]	Sensitivity of the minimized objective

Return to the original two-product LP. SciPy minimizes $-3x-2y$; max-profit shadow prices = −marginals = (1,1).

Check status, feasibility, objective, and a small RHS change.

3. Derive the sign instead of memorizing it

Where F* is differentiable: $m_i = \partial F^*/\partial b_i$.

Since $z^*(b) = -F^*(b)$: $\partial z^*/\partial b_i = -m_i$.

Here $m = (-1,-1)$, so $(u,v) = (1,1)$.

Extra capacity lowers F* and raises profit z*.

Units: $100 of profit per extra capacity unit.



4. Verify resource 1 with a small perturbation

Change only b_1: 4 → 4.1; keep $b_2 = 5$.

Prediction: $\Delta F^* = (-1)(0.1) = -0.1$.

Rerun: (x,y) = (0.9,3.2), $F^* = -9.1$.

Profit: 9 → 9.1, a gain of \$10/day.

(−9.1+9)/0.1 = −1; (9.1−9)/0.1 = +1.



5. Verify resource 2 independently

Reset $b_1 = 4$; change only b_2: $5 \to 5.1$.

Rerun: (x,y) = (1.1,2.9), F* = −9.1.

Resource use: 1.1+2.9=4; 2(1.1)+2.9=5.1.

ΔF*/0.1 = −1; Δz*/0.1 = +1.

Rows of ineqlin.marginals follow rows of A_ub.



6. A marginal is not a global prediction

Old prices apply for $2.5 \le b_1 \le 5$ ($b_2=5$).

From $b_1=4$ to 6, the old rule predicts $z^*=11$.

Rerun at $b_1=6$: $(x,y)=(0,5)$, $z^*=10$.

Residuals = (1,0); marginals = (0,−2).

New profit shadow prices are (0,2).



7. Row marginals and bound marginals differ

New example: max z=x+3y, with the same constraints.

Solution: (0,4), z*=12; row marginals=(−3,0).

lower.marginals = (2,0): this refers to variable bounds.

Raise the lower bound of x: 0 → 0.1.

Then (x,y)=(0.1,3.9), F*: −12 → −11.8.



8. Connect the bound marginal to reduced cost

For max z=x+3y, resource prices are (u,v)=(3,0).

Product x has max reduced cost: 1−(3+2·0)=−2.

Its min lower-bound marginal is +2.

Forcing x≥0.1 loses 0.2 profit units = $20/day.

Read status → objective → field/RHS → sign → local check.

1. Define the baseline and the purchase offer

Return to max $z=3x+2y$; capacities $(b_1,b_2)=(4,5)$.

Baseline: $(x,y)=(1,3)$, $z^*=9$ = \$900/day.

Let $q \geq 0$ be extra resource 1 capacity per day.

Offer A: $q=0.5$ costs \$30/day in total.

Compare incremental contribution with this new cost.



2. Convert the shadow-price range into a purchase limit

With $b_2=5$, the old basis needs $2.5 \le b_1 \le 5$.

Since $b_1=4+q$ and $q \ge 0$, its purchase range is $0 \le q \le 1$.

Old shadow price $u=1$ means \$100 per capacity unit.

For $0 \le q \le 1$: extra contribution $G(q)=\$100q$/day.

For $q>1$, reoptimize before valuing the purchase.



3. Add 0.5 capacity and calculate the gross gain

q=0.5 gives $b_1=4.5$, inside the old-price range.

$x=5-b_1=0.5$; $y=2b_1-5=4$.

$z^*=3(0.5)+2(4)=9.5$ = \$950/day.

Gross incremental contribution: \$950−\$900=\$50/day.

Shadow-price check: \$100×0.5=\$50/day.



4. Subtract the purchase cost exactly once

Gross gain G=$50/day; purchase cost C=$30/day.

Net benefit N=G−C=$20/day > 0: buy Offer A.

Contribution after the new cost: $950−$30=$920/day.

Compare $920 with the baseline $900.

Do not subtract existing variable costs again.



5. Buying 2 units crosses the useful range

q=2 gives $b_1=6$; the old rule predicts a $200 gain.

Reoptimize: (x,y)=(0,5), z*=10 = $1,000/day.

Actual gain: $1,000−$900=$100/day.

Resource 1 has slack 1; resource 2 is fully used.

G(q)=$100q for 0≤q≤1; G(q)=$100 for q≥1.



6. Use a break-even price for the whole offer

For q=2, maximum useful payment is $100/day.

If total cost C=$150/day: N=$100−$150=−$50/day.

Reject: $1,000−$150=$850 < $900 baseline.

If C=$100/day: N=0, so the model is indifferent.

If C<$100/day: N>0; buying improves the baseline.



7. If capacity is divisible, choose how much to buy

New pricing assumption:
C(q)=$60q/day; no fixed fee.

For $0 \le q \le 1$: $N(q)=100q-60q=40q$.

For $q \ge 1$: $N(q)=100-60q$.

Best purchase: $q^*=1$; net benefit=$40/day.

The unit price is attractive only while capacity adds value.



8. Compare complete offers by net benefit

Choose at most one package; each cost is a total daily cost.

A: q=0.5, C=$30 → G=$50, N=$20/day.

B: q=1, C=$70 → G=$100, N=$30/day.

C: q=2, C=$90 → G=$100, N=$10/day.

Choose B: $900+$30=$930/day after the new cost.



Exit ticket 1: can an entire edge be optimal?

Work independently. Attempt all four questions before the solutions.

For this question only: max z=2x+2y.

x+y≤4; 2x+y≤5; x,y≥0.

Identify the optimal edge and its endpoints.

Explain why every point on it gives the same best value.



Exit ticket 2: is this a valid dual bound?

Return to the original objective: max z=3x+2y.

Dual: min w=4u+5v; u,v≥0.

Constraints: u+2v≥3 and u+v≥2.

Test the candidate (u,v)=(1,0).

Compute w. Can it bound every feasible primal value? Why?



Exit ticket 3: construct an optimality certificate

Original LP: max $z=3x+2y$.

$x+y\le 4$; $2x+y\le 5$; $x,y\ge 0$.

Find a feasible primal point and feasible dual prices.

Show that their objective values are equal.

State why this proves both are optimal.



Exit ticket 4: when does the shadow price work?

Original LP: $b_1=4$, $b_2=5$; $z^*=9$ and $u=1$.

Keep $b_2=5$ and change only b_1.

A. Predict z^* at $b_1=4.2$ and check the production plan.

B. Would the same rule predict correctly at $b_1=6$?

Give the valid range and the actual optimum at 6.



Solution 1: combine a bound with an attainable edge

Step 1: $z=2(x+y)\le 8$ from $x+y\le 4$.

Step 2: attaining 8 requires $y=4-x$.

Step 3: $2x+(4-x)\le 5$ gives $x\le 1$.

With nonnegativity: $0\le x\le 1$.

The edge from (0,4) to (1,3) is optimal; $z^*=8$.



Solution 2: this dual candidate fails to give a bound

Step 1: (u,v)=(1,0) satisfies nonnegativity.

Step 2: u+2v=1<3 and u+v=1<2.

So the candidate is dual infeasible.

Step 3: w=4(1)+5(0)=4 is not a valid upper bound.

Counterexample: feasible (x,y)=(1,3) gives z=9>4.



Solution 3: verify both sides before equating values

Primal (x,y)=(1,3): resource use (4,5); x,y≥0.

Dual (u,v)=(1,1): u+2v=3; u+v=2; u,v≥0.

Primal value: z=3(1)+2(3)=9.

Dual value: w=4(1)+5(1)=9.

Weak duality gives z≤z*≤w. Hence z*=9.



Solution 4A: predict, then check at $b_1=4.2$

Step 1: $2.5 \le 4.2 \le 5$, so the old shadow price applies.

Step 2: $\Delta b_1=0.2$; $z^*=9+1(0.2)=9.2$.

Step 3: $x=5-4.2=0.8$; $y=2(4.2)-5=3.4$.

Check: $x+y=4.2$ and $2x+y=5$.

Direct value: $3(0.8)+2(3.4)=9.2$ = \$920/day.



Solution 4B: expose the failed extrapolation at 6

Old rule predicts $z=9+1(6-4)=11$.

But its basis gives $x=5-6=-1$, $y=7$: infeasible.

Actual optimum: $(x,y)=(0,5)$, $z^*=10$.

Dual $(u,v)=(0,2)$ is feasible and gives $w=6(0)+5(2)=10$.

Thus 10 is certified; the old price cannot cross $b_1=5$.



The big picture: one LP, four connected ideas



1. Primal proposes a plan; dual certifies its value

Primal: max 3x+2y; x+y≤4, 2x+y≤5; x,y≥0.

Dual: min 4u+5v; u+2v≥3, u+v≥2; u,v≥0.

Feasible plan (1,3) gives z=9.

Feasible prices (1,1) give w=9.

Weak duality: z≤z*≤w. Equality certifies optimality.



2. Complementary slackness closes the duality gap

$s_1 = 4 - x - y$; $s_2 = 5 - 2x - y$.

$r_x = u + 2v - 3$; $r_y = u + v - 2$.

$w - z = u \cdot s_1 + v \cdot s_2 + x \cdot r_x + y \cdot r_y$.

For feasible primal and dual points, all terms are nonnegative.

At optimum each product is zero: $u \cdot s_1 = v \cdot s_2 = x \cdot r_x = y \cdot r_y = 0$.

Two types of complementary pairs

Resource 1	$u \times s_1$	$1 \times 0 = 0$
Resource 2	$v \times s_2$	$1 \times 0 = 0$
Product x	$x \times r_x$	$1 \times 0 = 0$
Product y	$y \times r_y$	$3 \times 0 = 0$

Each term ≥ 0; zero gap makes every term zero.

3. Dual prices become local resource values

At capacities (4,5): (u,v)=(1,1).

Within the current basis region: $\Delta z^* = u\Delta b_1 + v\Delta b_2$.

One RHS at a time: $2.5 \le b_1 \le 5$; $4 \le b_2 \le 8$.

Example: $b_1 = 4.2$, $b_2 = 5$ gives $\Delta z^* = 0.2$.

This is \$20/day of gross gain, before any purchase cost.



4. When capacity changes, reconnect all four ideas



References and optional follow-up

Boyd and Vandenberghe, Convex Optimization: Sections 4.3, 5.2, 5.5, and 5.6.

SciPy documentation: linprog(method="highs"), marginals and residuals.

Optional Python practice: reproduce these examples with scipy.optimize.linprog.