
FÜKCH, L.:

COMPARING THE HOLOCENE AND THE RECENT MOLLUSK FAUNAS OF HUN-GARY - MAGYARORSZÁG HOLOCÉN ÉS RECENS MOLLUSCA-FAUNÁJÁNAK ÖSSZEHASONLITÁSA

ABSTRACT: Author compares the Holocene and recent faunas of present-day knowledge. The tabular illustration well shows that the two chronologically differentiated faunas have a similar combination in their main features. The existing differences arise from the continuous - natural or artificial - development of the fauna. Subsequent investigations on the faunas may reduce or enhance the identities or the differences between the two.

In a preceding paper /Fikch und KRCLOPP, 1985/ we have presented the list of Holocene faunas found in Hungarian collections. In the possession of these data we try to keep a breast with the fauna-development. An earlier paper of such a scope was restricted to the fauna-development of some areas of the country only /FÜKCH, 1982/. The papers which dealt with the spreading of the recent fauna of Hungary /PINTÉR, RICHNOVSZKY és SZIGNTHY, 1979; PINTÉR und SZIGETHY, 1979, 1980/ and which gave a taxonomic evaluation /PINTÉR, 1984/ of it were of great help.

The evolution and the spreading of our recent fauna is estimated to have taken place in the Quaternary. A considerable part of contemporary species can be found in Pleistocene deposits, too. The number of such elements which do not occour in our fauna, owing to extinction or the withdrawal of habitat, is very small. /The comparison of Pleistocene and recent faunas will be treated later./
In the course of the Holocene we have to reckon mostly with the influences of the climatic changes, anthropogenic activity which may result in habitat regrouping.



The malacological exploration of the Holocene deposits of our country is estimated to be moderate /Figure 1/. We have sufficient data from the areas mainly of the central range of mountains. The territory east of the river Tisza and the southern part of Transdanubia are white spots on our map. In spite of this explorative deficiency we can attempt to compare already the fauna of the two epochs /Holocene, recent/.

We have chosen the tabular illustration for the sake of a better understanding, in which the spesies appear in taxonomic grouping. The species which were found only in alluvial deposits are not figured in the table. In the "note" we refer to the possible reasons of the divergence of the two

feuna-lists. We do not consider data of the table as final, we collected them in the function of the actual time cross-section. The present Holocene fauna researches and the natural and/or artificial development of the recent fauna can approach or abduct the two fauna-lists.

The following may be established: The combination of the Holocene and recent fauna shows a remarkable conformity in the main features. At the same time we can point out that the development of our fauna is continuous in the present days, too, - as a result of anthropogenic activity /introductions, e. g. Alopia species/, and, on the other hand, by means of the natural transmission along the rivers /e. g. Helicigona banatica/. At the same time we have to reckon with a progressive speading of the species which today show isolated spreading picture.

## ÖSSZEFOGLALÁS

A dolgozat része annak a sorozatnak, mely a magyarországi csiga-fauna fejlődését mutatja be a harmadidőszaktól napjainkig. A korábbi dolgozatok /KROLOPP, 1983; KROLOPP, 1984; PÜKÖH und KROLOPP, 1985/ ismertették a pleisztocén és holocén faunát. Jelen dolgozat a faunisztika szintjén mutatja be a holocén és recens faunák közötti kapcsolatokat, az azonosságokat és különbségeket. A két faunalista egybevetése rámutat a jelenleg is folyó faunafejlődésre, mely egyrészt természetes folyamat, másrészt mesterséges tevékenység eredménye.

Family
Holocene Species
Neritidae
Theodoxus transversalis
Theodoxus prevostianus

Viviparidae Viviparus acerosus Viviparus contectus Recent Species

Theodoxus transversalis
Theoduxus prevostianus
Theoduxus fluviatilis 1.
Theoduxus danubialis 2.

Viviparus acerosus Viviparus contectus Valvatidae Valvata naticina Valvata piscinalis Valvata cristata Valvata pulchella

Pomatiasidae Pomatias elegans

Hydrobiidae Lithoglyphus naticoides Sadleriana pannonica

Bithyniidae Bithynia leachi Bithynia tentaculata

Aciculidae Acicula polita

Melanopsidae Fagotia acicularis Fagotia esperi

Ellobiidae Carychium minimum Carychium tridentatum

Acroloxidae Acroloxus lacustris

Lymnaeidae
Lymnaea stagnalis
Lymnaea palustris
Lymnaea truncatula
Lymnaea auricularia
Lymnaea peregra

Physidae Aclexa hypnorum Physa fontinalis Valvata naticina Valvata piscinalis Valvata cristata Valvata pulchella

Pomatias elegans Pomatias rivulere 3.

Lithoglyphus naticoides
Sadleriana pannonica
Paladilhia hungarica 4.
Bythinella austriaca 5.
Potamopyrgus jenkinsi 6.

Bithynia leachi Bithynia tentaculata

Acicula polita Acicula banatica 7.

Fagotia acicularis
Ragotia esperi
Melanopsis parreyssi 6.
Melanoides tuberculata 6.
Amphimelania holandri 8.

Carychium minimum Carychium tridentatum

Acroloxus lacustris

Lymnaea stagnalis
Lymnaea palustris
Lymnaea truncatula
Lymnaea auricularia
Lymnaea peregra
Lymnaea columella 6.

Aplexa hypnorum
Physa fontinalis
Physella heterostropha 9.
Physella acuta

Planorbidae
Planorbis corneus
Planorbis planorbis
Planorbis carinatus
Anisus septemgyratus
Anisus leucostoma
Anisus spirorbis
Anisus vortex
Anisus vortex
Anisus vorticulus
Bathyomphalus contortus
Gyraulus albus
Gyraulus crista
Hippeutis complanatus
Segmentina nitida

Ancylidae Ancylus fluviatilis

Cochlicopidae
Cochlicopa lubrice
Cochlicopa lubricella
Cochlicopa nitens

Pyramidulidae Pyramidula rupestris

Vertiginidae
Columella edentula
Truncatellina cylindrica
Truncatellina claustralis
Vertigo angustior
Vertigo pusilla
Vertigo antivertigo
Vertigo moulinsiana
Vertigo pygmaea
Vertigo alpestris

Orculidae
Orcula dolium
Fohyradium doliolum
Pagodulina pagodula

Chondrinidae Granaria frumentum Chrondrina clienta

Pupillidae
Pupilla muscorum
Pupilla triplicata

Planorbarius corneus
Planorbis planorbis
Planorbis carinatus
Anisus septemgyratus
Anisus leucostoma
Anisus soirorbis
Anisus vortex
Anisus vorticulus
Bathyomphalus contortus
Gyraulus albus
Gyraulus laevis
Gyraulus crista
Hippeutis complanatus
Eegmentina nitida
Helisoma duryi

Ancylus fluviatilis Ferrissia wautieri lo.

Cochlicopa lubrica Cochlicopa lubricella Cochlicopa nitens

Pyramidula rupestris

Columella edentula
Truncatellina cylindrica
Truncatellina claustralis
Vertigo angustior
Vertigo pisulla
Vertigo moulinsiana
Vertigo mygmaea
Vertigo alpestris
Truncatellina callicratis 11.
Vertigo substriata 10.

Orcula dolium Sphyradium doliolum Pagodulina pagodula

Cranaria frumentum Chondrina clienta

Pupilla muscorum Pupilla triplicata Spelaeodiscus triarius 12. Valloniidae
Vallonia pulchella
Vallonia costata
Vallonia enniensis
Achantinula aculeata

Enidae Chondrula tridens Ena montana Ena obscura Zebrina detrita

Succineidae
Succinea putris
Succinea oblonge
Oxyloma elegans

Clausiliidae
Cochlodina cerata
Cochlodina laminata
Cochlodina orthostoma
Ruthenica filograna
Macrogastra ventricosa
Macrogastra plicatula
Macrogastra latestriata
Clausilia dubia
Clausilia pumila
Clausilia cruciata
Laciniaria plicata
Balea biplicata
Bulgarica cana

Ferussaciidae Cecilioides acicula

Endodontidae
Punctum pygmaeum
Discus rotundatus
Discus perspectivus
Discus ruderatus

Vitrinidae Vitrina pellucida Vallonia pulchella Vallonia costata Vallonia enniensis Achantinula aculeata

Chondrula tridens
Ena montama
Ena obscura
Zebrina detrita

Succinea putris Succinea oblonga Oxyloma elegans

Cochlodina cerata Cochlodina laminata Cochlodina orthostoma Ruthenica filograna Macrogastra ventricosa Macrogastra plicatula Macrogastra latestriata Clausilia dubia Clausilia pumila Clausilia cruciata Laciniaria plicata Balea biplicata Bulgarica cana Alopia straminicollis monacha 6. Alopia livida bipalatalis 6. Pseudofusulus varians Clausilia parvula 13. Balea perversa lo. Vestia gulo 14. Vestia turgida 15. Bulgarica rugicollis 6.

Cecilioides acicula Cecilioides petitiana

Punctum pygmaeum
Discus rotundatus
Discus perspectivus
Discus ruderatus
Helicodiscus singleyanus

Vitrina pellucida <u>Semilimax semilimax</u> 13. <u>Phenacolimax annularis</u> 12. Zonitidae
Zonitoides nitidus
Vitrea diaphana
Vitrea subrimata
Vitrea crystallina
Vitrea contracta
Aegopinella minor
Aegopinella pura
Aegopinella ressmani
Aegopinella ressmani
Aegopis verticillus
Nesovitrea hammonis
Oxychilus orientalis
Oxychilus depressus
Cxychilus depressus
Daudebardia rufa
Daudebardia brevipes
Daudebardia helenae

Euconulidae Euconulus fulvus

Bradybaenidae Bradybaena fruticum

Helicidae
Helicella obvia
Helicopsis striata
Monacha cartusiana
Perforatella bidentata
Perforatella incarnata
Perforatella vicina
Perforatella rubiginosa
Trichia unidentata
Trichia unidentata
Trichia hispida
Euomphalia strigella
Helicigona faustina
Helicigona arbustorum
Isognomostoma isognomostoma
Cepaea vindobonensis
Cepaea hortensis
Helix pomatia

Zonitoides nitidus
Vitrea diaphana
Vitrea subrimata
Vitrea crystallina
Vitrea contracta
Aegopinella minor
Aegopinella pura
Aegopinella ressmani
Aegopis verticillus
Nesovitrea hammonis
Oxychilus orientalis
Oxychilus glaber
Oxychilus inovinatus
Oxychilus depressus
Daudebardia rufa
Daudebardia brevipes

Zonitoides arboreus 6. Oxychilus translucidus 6. Oxychilus dranarnaudi Oxychilus hydatinus 16.

Euconulus fulvus

Bradybaena fruticum

Helicella obvia Helicopsis striata Monacha cartusiana Perforatella bidentata Perforatella incarnata Perforatella vicina Perforatella rubiginosa Trichia unidentata Trichia hispida Euomphalia hispida Helicodonta obvoluta Helicigona faustina Helicigona arbustorum Isognomostoma isognomostoma Cepaea vindobonensis Cepaea nemoralis Cepaea hortensis Helix pomatia Candidula unifasciata 6. Perforatella dibothrion 14. Perforatella umbrosa 13. Hygromia cinctella 6. Hygromia transsylvanica 17. Hygromia kovacsi 18.

Trichia striolata 19.
Trichia filicina 13.
Trichia erjaveci 13.
Trichia lubomirskii 20.
Helicigona banatica 21.
Helicigona planospira 22.
Eobania vermiculata 6.
Helix aspersa 6.
Helix lutescens 10.

Localites of the Holocene Paunas of Hungary

1. In the Tisza 2. Eastward from the Danube only in the I-poly 3. Fen-wood of Bátorliget 4. In the caves of the Mecsek 5. In the springs of the Northern-Central Range of Mountains 6. Introduced 7. In the Mecsek 8. Westward from the lake Balaton 9. In Délegyháza lo, Insular-like spreading 11. Westward from the Danube 12. In the Bükk. 13. In Transdanubia 14. In Zemplén 15. Bükk, Karst of Aggtelek 16. In Békéscsaba 17. In Bükk and Mátra 18. In the county Békés 19. Along the Danube 20. Mátra and Zemplén 21. Near to Vásárosnamény 22. Along the Dráva

## REFERENCES

FÜKCH, L. /1982/: A holocén faunakutatás jelentősége a recens fauna feltárásában. Mal. Táj., 2: 12-13. - FÜKÜH, L. und KROLOPP, E. /1985/: Verzeichnis der holozänen Mollusken Ungarns. Soosiana, 13: 145-146. - KROLOPP, E. /1983/: Verzeichnis der pleistozinen Mollusken Ungarns. 10/11: 75-78. -KROLOPP, E. /1984/: A magyarországi pleisztocén Mollusca-fauna/jellemvonásai. Coosiana, 12: 7-10. - PINTÉR, L., RICHNOV-SZKY, A. és S. SZIGETHY, A. /1979/: A magyarországi recens puhatestuek elterjedése. Soosiana, /Suppl.I./: 1-351. -PINTÉR, L. und F. SZIGETHY, A. /1979/: Die Verbreitung der rezenten Mollusken Ungarns: Neunachweise und Berichtigungen, I. Soosiana, 7: 97-108. - PINTÉR, L. und S. SZICETHY, A. /1980/: Die Verbreitung der rezenten Mollusken Ungarns: Neunachweise und Berichtigungen, II. Soosiana, 8: 65-80. -PINTÉR, L. /1984/: Magyarország recens puhatestüinek revidealt katalógusa. /Mollusca/. Pol.Hist-nat.Mus.Matr.,9: 79-90.

DR. FÜKÜH LEVENTE Mátra Nuzeum Gyöngyös Kossuth u. 40. H-3200