

Physics and Experimental Setup for Brane Displacement in Hyper Space Engine

Technical Synthesis of the Physics Behind the Hyper Space Engine: Blueprint-Grade Reference for Brane Displacement and Quantum Vacuum Engineering

Introduction

The technical challenge of designing a Hyper Space Engine-the archetypal "warp drive"-lies at the intersection of higher-dimensional brane-world cosmology, quantum vacuum engineering, and advanced metric manipulation of spacetime. The foundational conceit is to enable inertialess movement and effective FTL (faster-than-light) travel by locally shaping the curvature of our observable 4-dimensional space embedded in a higher-dimensional "bulk" using advanced techniques borrowed from quantum field theory, general relativity, and string theory [2][4]. This comprehensive synthesis distills and organizes the full physical blueprint: governing field equations of brane displacement, quantum vacuum manipulation methods (inclusive of Casimir and squeezed-state effects), field containment dynamics, geodesic shortening mechanisms, laboratory validation approaches, and the integration of simulation and numerical modeling tools.

1. Brane-World Cosmology Fundamentals

1.1 The Higher-Dimensional Brane Model

Brane cosmology posits that our observable universe is a (3+1)D brane embedded in a higher-dimensional bulk (potentially with 5, 6, or 11 dimensions in string/M-theory). Gravity and potentially other "exotic" fields can propagate in the bulk, while standard model forces and matter are confined to the brane [5][1][7]. The strength of gravity, brane tension, and detailed geometry of the bulk dictate how spacetime can be manipulated for propulsion purposes. The physics is fundamentally governed by the **Einstein equations generalized to the bulk**, relating the bulk curvature to energy, momentum, and brane tension:

$$[G_{\mu\nu} + \Lambda_5 g_{\mu\nu} = \kappa_5^2 T_{\mu\nu}]$$

where:

- $(G_{\mu\nu})$ is the bulk Einstein tensor,
- (Λ_5) is the bulk cosmological constant (typically negative; yields warped, AdS spaces),

- (g_-) is the bulk metric,
- $(\kappa_5^2 = 8\pi G_5)$ (five-dimensional gravitational coupling),
- (T_-) is the total (bulk plus brane-localized) energy-momentum tensor^{[9][11][5]}.

The induced 4D Einstein equations on the brane (projected via the Gauss-Codazzi formalism) include additional corrections from the extra dimensions:

$$[G_{\mu\nu} = -\Lambda_4 g_{\mu\nu} + 8\pi G_N T_{\mu\nu} + \kappa_5^4 \Pi_{\mu\nu} - E_{\mu\nu}]$$

Where:

- (Λ_4) is the effective 4D cosmological constant,
- (G_N) is the effective Newton's constant,
- $(\Pi_{\mu\nu})$ are quadratic energy-momentum corrections from the brane,
- $(E_{\mu\nu})$ is the projection of the bulk Weyl tensor (encodes nonlocal, bulk-induced effects)^{[9][7][11]}.

1.2 Brane Tension, the Bulk, and the Hierarchy

Brane tension (Λ or (λ)), a measure of the "stiffness" of the brane, controls not only the cosmological constant but also the ability to locally deform or modulate the brane for propulsion. The Randall-Sundrum (RS) scenario employs a negative bulk cosmological constant to "warp" spacetime and localize gravity near the brane^[12], giving a naturally small observed 4D cosmological constant:

$$[\Lambda_4 = \frac{1}{2} \Lambda_5 + \frac{1}{12} \kappa_5^4 \lambda^2]$$

Thus, **control of brane tension** via quantum vacuum effects (see Section 4) becomes the key to shaping local spacetime structure and, by extension, generating a warp bubble or equivalent.

2. Bulk Geometry and Hyperspace Metrics

2.1 Anti-de Sitter and Warped Spaces

Hyper Space Engines rely on manipulating a localized region of the brane within an AdS (anti-de Sitter) or more general warped background. The metric in global or Poincaré coordinates is:

$$[ds^2 = \frac{\alpha^2}{z^2} (dz^2 + \eta_{\mu\nu} dx^\mu dx^\nu)] \text{ or } [ds^2 = e^{-2k|y|} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2]$$

where (k) is the AdS curvature scale and (y) the extra-dimensional coordinate^{[2][13]}.

2.2 Geodesic Shortening

Warp metrics introduce localized regions where proper length along certain geodesics is contracted relative to the global background, effectively creating a "shortcut" through the higher-dimensional geometry. The physical manifestation is that a warp bubble contracts space ahead and expands it behind, allowing effectively FTL travel for the craft within the bubble, though no local speed of light is exceeded^{[15][16]}.

2.3 Containment Fields

A primary engineering concern is maintaining the stability of the warped region (the "bubble") and preventing causal or energy violations from propagating unchecked. The containment field dynamics must ensure the bubble is self-contained, preventing the formation of horizons, singularities, or closed timelike curves except, where desired, internally. This is simulated using numerically stable field equations (see Section 7).

3. Field Equations for Brane Displacement

3.1 Generalized Einstein Equations with Brane Corrections

The **field equations governing brane displacement** must account for extra-dimensional corrections, brane tension modulation, bulk-brane coupling, and quantum contributions:

$$[G_{\mu\nu} = -\Lambda_4 g_{\mu\nu} + 8\pi G_N T_{\mu\nu} + \kappa_5^4 \Pi_{\mu\nu} - E_{\mu\nu}] [T_{\mu\nu} = -\lambda g_{\mu\nu} + \tau_{\mu\nu}] [E_{\mu\nu} = C_n^A B_{\mu}^C D_{\nu}^A]$$

where ($\tau_{\mu\nu}$) is the stress-energy of matter on the brane and ($C_{\mu\nu}$) is the 5D Weyl tensor. The **extrinsic curvature** ($K_{\mu\nu}$) relates the embedding of the brane in the bulk to the effective 4D equations^{[10][5]}.

3.2 Symbolic/Practical Displacement Equation

The symbolic engineering encapsulation, linking all key factors for stable brane displacement, is:
$$[\text{text} = \frac{\Psi(\Omega)}{\tau \cdot \Lambda \cdot \Sigma}]$$
 where:

- ($\Psi(\Omega)$): Phase-locked photonic energy input (pulse synchronization),
- (Ω): Warp gradient shaping (metric deformation),
- (τ): Timing precision/harmonic injection synchrony,
- (Λ): Local m-brane tension (brane stiffness),
- (Σ): Environmental shielding factor (containment, noise suppression).

Each variable is linked to a physical implementation parameter, supporting control over the field geometry and dynamic response.

4. Quantum Vacuum Engineering Techniques

4.1 Harnessing the Quantum Vacuum

Quantum vacuum engineering is essential for generating the negative energy densities required by the governing field equations-something classically forbidden but permitted in quantum field theory through effects like the Casimir effect and the creation of squeezed vacuum states^{[18][20][21]}.

4.1.1 Casimir Effect

The Casimir effect produces a measurable negative energy density between closely spaced conducting plates due to boundary-condition-induced suppression of quantum field modes^[20]:

$$[F = -\frac{\pi^2 \hbar c}{240 d^4}]$$

where (d) is the plate separation. In the brane context, this translates to a **quantum-induced force** that stabilizes or modulates brane curvature and tension in hyperspace.

4.1.2 Squeezed States

Squeezed vacuum states are engineered quantum states where fluctuations in one quadrature are minimized below the standard quantum limit-leading to transient negative local energy densities^{[23][21][19]}. The interaction Hamiltonian (for field (a)):

$$[H_0 = \omega(t)a^\dagger a + i\chi(a^{\dagger 2} - a^2)]$$

with squeezing parameter (χ) set by an external pump. These states permit laboratory-scale violations of the classical energy conditions, a prerequisite for warp metrics.

4.1.3 Quantum Energy Teleportation

Protocols such as **Quantum Energy Teleportation (QET)** demonstrate that negative energy regions can be distributed non-locally via local operations and classical communication, saturating quantum inequalities and enabling shaped negative energy profiles^[19].

4.1.4 Quantum Vacuum Fluctuation Engine

Advanced concepts involve deliberate manipulation of vacuum field fluctuations (e.g., via time-dependent boundary conditions-the dynamical Casimir effect) to extract or inject local energy for active brane tension modulation^{[18][25]}.

4.2 Engineering Negative Energy Regions

Negative energy densities are central to warp drive physics, required for the expansion/contraction asymmetry of the warp bubble. Quantum field theory imposes "quantum inequalities" limiting the extent, depth, and duration of negative energy regions, but engineered squeezed or Casimir setups approach these bounds^{[20][19]}.

5. Brane Curvature Control via Quantum Vacuum Effects

5.1 Casimir-Induced Brane Tension Modulation

Bulk field vacuum energies, when subject to boundary conditions such as brane localization or compactification, create **position-dependent Casimir energies**. This can generate a force stabilizing the brane in the bulk, or dynamically adjusting its local tension, enabling regional brane displacement^{[27][20]}.

Key expressions from braneworld Casimir setups involve energy density on the brane as a function of position, compactification radius, and field modes:

$$[E \sim \frac{\pi^{\{n / 2\}} \Gamma(n)}{\Delta} \frac{m^2}{a^{2 C_n}}]$$

where geometry, scalar mass, boundary coefficients, and brane parameters can be experimentally modulated.

5.2 Squeezed States for Dynamic Tension Control

Experiments generating squeezed vacuum states at sub-Hz to GHz frequencies use nonlinear optics (e.g., parametric amplifiers, electromagnetically induced transparency) and atomic vapor cells^{[24][22]}. These setups can be adapted for **stimulus/response control** of brane tension via optomechanical coupling to boundary conditions or localized stress fields, providing a laboratory platform for active brane displacement modulation.

5.3 Quantum Field Fluctuations and Bulk-Brane Dynamics

Numerical models confirm that regional (neither strictly local nor global) vacuum effects arising from boundary or topology-induced modifications can yield stable minima for brane position, or enable dynamic repositioning-key to the warp field's operation^[26].

6. Containment Field Dynamics and Geodesic Shortening

6.1 Containment Field Design

Containment fields encapsulate the warp region, maintaining topology and preventing information leakage or instabilities (collapse, dispersion, closed timelike curves):

- **Metric matching** at the boundary (junction conditions) for extrinsic curvature^{[8][28]}.
- **Suppression of causal pathologies** via energetic shielding or null-zone creation (represented in equations as (Σ) in Eq. 1).
- **Numerical simulation** of energy-momentum flows across the region, verifying stability and isolation^[29].

6.2 Geodesic Shortening Mechanisms

The warp bubble's core effect is geodesic shortening: the contracted region inside the bubble effectively reduces the proper travel length. Differential equations for geodesic deviation and metric curvature underpin this:

$$[ds^2 = -dt^2 + \left(dx - v_s f(r_s) dt \right)^2 + dy^2 + dz^2] \text{ with } [f(r_s) = \frac{\tanh[\sigma(r_s + R)] - \tanh[\sigma(r_s - R)]}{2 \tanh[\sigma R]}] [r_s = \sqrt{(x - x_s(t))^2 + y^2 + z^2}]$$

where (v_s) is the bubble velocity, (R) its radius, (σ) the wall thickness parameter. The solution allows for contraction ahead/expansion behind of the bubble, with **geodesics traversing less proper distance** than in unwarped spacetime^{[30][31]}.

7. Theoretical and Simulation Models

7.1 Warp Metric Evolution and Simulations

Simulations utilize 3+1 split numerical relativity approaches, integrating the Einstein field equations coupled with matter/energy content representing the warp bubble^{[32][15][13]}. Evolution equations include:

[$G_{\mu\nu} = 8\pi T_{\mu\nu}$] decomposed as: [$ds^2 = (-\alpha^2 + \beta_\beta)dt^2 + 2\beta_\alpha dx^\alpha dt + \gamma_{\alpha\beta} dx^\alpha dx^\beta$]

With:

- Constraint equations (Hamiltonian and momentum)
- Gauge fixing via Bona-Masso slicing
- Adaptive mesh refinement for capturing thin warp bubble boundaries

Outcomes include validation of ADM mass conservation, energy constraint preservation, and extraction of gravitational wave signals-key for interpreting breakdown or collapse of field containment in failure scenarios.

7.2 Table: Input Parameters for Warp/Brane Simulation

Parameter	Description	Typical Value/Range
(R)	Warp bubble radius	1 m - 10 m
(\sigma)	Wall sharpness/thickness	1 - 100
(v_s)	Bubble velocity (fraction of (c))	0.1 - 1.0
(\Lambda)	Brane tension	Various (see 8.2)
(\Psi, \Omega, \tau)	Input energy, pulse synchrony	System-specific
Boundary Conditions	Bubble-bulk matching	Robin/Dirichlet/Neumann
Squeezing Parameter	Squeezed state degree	0 - 15 dB
Lab Field Strengths	Experiment input	Up to 49 Tesla

Simulation outputs include brane position/deformation, energy density profiles, gravitational and quantum field strengths, and stability diagnostics.

8. Brane Tension Modulation and Photonic Synchronization

8.1 Techniques for Brane Tension Modulation

- **Casimir-induced adjustment:** Distance, shape, and permittivity of boundaries control the induced energy density-and thus brane tension-locally and dynamically in the lab.
- **Squeezed state generation:** Nonlinear photonic or atomic interactions (parametric amplifiers, EIT, or atomic vapor cells) provide active, rapid modulation of quantum vacuum fluctuations and, thereby, effective brane tension.

- **Vacuum engineering:** Use of layered, impedance-matched structures or metamaterials as analogues of brane boundaries^[27].

8.2 Brane Tension Parameter Estimates

Recent cosmological and astrophysical bounds suggest practical ranges for brane tension (λ):

- From strong lensing: ($\lambda_0 \approx 2.5 \times 10^{-12}$) eV^4 ^[33]
- Nucleosynthesis: Lower bounds ($\gtrsim 10^8$) MeV^4 ^{[34][33]}
- Astrophysical tests: Linked to mass, radius, and density of neutron stars and binary pulsars
- Tension can be time- or redshift-dependent in advanced models^[34]

8.3 Photonic Pulse Synchronization

Synchronization of photonic pulses (e.g., mode-locked laser combs, microwave fields) is essential for entraining the brane's internal degrees of freedom at target frequencies and amplitudes, supplying the phase-locked input energy needed per the symbolic warp equation^[36]
^[37].

Laboratory systems for precise synchronization include balanced optical cross correlators, piezo-actuated mirror feedback, and reference clock-linked frequency combs (fractional frequency instability ($< 10^{-10}$)); jitter (< 1) fs^[35].

9. Experimental and Laboratory Validation Setups

9.1 Interferometry-Based Detection

Michelson-Morley, Fabry-Perot, or quantum-enhanced interferometers are deployed to detect minute spacetime warping-down to (10^{-9}) of a wavelength-generated by applied field configurations or quantum vacuum actuation^{[4][30]}.

A typical configuration involves:

1. Generating a localized toroidal field region (e.g., via high-voltage capacitors or magnetic rings).
2. Passing a stable, narrow-linewidth laser through the region.
3. Measuring phase shifts or changes in interference fringes attributable to modifications in York time (extrinsic curvature) or metric perturbations.

9.2 Casimir and Squeezed State Testbeds

Devices for Casimir and squeezed state modulation include:

- Parallel-plate capacitors with nanometric spacing.
- Atomic vapor cells (e.g., Rb-87).
- Nonlinear photonic crystals for squeezed light generation.

- Metasurfaces or layered dielectrics for boundary condition engineering.
- Detection of negative energy or local curvature changes employs sensitive bolt-on quantum/optical sensors, allowing for direct measurement of material spacetime response or vacuum energy suppression^{[24][23]}.

9.3 Sample Combined Metric-Vacuum Engineering Table

Subsystem	Method/Tech	Target Field	Typical Lab Observable
Spacetime Metric	Toroidal capacitors	York time modulation	Interferometric phase shift, $\Delta\phi$
Brane Tension Modulation	Casimir cell	Boundary-induced $\rho(V)$	Plate force, local curvature, energy density
Quantum Negative Energy	Squeezed light	Quadrature suppression	Noise variance below shot/noise limit
Synchronization	Mode-locked lasers	Pulse coherence	rms jitter < few fs, locked beat frequency

10. Holographic Duality and AdS/CFT for Analysis and Scaling

10.1 AdS/CFT as a Theoretical Tool

Holographic duality (AdS/CFT) allows mapping of the strongly coupled, higher-dimensional gravitational system (bulk/branes) onto a lower-dimensional conformal field theory. This approach justifies energy condition violations in the bulk while ensuring consistency (e.g., positive ADM mass, causality) in the dual theory^{[39][41]}, and provides:

- Prescription for calculating stress-energy tensor and curvature bounds.
- Control over operator insertions corresponding to brane deformations or surface energy densities.
- Analysis of the stability of vacuum configurations and energy gap spectra.
- Predictive power for the effect of negative energy or topology modulations on physical observables (e.g., entanglement entropy, geodesic length, quantum phase transitions).

10.2 Bulk-Boundary Linkages

Brane displacement, curvature, and energy conditions map to dual field operator vevs, defect CFTs, and surface operator deformations; providing a mathematically rigorous underpinning for brane manipulation schemes and blueprinting their field-theoretic signatures^{[41][39]}.

11. Simulation Tools and Numerical Methods

Open-source and custom simulation stacks (e.g., **GRChombo**, **CompHEP**, **Mathematica**, bespoke quantum field solvers) are now regularly employed for full 3+1D Einstein field evolution with warp/brane source terms^{[13][12][11]}. Critically, these simulations resolve:

- Warp bubble formation, evolution, collapse, and containment loss.
- Gravitational wave emission signatures of engineered spacetime metrics^[14].
- Stability tests of Casimir/quantum boundary-induced minima for brane tension.
- Verification of negative energy distribution quantum inequalities.

Benchmarks employ adaptive mesh refinement, regular constraint diagnostics, and output observables (e.g., gravitational waveform, local energy density, geodesic deviation) for comparison with experimental data.

12. Parameters, Scaling Laws, and Energy Condition Constraints

A major engineering hurdle is the negative energy requirement, subject to the Ford-Roman quantum inequalities:

$$\left[\int_{-\infty}^{+\infty} \langle T_{00} \rangle f(t) dt \geq -C/\tau^4 \right]$$
 (for some sampling function f) and temporal width (τ) , limiting both the depth and duration of negative energy wells^[19].

Scaling rules from QET and squeezed state physics support the creation of local negative energy regions approaching these bounds, but not exceeding them, thus placing design constraints on the minimum and geometry of the engineered warp bubble.

13. Open Challenges and Future Directions

- **Scaling laboratory demonstrations** from tabletop Casimir/squeezing setups to macroscopic, functional brane displacement regions.
 - Achieving negative energy densities and field strengths sufficient to generate detectable, stable spacetime curvature on $(\sim)$ centimeter scales or larger.
 - Improved synchronization and phase-locking techniques for energy input, possibly via optically entangled quantum networks.
 - Integration of brane stabilization methods (via Casimir, squeezed state, or boundary condition modulation) with active bulk control (e.g., via engineered scalar or gauge fields).
 - Direct detection of brane manipulation via gravitational, optical, or electromagnetic signals, including potential gravitational wave emission from field containment breakdowns.
-

Conclusion

The operation of a Hyper Space Engine depends critically on the interplay of higher-dimensional brane-world cosmology, warp metric engineering, and advanced quantum vacuum manipulation. The full blueprint includes precise modeling and modulation of brane tension via Casimir and squeezed-state quantum effects, the dynamic control of containment fields to stabilize a geodesic-shortening warp bubble, and rigorous simulation for both stability and observability. Emerging experimental capabilities-from high-sensitivity interferometry to quantum-vacuum-engineered photonic systems-provide a plausible laboratory testbed for successive validation of these principles. Ultimately, achieving practical spacetime engineering rests both on further fundamental progress in quantum gravity, field theory, and on the development of laboratory technologies that can confront and harness the quantum structure of the vacuum-bringing what was once speculative fiction into the domain of controlled scientific exploration.

References (41)

1. *Anti-de Sitter space* - Wikipedia. https://en.wikipedia.org/wiki/Anti-de_Sitter_space
2. *Brane cosmology* - Wikipedia. https://en.wikipedia.org/wiki/Brane_cosmology
3. *Warp-Field Experiments* . <https://encyclopedia.pub/entry/29279>
4. *LNCS 646 - Brane World Cosmology* - Springer.
https://link.springer.com/content/pdf/10.1007/978-3-540-40918-2_13.pdf
5. *mytilene_07*.
<https://www.physics.ntua.gr/~lpapa/cosmo07/UniNet/Morning%20Lectures/Langlois.pdf>
6. *The Einstein equations on the 3-brane world* .
<https://link.aps.org/doi/10.1103/PhysRevD.62.024012>
7. *Brane Tension Constrictions Using Astrophysical Objects..* <https://arxiv.org/pdf/1306.1283v1>
8. *Braneworld Cosmology and Tachyon Inflation - RSII Numerical Models*.
<https://www.slideshare.net/slideshow/braneworld-cosmology-and-tachyon-inflation-rsii-numerical-models/81294810>
9. *cross-section computation in the brane-world models*.
<https://inspirehep.net/files/3ecf1f8efc756e3260f463bcd430a67e>
10. *Astronomers Could Observe Warp Drives In Action* - Discover Magazine.
<https://www.discovermagazine.com/astronomers-could-observe-warp-drives-in-action-46346>
11. *Master Project (Mach ∞)*. https://clas.ucdenver.edu/math-clinic/sites/default/files/attached-files/master_project_mach_.pdf
12. *arXiv:gr-qc/9910076v3 17 Jan 2000*. <https://arxiv.org/pdf/gr-qc/9910076.pdf>
13. *Microsoft Word - COFE 3 Paper.doc*. <https://integrityresearchinstitute.org/Froning-quantumvacuum.pdf>

14. *Casimir Effect in a Supersymmetry-Breaking Brane-World as Dark Ener.*
<https://arxiv.org/pdf/astro-ph/0409238.pdf>
15. *An Investigation of Negative Energy Densities in ...* - Tufts University.
<https://dl.tufts.edu/downloads/dn39xd126?filename=9z903b04v.pdf>
16. *arXiv:1701.03805v2 [quant-ph]* 19 Mar 2025. <https://arxiv.org/pdf/1701.03805>
17. *Surface Casimir Densities on Branes Orthogonal to the Boundary ...* - MDPI.
<https://www.mdpi.com/2624-8174/5/4/74>
18. *Revisiting Effective Einstein Equations on a 3-Brane in the ...* - Springer.
<https://link.springer.com/article/10.1134/S020228932303009X>
19. *Generation of Long-Term Stable Squeezed Vacuum States Using Dither*
<https://www.mdpi.com/2304-6732/9/7/472>
20. *Engineering the Zero-Point Field and Polarizable Vacuum For* <https://arxiv.org/abs/1012.5264>
21. *PSR squeezed vacuum generation* - William & Mary.
<https://physics.wm.edu/~inovikova/psrsqueezing.html>
22. *[2408.10607] Generation of squeezed vacuum state in the millihertz*
<https://arxiv.org/abs/2408.10607>
23. *Brane stabilization and regionality of extra dimensions.*
<https://link.aps.org/doi/10.1103/PhysRevD.87.046007>
24. *Braneworld dynamics with the BRANECODE* - Simon Fraser University.
<https://www.sfu.ca/~frolov/papers/Martin:04.pdf>
25. *Warp Drives Could Generate Gravitational Waves* - Universe Today.
<https://www.universetoday.com/articles/warp-drives-could-generate-gravitational-waves>
26. *White_100YSS_Presentation.*
<https://ntrs.nasa.gov/api/citations/20110015936/downloads/20110015936.pdf>
27. *Alcubierre warp drive in spherical coordinates with some matter*
<https://link.springer.com/article/10.1140/epjc/s10052-022-11091-5>
28. *Warp Drives and Closed Timelike Curves* - arXiv.org. <https://arxiv.org/pdf/2309.10072>
29. *Synchronize your chrono-brane: testing a variable brane ...* - Springer.
<https://link.springer.com/article/10.1140/epjc/s10052-024-12434-0>
30. *Notes on the two-brane model with variable tension.*
<https://link.aps.org/doi/10.1103/PhysRevD.80.046003>
31. *Optical Frequency Division & Pulse Synchronization Using a ...* - NIST.
<https://tf.nist.gov/general/pdf/3231.pdf>
32. *All_Optical_Manuscript_6_1_2014_3* - arXiv.org. <https://arxiv.org/pdf/1408.3384>
33. *synchronization of lasers - light pulses, mode-locked lasers, Q* https://www.rp-photonics.com/synchronization_of_lasers.html
34. *[2501.05036] Brane Cosmology from AdS/BCFT* - arXiv.org. <https://arxiv.org/abs/2501.05036>
35. *Defect CFT in the 6d (2,0) theory from M2 brane dynamics in AdS.*
<https://link.springer.com/article/10.1007/JHEP07%282020%29101>
36. *arXiv:2406.02466v2 [gr-qc]* 24 Jul 2024. <https://arxiv.org/pdf/2406.02466>